


## 5. Agriculture

**A**gricultural activities contribute directly to emissions of greenhouse gases through a variety of processes. This chapter provides an assessment of non-carbon dioxide emissions from the following source categories: enteric fermentation in domestic livestock, livestock manure management, rice cultivation, agricultural soil management, and agricultural residue burning (see Figure 5-1). Carbon dioxide (CO<sub>2</sub>) emissions and removals from agriculture-related land-use activities, such as conversion of grassland to cultivated land, are discussed in the Land-Use Change and Forestry chapter.

In 2000, agricultural activities were responsible for emissions of 485.1 Tg CO<sub>2</sub> Eq., or 6.9 percent of total U.S. greenhouse gas emissions. Methane (CH<sub>4</sub>) and nitrous oxide (N<sub>2</sub>O) were the primary greenhouse gases emitted by agricultural activities. Methane emissions from enteric fermentation and manure management represent about 20 percent and 6 percent of total CH<sub>4</sub> emissions from anthropogenic activities, respectively. Of all domestic animal types, beef and dairy cattle were by far the largest emitters of methane. Rice cultivation and agricultural crop residue burning were minor sources of methane. Agricultural soil management activities such as fertilizer application and other cropping practices were the largest source of U.S. N<sub>2</sub>O emissions, accounting for 70 percent. Manure management and agricultural residue burning were also small sources of N<sub>2</sub>O emissions.

Table 5-1 and Table 5-2 present emission estimates for the Agriculture chapter. Between 1990 and 2000, CH<sub>4</sub> emissions from agricultural activities increased by 2.9 percent while N<sub>2</sub>O emissions increased by 11.3 percent. In addition to CH<sub>4</sub> and N<sub>2</sub>O, agricultural residue burning was also a minor source of the ambient air pollutants carbon monoxide (CO) and nitrogen oxides (NO<sub>x</sub>).

Figure 5-1



**Table 5-1: Emissions from Agriculture (Tg CO<sub>2</sub> Eq.)**

Gas/Source	1990	1995	1996	1997	1998	1999	2000
<b>CH<sub>4</sub></b>	<b>164.9</b>	<b>176.2</b>	<b>171.5</b>	<b>170.9</b>	<b>171.6</b>	<b>171.1</b>	<b>169.6</b>
Enteric Fermentation	127.9	133.2	129.6	126.8	124.9	124.5	123.9
Manure Management	29.2	34.8	34.2	35.8	38.0	37.6	37.5
Rice Cultivation	7.1	7.6	7.0	7.5	7.9	8.3	7.5
Agricultural Residue Burning	0.7	0.7	0.7	0.8	0.8	0.8	0.8
<b>N<sub>2</sub>O</b>	<b>283.5</b>	<b>300.2</b>	<b>309.8</b>	<b>315.0</b>	<b>316.0</b>	<b>313.9</b>	<b>315.5</b>
Agricultural Soil Management	267.1	283.4	292.6	297.5	298.4	296.3	297.6
Manure Management	16.0	16.4	16.8	17.1	17.1	17.1	17.5
Agricultural Residue Burning	0.4	0.4	0.4	0.4	0.5	0.4	0.5
<b>Total</b>	<b>448.4</b>	<b>476.4</b>	<b>481.3</b>	<b>485.9</b>	<b>487.6</b>	<b>485.0</b>	<b>485.1</b>

Note: Totals may not sum due to independent rounding.

**Table 5-2: Emissions from Agriculture (Gg)**

Gas/Source	1990	1995	1996	1997	1998	1999	2000
<b>CH<sub>4</sub></b>	<b>7,851</b>	<b>8,392</b>	<b>8,166</b>	<b>8,136</b>	<b>8,172</b>	<b>8,149</b>	<b>8,076</b>
Enteric Fermentation	6,089	6,342	6,171	6,037	5,948	5,929	5,898
Manure Management	1,390	1,657	1,628	1,707	1,811	1,788	1,784
Rice Cultivation	339	363	332	356	376	395	357
Agricultural Residue Burning	33	31	36	36	37	36	37
<b>N<sub>2</sub>O</b>	<b>914</b>	<b>968</b>	<b>999</b>	<b>1,016</b>	<b>1,019</b>	<b>1,012</b>	<b>1,018</b>
Agricultural Soil Management	862	914	944	960	963	956	960
Manure Management	52	53	54	55	55	55	57
Agricultural Residue Burning	1	1	1	1	1	1	1

Note: Totals may not sum due to independent rounding.

## Enteric Fermentation

Methane (CH<sub>4</sub>) is produced as part of normal digestive processes in animals. During digestion, microbes resident in an animal's digestive system ferment food consumed by the animal. This microbial fermentation process, referred to as enteric fermentation, produces methane as a by-product, which can be exhaled or eructated by the animal. The amount of methane produced and excreted by an individual animal depends primarily upon the animal's digestive system, and the amount and type of feed it consumes.

Among domesticated animal types, ruminant animals (e.g., cattle, buffalo, sheep, goats, and camels) are the major emitters of methane because of their unique digestive system. Ruminants possess a rumen, or large "fore-stomach," in which microbial fermentation breaks down the feed they consume into products that can be metabolized. The microbial fermentation that occurs in the rumen enables

them to digest coarse plant material that non-ruminant animals cannot. Ruminant animals, consequently, have the highest methane emissions among all animal types.

Non-ruminant domesticated animals (e.g., swine, horses, mules, and goats) also produce methane emissions through enteric fermentation, although this microbial fermentation occurs in the large intestine. These non-ruminants emit significantly less methane on a per-animal basis than ruminants because the capacity of the large intestine to produce methane is lower.

In addition to the type of digestive system, an animal's feed intake also affects methane emissions. In general, a higher feed intake leads to higher methane emissions. Feed intake is positively related to animal size, growth rate, and production (e.g., milk production, wool growth, pregnancy, or work). Therefore, feed intake varies among animal types as well as among different management practices for individual animal types.

**Table 5-3: CH<sub>4</sub> Emissions from Enteric Fermentation (Tg CO<sub>2</sub> Eq.)**

Livestock Type	1990	1995	1996	1997	1998	1999	2000
Beef Cattle	93.3	100.1	98.1	95.4	93.5	93.0	91.7
Dairy Cattle	28.7	27.5	26.1	26.0	25.9	26.2	26.9
Horses	1.9	2.0	2.0	2.0	2.0	2.0	2.0
Sheep	1.9	1.5	1.4	1.3	1.3	1.2	1.2
Swine	1.7	1.9	1.8	1.8	2.0	1.9	1.9
Goats	0.3	0.2	0.2	0.2	0.2	0.2	0.2
<b>Total</b>	<b>127.9</b>	<b>133.2</b>	<b>129.6</b>	<b>126.8</b>	<b>124.9</b>	<b>124.5</b>	<b>123.9</b>

Note: Totals may not sum due to independent rounding.

**Table 5-4: CH<sub>4</sub> Emissions from Enteric Fermentation (Gg)**

Livestock Type	1990	1995	1996	1997	1998	1999	2000
Beef Cattle	4,444	4,768	4,673	4,541	4,453	4,429	4,365
Dairy Cattle	1,369	1,308	1,241	1,240	1,234	1,246	1,283
Horses	93	94	94	94	95	96	96
Sheep	91	72	68	64	63	58	56
Swine	81	88	84	88	93	90	88
Goats	13	11	10	10	10	10	10
<b>Total</b>	<b>6,089</b>	<b>6,342</b>	<b>6,171</b>	<b>6,037</b>	<b>5,948</b>	<b>5,929</b>	<b>5,898</b>

Note: Totals may not sum due to independent rounding.

Methane emission estimates from enteric fermentation are provided in Table 5-3 and Table 5-4. Total livestock methane emissions in 2000 were 123.9 Tg CO<sub>2</sub> Eq. (5,898 Gg), decreasing slightly since 1999. Beef cattle remain the largest contributor of methane emissions from enteric fermentation, accounting for 74 percent in 2000. Emissions from dairy cattle in 2000 accounted for 22 percent, and the remaining 4 percent was from horses, sheep, swine, and goats.

## Methodology

Livestock emission estimates fall into two categories: cattle and other domesticated animals. Cattle, due to their large population, large size, and particular digestive characteristics, account for the majority of methane emissions from livestock in the United States. Cattle production systems in the United States are better characterized in comparison with other livestock management systems. A more detailed methodology (i.e., IPCC Tier 2) was therefore applied to estimating emissions for cattle. Emission estimates for other domesticated animals were handled using a less detailed approach (i.e., IPCC Tier 1).

While the large diversity of animal management practices cannot be precisely characterized and evaluated, significant scientific literature exists that describes the quantity of methane produced by individual ruminant animals, particularly cattle. A detailed model that incorporates this information and other analyses of livestock population, feeding practices and production characteristics was used to estimate emissions from cattle populations.

National cattle population statistics were disaggregated into the following cattle sub-populations:

### Dairy Cattle

- Calves
- Heifer Replacements
- Cows

### Beef Cattle

- Calves
- Heifer Replacements
- Heifer and Steer Stockers
- Animals in Feedlots
- Cows
- Bulls

Calf birth estimates, end of year population statistics, detailed feedlot placement information, and slaughter weight data were used in the model to initiate and track cohorts of individual animal types having distinct emissions profiles. The key variables tracked for each of the cattle population categories are described in Annex K. These variables include performance factors such as pregnancy and lactation as well as average weights and weight gain.

Diet characteristics were estimated by State and region for U.S. dairy, beef, and feedlot cattle. These estimates were used to calculate Digestible Energy (DE) values and methane conversion rates ( $Y_m$ ) for each population category. The IPCC recommends  $Y_m$  values of 3.5 to 4.5 percent for feedlot cattle and 5.5 to 6.5 percent for all other cattle. Given the availability of detailed diet information for different regions and animal types in the United States, DE and  $Y_m$  values unique to the United States were developed, rather than using the recommended IPCC values. The diet characterizations and estimation of DE and  $Y_m$  values were based on contact with state agricultural extension specialists, a review of published forage quality studies, expert opinion, and modeling of animal physiology. See Annex K for more details on the method used to characterize cattle diets in the United States.

In order to estimate methane emissions from cattle, the population was divided into region, age, sub-type (e.g., calves, heifer replacements, cows, etc.), and production (i.e., pregnant, lactating, etc.) groupings to more fully capture any differences in methane emissions from these animal types. Cattle diet characteristics developed under Step 2 were used to develop regional emission factors for each sub-category. Tier 2 equations from IPCC (2000) were used to produce methane emission factors for the following cattle types: dairy cows, beef cows, dairy replacements, beef replacements, steer stockers, heifer stockers, steer feedlot animals, heifer feedlot animals, and steer and heifer feedlot step-up diet animals. To estimate emissions from cattle, population data were multiplied by the emission factor for each cattle type. More details can be found in Annex K.

Emission estimates for other animal types were based upon average emission factors representative of entire populations of each animal type. Methane emissions from these animals accounted for a minor portion of total methane emissions from livestock in the United States from 1990

through 2000. Also, the variability in emission factors for each of these other animal types (e.g. variability by age, production system, and feeding practice within each animal type) is less than that for cattle.

See Annex K for more detailed information on the methodology and data used to calculate methane emissions from enteric fermentation.

## Data Sources

Annual cattle population data were obtained from the U.S. Department of Agriculture's National Agricultural Statistics Service (1995a-d, 1996b, 1997, 1998a, 1999a-c,f-g, 2000a,c,d,f, 2001a,c,d,g). DE and  $Y_m$  values were used to calculate emissions from cattle populations. DE and  $Y_m$  for dairy and beef cows, and for beef stockers, were calculated from diet characteristics using a model simulating ruminant digestion in growing and/or lactating cattle (Donovan and Baldwin 1999). For feedlot animals, DE and  $Y_m$  values recommended by Johnson (1999) were used. Values from EPA (1993) were used for dairy replacement heifers. Weight data were estimated from Feedstuffs (1998), Western Dairyman (1998), and expert opinion. Annual livestock population data for other livestock types, except horses, as well as feedlot placement information were obtained from the U.S. Department of Agriculture's National Agricultural Statistics Service (USDA 1994a-b, 1998b-c, 1999d,e,h, 2000b,e, 2001b,e,f). Horse data were obtained from the Food and Agriculture Organization (FAO) statistical database (FAO 2000, 2001). Methane emissions from sheep, goats, swine, and horses were estimated by using emission factors utilized in Crutzen et al. (1986). These emission factors are representative of typical animal sizes, feed intakes, and feed characteristics in developed countries. The methodology is the same as that recommended by IPCC (IPCC/UNEP/OECD/IEA 1997, IPCC 2000).

## Uncertainty

The basic uncertainties associated with estimating emissions from enteric fermentation are the range of emission factors possible for the different animal types and the number of animals with a particular emissions profile that exist during the year. Although determining an emission factor for all possible cattle sub-groupings and diet characterizations in the United States is not possible, the

enteric fermentation model that was used estimates the likely emission factors for the major animal types and diets. The model generates estimates for dairy and beef cows, dairy and beef replacements, beef stockers, and feedlot animals. The analysis departs from the recommended IPCC (2000) DE and  $Y_m$  values to account for diets for these different animal types regionally. Based on expert opinion and peer reviewer recommendations, it is believed that the values supporting the development of emission factors for the animal types studied are appropriate for the situation in the United States.

In addition to the uncertainty associated with developing emission factors for different cattle population categories based on estimated energy requirements and diet characterizations, there is uncertainty in the estimation of animal populations by animal type. The model estimates the movement of animal cohorts through the various monthly age and weight classes by animal type. Several inputs affect the precision of this approach, including estimates of births by month, weight gain of animals by age class, and placement of animals into feedlots based on placement statistics and slaughter weight data. However, it is believed that the model sufficiently characterizes the U.S. cattle population and captures the potential differences related to the emission factors used for different animal types.

In order to ensure the quality of the emission estimates from enteric fermentation, the IPCC Tier 1 and Tier 2 QA/QC procedures were implemented. Tier 1 procedures included quality checks on data gathering, input, and documentation, as well as checks on the actual emission calculations. Additionally, Tier 2 procedures included quality checks on emission factors, activity data, and emissions.

## **Manure Management**

The management of livestock manure can produce anthropogenic methane ( $CH_4$ ) and nitrous oxide ( $N_2O$ ) emissions. Methane is produced by the anaerobic decomposition of manure. Nitrous oxide is produced as part of the nitrogen cycle through the nitrification and denitrification of the organic nitrogen in livestock manure and urine.

When livestock or poultry manure are stored or treated in systems that promote anaerobic conditions (e.g., as a liquid in lagoons, ponds, tanks, or pits), the decomposition of materials in the manure tends to produce  $CH_4$ . When manure is handled as a solid (e.g., in stacks or pits) or deposited on pasture, range, or paddock lands, it tends to decompose aerobically and produce little or no  $CH_4$ . A number of other factors related to how the manure is handled also affect the amount of  $CH_4$  produced: 1) ambient temperature and moisture affect the amount of  $CH_4$  produced because they influence the growth of the bacteria responsible for methane formation; 2) methane production generally increases with rising temperature and residency time; and 3) for non-liquid based manure systems, moist conditions (which are a function of rainfall and humidity) favor  $CH_4$  production. Although the majority of manure is handled as a solid, producing little  $CH_4$ , the general trend in manure management, particularly for large dairy and swine producers, is one of increasing use of liquid systems. In addition, use of daily spread systems at smaller dairies is decreasing, due to new regulations limiting the application of manure nutrients, which has resulted in an increase of manure managed and stored on site at these smaller dairies.

The composition of the manure also affects the amount of methane produced. Manure composition varies by animal type, including the animal's digestive system and diet. In general, the greater the energy content of the feed, the greater the potential for  $CH_4$  emissions. For example, feedlot cattle fed a high energy grain diet generate manure with a high  $CH_4$ -producing capacity (represented by  $B_o$ ). Range cattle fed a low energy diet of forage material produce manure with about 50 percent of the  $CH_4$ -producing potential of feedlot cattle manure. In addition, there is a trend in the dairy industry indicating that dairy cows are producing more milk per year. These high-production milk cows tend to produce more volatile solids in their manure as milk production increases, which increases the probability of  $CH_4$  production.

A very small portion of the total nitrogen excreted is expected to convert to nitrous oxide in the waste management system. The production of nitrous oxide from livestock manure depends on the composition of the manure and urine, the type of bacteria involved in the process, and the amount of oxygen and liquid in the manure system. For

N<sub>2</sub>O emissions to occur, the manure must first be handled aerobically where ammonia or organic nitrogen is converted to nitrates and nitrites (nitrification), and then handled anaerobically where the nitrates and nitrites are reduced to nitrogen gas (N<sub>2</sub>), with intermediate production of nitrous oxide (N<sub>2</sub>O) and nitric oxide (NO) (denitrification) (Groffman, et al. 2000).

These emissions are most likely to occur in dry manure handling systems that have aerobic conditions, but that also contain pockets of anaerobic conditions due to saturation. For example, manure at cattle drylots is deposited on soil, oxidized to nitrite and nitrate, and has the potential to encounter saturated conditions following rain events.

Certain N<sub>2</sub>O emissions are accounted for and discussed under Agricultural Soil Management. These are emissions from livestock manure and urine deposited on pasture, range, or paddock lands, as well as emissions from manure and urine that is spread onto fields either directly as “daily spread” or after it is removed from manure management systems (e.g., lagoon, pit, etc.).

Table 5-5 and Table 5-6 provide estimates of CH<sub>4</sub> and N<sub>2</sub>O emissions from manure management by animal category. Estimates for methane emissions in 2000 were 37.5 Tg CO<sub>2</sub> Eq. (1,783 Gg), 28 percent higher than in 1990. The majority of this increase was from swine and dairy cow manure and is attributed to shifts by the swine and dairy industries towards larger facilities. Larger swine and dairy farms tend to use flush or scrape liquid systems to manage and store manure. Thus the shift towards larger facilities is translated into an increasing use of liquid manure management systems. This shift was accounted for by incorporating state-specific weighted methane conversion factor (MCF) values calculated from the 1992 and 1997 farm-size distribution reported in the *Census of Agriculture* (USDA 1999e). In 2000, swine CH<sub>4</sub> emissions decreased from 1999 due to a decrease in those animal populations.

As stated previously, smaller dairies are moving away from daily spread systems. Therefore, more manure is managed and stored on site, contributing to additional CH<sub>4</sub> emissions over the time series. The CH<sub>4</sub> estimates also account for changes in volatile solids production from dairy cows correlated to their generally increasing milk production. A description of the methodology is provided in Annex L.

Total N<sub>2</sub>O emissions from manure management systems in 2000 were estimated to be 17.5 Tg CO<sub>2</sub> Eq. (56.5 Gg). The 9 percent increase in N<sub>2</sub>O emissions from 1990 to 2000 can be partially attributed to a shift in the poultry industry away from the use of liquid manure management systems, in favor of litter-based systems and high rise houses. In addition, there was an overall increase in the population of poultry and swine from 1990 to 2000, although swine populations declined slightly in 1993, 1995, 1996, 1999, and 2000 from previous years and poultry populations decreased in 1995.

The population of beef cattle in feedlots, which tend to store and manage manure on site, also increased, resulting in increased N<sub>2</sub>O emissions from this animal category. Although dairy cow populations decreased overall, the population of dairies managing and storing manure on site—as opposed to using pasture, range, or paddock or daily spread systems—increased. Therefore, the increase in dairies using on-site storage to manage their manure results in increased N<sub>2</sub>O emissions. As stated previously, N<sub>2</sub>O emissions from livestock manure deposited on pasture, range, or paddock land and manure immediately applied to land in daily spread systems are accounted for under Agricultural Soil Management.

## Methodology

The methodologies presented in *Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories* (IPCC 2000) form the basis of the CH<sub>4</sub> and N<sub>2</sub>O emissions estimates for each animal type. The calculation of emissions requires the following information:

- Animal population data (by animal type and state)
- Amount of nitrogen produced (amount per 1000 pound animal times average weight times number of head)
- Amount of volatile solids produced (amount per 1000 pound animal times average weight times number of head)
- Methane producing potential of the volatile solids (by animal type)
- Extent to which the methane producing potential is realized for each type of manure management system (by State and manure management system)

**Table 5-5: CH<sub>4</sub> and N<sub>2</sub>O Emissions from Manure Management (Tg CO<sub>2</sub> Eq.)**

Gas/Animal Type	1990	1995	1996	1997	1998	1999	2000
<b>CH<sub>4</sub></b>	<b>29.2</b>	<b>34.8</b>	<b>34.2</b>	<b>35.9</b>	<b>38.0</b>	<b>37.6</b>	<b>37.5</b>
Dairy Cattle	9.6	12.0	12.1	12.7	13.1	13.3	13.7
Beef Cattle	3.2	3.5	3.5	3.4	3.4	3.4	3.4
Swine	13.0	16.0	15.3	16.4	18.1	17.6	17.1
Sheep	0.1	0.1	+	+	+	+	+
Goats	+	+	+	+	+	+	+
Poultry	2.7	2.6	2.6	2.7	2.7	2.6	2.6
Horses	0.6	0.6	0.6	0.6	0.6	0.6	0.6
<b>N<sub>2</sub>O</b>	<b>16.0</b>	<b>16.4</b>	<b>16.8</b>	<b>17.1</b>	<b>17.1</b>	<b>17.2</b>	<b>17.5</b>
Dairy Cattle	4.2	4.0	3.9	3.9	3.8	3.8	3.8
Beef Cattle	4.9	5.3	5.1	5.4	5.5	5.5	5.9
Swine	0.3	0.3	0.3	0.4	0.4	0.4	0.4
Sheep	+	+	+	+	+	+	+
Goats	+	+	+	+	+	+	+
Poultry	6.3	6.5	7.2	7.2	7.2	7.2	7.2
Horses	0.2	0.2	0.2	0.2	0.2	0.2	0.2
<b>Total</b>	<b>45.2</b>	<b>51.2</b>	<b>51.0</b>	<b>53.0</b>	<b>55.1</b>	<b>54.8</b>	<b>55.0</b>

+ Does not exceed 0.05 Tg CO<sub>2</sub> Eq.

Note: Totals may not sum due to independent rounding.

**Table 5-6: CH<sub>4</sub> and N<sub>2</sub>O Emissions from Manure Management (Gg)**

Gas/Animal Type	1990	1995	1996	1997	1998	1999	2000
<b>CH<sub>4</sub></b>	<b>1,390</b>	<b>1,657</b>	<b>1,628</b>	<b>1,707</b>	<b>1,811</b>	<b>1,788</b>	<b>1,783</b>
Dairy Cattle	457	571	577	604	624	634	653
Beef Cattle	151	166	165	163	161	160	161
Swine	621	763	729	782	864	839	814
Sheep	3	2	2	2	2	2	2
Goats	1	1	1	1	1	1	1
Poultry	128	124	125	127	130	124	124
Horses	29	29	29	29	29	30	30
<b>N<sub>2</sub>O</b>	<b>52</b>	<b>53</b>	<b>54</b>	<b>55</b>	<b>55</b>	<b>55</b>	<b>57</b>
Dairy Cattle	14	13	13	12	12	12	12
Beef Cattle	16	17	17	17	18	18	19
Swine	1	1	1	1	1	1	1
Sheep	+	+	+	+	+	+	+
Goats	+	+	+	+	+	+	+
Poultry	21	21	23	23	23	23	23
Horses	1	1	1	1	1	1	1

+ Does not exceed 0.5 Gg

Note: Totals may not sum due to independent rounding.

- Portion of manure managed in each manure management system (by State and animal type)
- Portion of manure deposited on pasture, range, or paddock or used in daily spread systems

Both CH<sub>4</sub> and N<sub>2</sub>O emissions were estimated by first determining activity data, including animal population, waste characteristics, and manure management system usage. For swine and dairy cattle, manure management system usage was determined for different farm size

categories using data from USDA (USDA 1996b, 1998d, 2000h) and EPA (ERG 2000a, EPA 2001a, 2001b). For beef cattle and poultry, manure management system usage data was not tied to farm size (ERG 2000a, USDA 2000i). For other animal types, manure management system usage was based on previous EPA estimates (EPA 1992).

Next, MCFs and N<sub>2</sub>O emission factors were determined for all manure management systems. MCFs for dry systems and N<sub>2</sub>O emission factors for all systems were set equal to

default IPCC factors (IPCC 2000). MCFs for liquid/slurry, anaerobic lagoon, and deep pit systems were calculated based on the forecast performance of biological systems relative to temperature changes as predicted in the van't Hoff-Arrhenius equation (see Annex L for detailed information on MCF derivations for liquid systems). The MCF calculations model the average monthly ambient temperature, a minimum system temperature, the carryover of volatile solids in the system from month to month due to long storage times exhibited by anaerobic lagoon systems, and a factor to account for management and design practices that result in the loss of volatile solids from lagoon systems.

For each animal group—except sheep, goats, and horses—the base emission factors were then weighted to incorporate the distribution of management systems used within each state and thereby to create an overall state-specific weighted emission factor. To calculate this weighted factor, the percent of manure for each animal group managed in a particular system in a state was multiplied by the emission factor for that system and state, and then summed for all manure management systems in the state.

Methane emissions were estimated by calculating the volatile solids (VS) production for all livestock. For each animal group except dairy cows, VS production was calculated using a national average VS production rate from the *Agricultural Waste Management Field Handbook* (USDA 1996a), which was then multiplied by the average weight of the animal and the State-specific animal population. For dairy cows, the national average VS constant was replaced with a mathematical relationship between milk production and VS (USDA 1996a), which was then multiplied by State-specific average annual milk production (USDA 2001g). The resulting VS for each animal group was then multiplied by the maximum methane producing capacity of the waste ( $B_0$ ), and the State-specific methane conversion factors.

Nitrous oxide emissions were estimated by determining total Kjeldahl nitrogen (TKN)<sup>1</sup> production for all livestock wastes using livestock population data and nitrogen excretion rates. For each animal group, TKN production was calculated using a national average nitrogen excretion rate from the *Agricultural Waste Management Field*

*Handbook* (USDA 1996a), which was then multiplied by the average weight of the animal and the State-specific animal population. State-specific weighted  $N_2O$  emission factors specific to the type of manure management system were then applied to total nitrogen production to estimate  $N_2O$  emissions.

See Annex L for more detailed information on the methodology and data used to calculate methane and nitrous oxide emissions from manure management.

## Data Sources

Animal population data for all livestock types, except horses and goats, were obtained from the U.S. Department of Agriculture's National Agricultural Statistics Service (USDA 1994a-b, 1995a-b, 1998a-b, 1999a-c, 2000a-g, 2001a-f). Horse population data were obtained from the FAOSTAT database (FAO 2001). Goat population data were obtained from the Census of Agriculture (USDA 1999d). Information regarding poultry turnover (i.e., slaughter) rate was obtained from State Natural Resource Conservation Service (NRCS) personnel (Lange 2000). Dairy cow and swine population data by farm size for each state, used for the weighted MCF and emission factor calculations, were obtained from the *Census of Agriculture*, which is conducted every five years (USDA 1999e).

Manure management system usage data for dairy and swine operations were obtained from USDA's Centers for Epidemiology and Animal Health (USDA 1996b, 1998d, 2000h) for small operations and from preliminary estimates for EPA's Office of Water regulatory effort for large operations (ERG 2000a; EPA 2001a, 2001b). Data for layers were obtained from a voluntary United Egg Producers' survey (UEP 1999), previous EPA estimates (EPA 1992), and USDA's Animal Plant Health Inspection Service (USDA 2000i). Data for beef feedlots were also obtained from EPA's Office of Water (ERG 2000a; EPA 2001a, 2001b). Manure management system usage data for other livestock were taken from previous EPA estimates (EPA 1992). Data regarding the use of daily spread and pasture, range, or paddock systems for dairy cattle were obtained from personal communications with personnel from several organizations, and data provided by those personnel (Poe

<sup>1</sup> Total Kjeldahl nitrogen is a measure of organically bound nitrogen and ammonia nitrogen.

et al. 1999). These organizations include State NRCS offices, State extension services, State universities, USDA National Agriculture Statistics Service (NASS), and other experts (Deal 2000, Johnson 2000, Miller 2000, Stettler 2000, Sweeten 2000, and Wright 2000). Additional information regarding the percent of beef steer and heifers on feedlots was obtained from contacts with the national USDA office (Milton 2000).

Methane conversion factors for liquid systems were calculated based on average ambient temperatures of the counties in which animal populations were located. The average county and state temperature data were obtained from the National Climate Data Center (NOAA 2001), and the county population data were based on 1992 and 1997 Census data (USDA 1999e). County population data for 1990 and 1991 were assumed to be the same as 1992; county population data for 1998 through 2000 were assumed to be the same as 1997; and county population data for 1993 through 1996 were extrapolated based on 1992 and 1997 data.

The maximum methane producing capacity of the volatile solids, or  $B_0$ , was determined based on data collected in a literature review (ERG 2000b).  $B_0$  data were collected for each animal type for which emissions were estimated.

Volatile solids and nitrogen excretion rate data from the USDA *Agricultural Waste Management Field Handbook* (USDA 1996a) were used for all livestock except sheep, goats, and horses. Data from the American Society of Agricultural Engineers (ASAE 1999) were used for these animal types. In addition, annual NASS data for average milk production per cow per State (USDA 2001g) were used to calculate state-specific volatile solids production rates for dairy cows for each year of the inventory. Nitrous oxide emission factors and MCFs for dry systems were taken from *Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories* (IPCC 2000).

## Uncertainty

The primary factors contributing to the uncertainty in emission estimates are a lack of information on the usage of various manure management systems in each regional location and the exact methane generating characteristics of each type of manure management system. Because of significant shifts in the swine and dairy sectors toward larger

farms, it is believed that increasing amounts of manure are being managed in liquid manure management systems. The existing estimates reflect these shifts in the weighted MCFs based on the 1992 and 1997 farm-size data. However, the assumption of a direct relationship between farm size and liquid system usage may not apply in all cases and may vary based on geographic location. In addition, the  $CH_4$  generating characteristics of each manure management system type are based on relatively few laboratory and field measurements, and may not match the diversity of conditions under which manure is managed nationally.

*Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories* (IPCC 2000) published a default range of MCFs for anaerobic lagoon systems of 0 to 100 percent, which reflects the wide range in performance that may be achieved with these systems. There exist relatively few data points on which to determine country-specific MCFs for these systems. In the United States, many livestock waste treatment systems classified as anaerobic lagoons are actually holding ponds that are substantially organically overloaded and therefore not producing methane at the same rate as a properly designed lagoon. In addition, these systems may not be well operated, contributing to higher loading rates when sludge is allowed to enter the treatment portion of the lagoon or the lagoon volume is pumped too low to allow treatment to occur. Rather than setting the MCF for all anaerobic lagoon systems in the United States based on data available from optimized lagoon systems, an MCF methodology was developed that more closely matches observed system performance and accounts for the affect of temperature on system performance.

However, there is uncertainty related to the new methodology. The MCF methodology used in the inventory includes a factor to account for management and design practices that result in the loss of volatile solids from the management system. This factor is currently estimated based on data from anaerobic lagoons in temperate climates, and from only three systems. However, this methodology is intended to account for systems across a range of management practices. Future work in gathering measurement data from animal waste lagoon systems across the country will contribute to the verification and refinement of this methodology. It will also be evaluated whether lagoon

temperatures differ substantially from ambient temperatures and whether the lower bound estimate of temperature established for lagoons and other liquid systems should be revised for use with this methodology.

The IPCC provides a suggested MCF for poultry waste management operations of 1.5 percent. Additional study is needed in this area to determine if poultry high-rise houses promote sufficient aerobic conditions to warrant a lower MCF.

The default N<sub>2</sub>O emission factors published in *Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories* (IPCC 2000) were derived using limited information. The IPCC factors are global averages; U.S.-specific emission factors may be significantly different. Manure and urine in anaerobic lagoons and liquid/slurry management systems produce methane at different rates, and would in all likelihood produce nitrous oxide at different rates, although a single N<sub>2</sub>O emission factors was used for both system types. In addition, there are little data available to determine the extent to which nitrification-denitrification occurs in animal waste management systems. Ammonia concentrations that are present in poultry and swine systems suggest that N<sub>2</sub>O emissions from these systems may be lower than predicted by the IPCC default factors. At this time, there are insufficient data available to develop U.S.-specific N<sub>2</sub>O emission factors; however, this is an area of on-going research, and warrants further study as more data become available.

Although an effort was made to introduce the variability in volatile solids production due to differences in diet for dairy cows, additional work is needed to establish the relationship between milk production and volatile solids production. In addition, the corresponding dairy methane emissions may be underestimated because milk production was unable to be correlated to specific manure management systems in each state. A methodology to assess variability in swine volatile solids production would be useful in future inventory estimates.

Uncertainty also exists with the maximum CH<sub>4</sub> producing potential of volatile solids excreted by different animal groups (i.e., B<sub>0</sub>). The B<sub>0</sub> values used in the CH<sub>4</sub> calculations are published values for U.S. animal waste. However, there are several studies that provide a range of

B<sub>0</sub> values for certain animals, including dairy and swine. The B<sub>0</sub> values chosen for dairy assign separate values for dairy cows and dairy heifers to better represent the feeding regimens of these animal groups. For example, dairy heifers do not receive an abundance of high energy feed and consequently, dairy heifer manure will not produce as much methane as manure from a milking cow. However, the data available for B<sub>0</sub> values are sparse, and do not necessarily reflect the rapid changes that have occurred in this industry with respect to feed regimens. Current research is being conducted to evaluate the usefulness of incorporating animal excretion data developed for the enteric fermentation emissions inventory into the manure management inventory calculations.

## Rice Cultivation

Most of the world's rice, and all rice in the United States, is grown on flooded fields. When fields are flooded, aerobic decomposition of organic material gradually depletes the oxygen present in the soil and floodwater, causing anaerobic conditions in the soil to develop. Once the environment becomes anaerobic, methane is produced through anaerobic decomposition of soil organic matter by methanogenic bacteria. As much as 60 to 90 percent of the methane produced is oxidized by aerobic methanotrophic bacteria in the soil (Holzapfel-Pschorn et al. 1985, Sass et al. 1990). Some of the CH<sub>4</sub> is also leached away as dissolved methane in floodwater that percolates from the field. The remaining un-oxidized CH<sub>4</sub> is transported from the submerged soil to the atmosphere primarily by diffusive transport through the rice plants. Minor amounts of methane also escape from the soil via diffusion and bubbling through floodwaters.

The water management system under which rice is grown is one of the most important factors affecting CH<sub>4</sub> emissions. Upland rice fields are not flooded, and therefore are not believed to produce CH<sub>4</sub>. In deepwater rice fields (i.e., fields with flooding depths greater than one meter), the lower stems and roots of the rice plants are dead so the primary CH<sub>4</sub> transport pathway to the atmosphere is blocked. The quantities of CH<sub>4</sub> released from deepwater fields, therefore, are believed to be significantly less than the quantities released from areas with more shallow flooding

depths. Some flooded fields are drained periodically during the growing season, either intentionally or accidentally. If water is drained and soils are allowed to dry sufficiently, CH<sub>4</sub> emissions decrease or stop entirely. This is due to soil aeration, which not only causes existing soil methane to oxidize but also inhibits further CH<sub>4</sub> production in soils. All rice in the United States is grown under continuously flooded conditions; none is grown under deepwater conditions. Mid-season drainage does not occur except by accident (e.g., due to levee breach).

Other factors that influence CH<sub>4</sub> emissions from flooded rice fields include fertilization practices (especially the use of organic fertilizers), soil temperature, soil type, rice variety, and cultivation practices (e.g., tillage, and seeding and weeding practices). The factors that determine the amount of organic material that is available to decompose (i.e., organic fertilizer use, soil type, rice variety,<sup>2</sup> and cultivation practices) are the most important variables influencing the amount of CH<sub>4</sub> emitted over an entire growing season because the total amount of CH<sub>4</sub> released depends primarily on the amount of organic substrate available. Soil temperature is known to be an important factor regulating the activity of methanogenic bacteria, and therefore the rate of CH<sub>4</sub> production. However, although temperature controls the amount of time it takes to convert a given amount of organic material to CH<sub>4</sub>, that time is short relative to a growing season, so the dependence of total emissions over an entire growing season on soil temperature is weak. The application of synthetic fertilizers has also been found to influence CH<sub>4</sub> emissions; in particular, both nitrate and sulfate fertilizers (e.g., ammonium nitrate, and ammonium sulfate) appear to inhibit CH<sub>4</sub> formation.

Rice is cultivated in seven states: Arkansas, California, Florida, Louisiana, Mississippi, Missouri, and Texas. Soil types, rice varieties, and cultivation practices for rice vary from state to state, and even from farm to farm. However, most rice farmers utilize organic fertilizers in the form of rice residue from the previous crop, which is left standing, disked, or rolled into the fields. Most farmers also apply synthetic fertilizer to their fields, usually urea. Nitrate and sulfate fertilizers are not commonly used in rice cultivation in the United States. In addition, the climatic conditions of

Arkansas, southwest Louisiana, Texas, and Florida allow for a second, or ratoon, rice crop. This second rice crop is produced from regrowth of the stubble after the first crop has been harvested. Because the first crop's stubble is left behind in ratooned fields, and there is no time delay between cropping seasons (which would allow for the stubble to decay aerobically), the amount of organic material that is available for decomposition is considerably higher than with the first (i.e., primary) crop. Methane emissions from ratoon crops have been found to be considerably higher than those from the primary crop.

Rice cultivation is a small source of CH<sub>4</sub> in the United States (Table 5-7 and Table 5-8). In 2000, CH<sub>4</sub> emissions from rice cultivation were 7.5 Tg CO<sub>2</sub> Eq. (357 Gg). Although annual emissions fluctuated considerably between the years 1990 and 2000, there was an overall increase of 5 percent over the ten year period due to an overall increase in harvested area. However, between 1990 and 1999 there was a 17 percent increase in emissions, which highlights the annual variability in the estimates. In 2000, the harvest area was the largest since 1996, as seen in Table 5-9.

The factors that affect the rice acreage in any year vary from State to State, although the price of rice relative to competing crops is the primary controlling variable in most States. Price is the primary factor affecting rice area in Arkansas, as farmers will plant more of what is most lucrative amongst soybeans, rice, and cotton. Government support programs have also been influential in so much as they affect the price received for a rice crop (Slaton 2001b, Mayhew 1997). California rice area is primarily influenced by price and government programs, but is also affected by water availability (Mutters 2001). In Florida, the State having the smallest harvested rice area, rice acreage is largely a function of the price of rice relative to sugarcane and corn. Most rice in Florida is rotated with sugarcane, but sometimes it is more profitable for farmers to follow their sugarcane crop with sweet corn or more sugarcane instead of rice (Schueneman 1997, 2001b). In Louisiana, rice area is influenced by government support programs, the price of rice relative to cotton, soybeans, and corn, and in some years, weather (Saichuk 1997, Linscombe 2001b). For example, a drought in 2000 caused extensive saltwater

<sup>2</sup> The roots of rice plants shed organic material, which is referred to as "root exudate." The amount of root exudate produced by a rice plant over a growing season varies among rice varieties.

**Table 5-7: CH<sub>4</sub> Emissions from Rice Cultivation (Tg CO<sub>2</sub> Eq.)**

State	1990	1995	1996	1997	1998	1999	2000
Arkansas	2.1	2.4	2.1	2.5	2.7	2.9	2.5
California	0.7	0.8	0.9	0.9	0.8	0.9	1.0
Florida	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Louisiana	2.1	2.2	2.0	2.2	2.3	2.3	2.1
Mississippi	0.4	0.5	0.4	0.4	0.5	0.6	0.4
Missouri	0.1	0.2	0.2	0.2	0.3	0.3	0.3
Texas	1.6	1.4	1.3	1.1	1.3	1.1	1.1
<b>Total</b>	<b>7.1</b>	<b>7.6</b>	<b>7.0</b>	<b>7.5</b>	<b>7.9</b>	<b>8.3</b>	<b>7.5</b>

Note: Totals may not sum due to independent rounding.

**Table 5-8: CH<sub>4</sub> Emissions from Rice Cultivation (Gg)**

Site	1990	1995	1996	1997	1998	1999	2000
Arkansas	102	114	99	118	126	138	120
California	34	40	42	44	39	43	47
Florida	3	6	5	5	5	5	4
Louisiana	98	102	96	105	111	111	101
Mississippi	21	24	18	20	23	27	19
Missouri	7	10	8	10	12	16	15
Texas	75	67	63	55	60	55	52
<b>Total</b>	<b>339</b>	<b>363</b>	<b>332</b>	<b>356</b>	<b>376</b>	<b>395</b>	<b>357</b>

Note: Totals may not sum due to independent rounding.

**Table 5-9: Rice Areas Harvested (Hectares)**

State/Crop	1990	1995	1996	1997	1998	1999	2000
Arkansas							
Primary	485,633	542,291	473,493	562,525	600,971	657,628	570,619
Ratoon*	NO	NO	NO	NO	202	202	NO
California	159,854	188,183	202,347	208,822	185,350	204,371	221,773
Florida							
Primary	4,978	9,713	8,903	7,689	8,094	7,229	7,801
Ratoon	2,489	4,856	4,452	3,845	4,047	4,673	3,193
Louisiana							
Primary	220,558	230,676	215,702	235,937	250,911	249,292	194,253
Ratoon	66,168	69,203	64,711	70,781	75,273	74,788	77,701
Mississippi	101,174	116,552	84,176	96,317	108,458	130,716	88,223
Missouri	32,376	45,326	38,446	47,349	57,871	74,464	70,417
Texas							
Primary	142,857	128,693	120,599	104,816	114,529	104,816	86,605
Ratoon	57,143	51,477	48,240	41,926	45,811	41,926	43,302
<b>Total</b>	<b>1,273,229</b>	<b>1,386,969</b>	<b>1,261,068</b>	<b>1,380,008</b>	<b>1,451,518</b>	<b>1,550,106</b>	<b>1,363,888</b>

Note: Totals may not sum due to independent rounding.

\* Arkansas ratooning occurred only in 1998 and 1999.

NO (Not Occurring)

intrusion along the Gulf Coast, making over 32,000 hectares unplanted. In Mississippi, rice is usually rotated with soybeans, but if soybean prices increase relative to rice prices, then some of the acreage that would have been planted in rice, is instead planted in soybeans (Street 1997, 2001). In Missouri, rice acreage is affected by weather (e.g., rain during the planting season may prevent the planting of rice), the price differential between rice and soybeans or cotton, and government support programs (Stevens 1997, Guethle 2001). In Texas, rice area is affected mainly by the price of rice, government support programs, and water availability (Klosterboer 1997, 2001b).

## Methodology

The *Revised 1996 IPCC Guidelines* (IPCC/UNEP/OECD/IEA 1997) recommends utilizing annual harvested rice areas and area-based seasonally integrated emission factors (i.e., amount of CH<sub>4</sub> emitted over a growing season per unit harvested area) to estimate annual CH<sub>4</sub> emissions from rice cultivation. This methodology is followed but a United States specific emission factor is used in the calculations. In addition, because daily average emissions have been found to be much higher for ratooned crops than for primary crops, emissions from ratooned and primary areas are estimated separately. This is consistent with IPCC *Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories* (IPCC 2000).

## Data Sources

The harvested rice areas for the primary and ratoon crops in each state are presented in Table 5-9. Data for 1990 through 2000 for all states except Florida were taken from U.S. Department of Agriculture's *Field Crops Final Estimates 1987-1992* (USDA 1994), *Field Crops Final Estimates 1992-1997* (USDA 1998), *Crop Production 1999 Summary* (USDA 2000), and *Crop Production 2000 Summary* (USDA 2001). Harvested rice areas in Florida, which are not reported by USDA, were obtained from Tom Schueneman (1999b, 1999c, 2000, 2001a), a Florida agricultural extension agent. Acreages for the ratoon crops

were derived from conversations with the agricultural extension agents in each state. In Arkansas, ratooning occurred only in 1998 and 1999, when the ratooned area was less than 1 percent of the primary area (Slaton 1999, 2000, 2001a). In Florida, the ratooned area was 50 percent of the primary area from 1990 to 1998 (Schueneman 1999a), about 65 percent of the primary area in 1999 (Schueneman 2000), and around 41 percent of the primary area in 2000 (Schueneman 2001a). In Louisiana, the percentage of the primary area that was ratooned was constant at 30 percent over the 1990 to 1999 period, but increased to approximately 40 percent in 2000 (Linscombe 1999a, 2001a and Bollich 2000). In Texas, the percentage of the primary area that was ratooned was constant at 40 percent over the entire 1990 to 1999 period, but increased to 50 percent in 2000 due to an early primary crop (Klosterboer 1999, 2000, 2001a).

To determine what seasonal CH<sub>4</sub> emission factors should be used for the primary and ratoon crops, CH<sub>4</sub> flux information from rice field measurements in the United States was collected. Experiments which involved the application of nitrate or sulfate fertilizers, or other substances believed to suppress CH<sub>4</sub> formation, as well as experiments in which measurements were not made over an entire flooding season or in which floodwaters were drained mid-season, were excluded from the analysis. The remaining experimental results<sup>3</sup> were then sorted by season (i.e., primary and ratoon) and type of fertilizer amendment (i.e., no fertilizer added, organic fertilizer added, and synthetic and organic fertilizer added). The experimental results from primary crops with synthetic and organic fertilizer added (Bossio et al. 1999, Cicerone et al. 1992, Sass et al. 1991a and 1991b) were averaged to derive an emission factor for the primary crop, and the experimental results from ratoon crops with synthetic fertilizer added (Lindau and Bollich 1993, Lindau et al. 1995) were averaged to derive an emission factor for the ratoon crop. The resultant emission factor for the primary crop is 210 kg CH<sub>4</sub>/hectare-season, and the resultant emission factor for the ratoon crop is 780 kg CH<sub>4</sub>/hectare-season.

<sup>3</sup> In some of these remaining experiments, measurements from individual plots were excluded from the analysis because of the reasons just mentioned. In addition, one measurement from the ratooned fields (i.e., the flux of 2.041 g/m<sup>2</sup>/day in Lindau and Bollich 1993) was excluded since this emission rate is unusually high compared to other flux measurements in the United States, as well as in Europe and Asia (IPCC/UNEP/OECD/IEA 1997).

## Uncertainty

The largest uncertainty in the calculation of CH<sub>4</sub> emissions from rice cultivation is associated with the emission factors. Seasonal emissions, derived from field measurements in the United States, vary by more than one order of magnitude. This variability is due to differences in cultivation practices, particularly the type, amount, and mode of fertilizer application; differences in cultivar type; and differences in soil and climatic conditions. Some of this variability is accounted for by separating primary from ratooned areas. However, even within a cropping season, measured emissions vary significantly. Of the experiments that were used to derive the emission factors used here, primary emissions ranged from 22 to 479 kg CH<sub>4</sub>/hectare-season and ratoon emissions ranged from 481 to 1,490 kg CH<sub>4</sub>/hectare-season. Based on these emission ranges, total CH<sub>4</sub> emissions from rice cultivation in 2000 were estimated to range from 1.8 to 16 Tg CO<sub>2</sub> Eq. (87 to 779 Gg).

A second source of uncertainty is the ratooned area data, which are not compiled regularly. However, this is a relatively minor source of uncertainty, as these areas account for less than 10 percent of the total area. Expert judgment was used to estimate these areas.

The last source of uncertainty is in the practice of flooding outside of the normal rice season. According to agriculture extension agents, all of the rice-growing States practice this on some part of their rice acreage. Estimates of these areas range from 5 to 33 percent of the rice acreage. Fields are flooded for a variety of reasons: to provide habitat for waterfowl, to provide ponds for crawfish production, and to aid in rice straw decomposition. To date, methane flux measurements have not been undertaken in these flooded areas, so this activity is not included in the emission estimates presented here.

## Agricultural Soil Management

Nitrous oxide (N<sub>2</sub>O) is produced naturally in soils through the microbial processes of nitrification and denitrification.<sup>4</sup> A number of agricultural activities add nitrogen to soils, thereby increasing the amount of nitrogen available for nitrification and denitrification, and ultimately the amount of N<sub>2</sub>O emitted. These activities may add nitrogen to soils either directly or indirectly (Figure 5-2). Direct additions occur through various soil management practices and from the deposition of manure on soils by animals on pasture, range, and paddock (i.e., by animals whose manure is not managed). Soil management practices that add nitrogen to soils include fertilizer use, application of managed livestock manure, disposal of sewage sludge, production of nitrogen-fixing crops, retention of crop residues, and cultivation of histosols (i.e., soils with a high organic matter content, otherwise known as organic soils).<sup>5</sup> Indirect additions of nitrogen to soils occur through two mechanisms: 1) volatilization and subsequent atmospheric deposition of applied nitrogen;<sup>6</sup> and 2) surface runoff and leaching of applied nitrogen into groundwater and surface water. Other agricultural soil management practices, such as irrigation, drainage, tillage practices, and fallowing of land, can affect fluxes of N<sub>2</sub>O, as well as other greenhouse gases, to and from soils. However, because there are significant uncertainties associated with these other fluxes, they have not been estimated.

Agricultural soil management is the largest source of N<sub>2</sub>O in the United States.<sup>7</sup> Estimated emissions from this source in 2000 were 297.4 Tg CO<sub>2</sub> Eq. (959 Gg N<sub>2</sub>O) (see Table 5-10 and Table 5-11). Although annual agricultural soil management emissions fluctuated between 1990 and 2000, there was a general increase in emissions over the


<sup>4</sup> Nitrification and denitrification are two processes within the nitrogen cycle that are brought about by certain microorganisms in soils. Nitrification is the aerobic microbial oxidation of ammonium (NH<sub>4</sub>) to nitrate (NO<sub>3</sub>), and denitrification is the anaerobic microbial reduction of nitrate to dinitrogen gas (N<sub>2</sub>). Nitrous oxide is a gaseous intermediate product in the reaction sequence of denitrification, which leaks from microbial cells into the soil and then into the atmosphere. Nitrous oxide is also produced during nitrification, although by a less well understood mechanism (Nevison 2000).

<sup>5</sup> Cultivation of histosols does not, *per se*, “add” nitrogen to soils. Instead, the process of cultivation enhances mineralization of old, nitrogen-rich organic matter that is present in histosols, thereby enhancing N<sub>2</sub>O emissions from histosols.

<sup>6</sup> These processes entail volatilization of applied nitrogen as ammonia (NH<sub>3</sub>) and oxides of nitrogen (NO<sub>x</sub>), transformations of these gases within the atmosphere (or upon deposition), and deposition of the nitrogen primarily in the form of particulate ammonium (NH<sub>4</sub>), nitric acid (HNO<sub>3</sub>), and oxides of nitrogen.

<sup>7</sup> Note that the emission estimates for this source category include applications of nitrogen to *all* soils, but the term “Agricultural Soil Management” is kept for consistency with the reporting structure of the *Revised 1996 IPCC Guidelines* (IPCC/UNEP/OECD/IEA 1997).

Figure 5-2



eleven-year period (see Annex M for a complete time series of emission estimates). This general increase in emissions was due primarily to an increase in synthetic fertilizer use, manure production, and crop production over this period. The year-to-year fluctuations are largely a reflection of annual variations in synthetic fertilizer consumption and crop production. Over the eleven-year period, total emissions of N<sub>2</sub>O from agricultural soil management increased by approximately 11 percent.

Estimated direct and indirect N<sub>2</sub>O emissions, by subsourse, are provided in Table 5-12, Table 5-13, and Table 5-14.

### Methodology

The methodology used to estimate emissions from agricultural soil management is consistent with the *Revised 1996 IPCC Guidelines* (IPCC/UNEP/OECD/IEA 1997), as amended by the *IPCC Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories* (IPCC 2000). The *Revised 1996 IPCC Guidelines* divide this N<sub>2</sub>O source category into three components: (1) direct emissions from managed soils due to applied nitrogen and cultivation of histosols; (2) direct emissions from soils due to the deposition of manure by livestock on pasture, range, and paddock; and (3) indirect emissions from soils induced by applied nitrogen.

Table 5-10: N<sub>2</sub>O Emissions from Agricultural Soil Management (Tg CO<sub>2</sub> Eq.)

Activity	1990	1995	1996	1997	1998	1999	2000
<b>Direct</b>	<b>193.5</b>	<b>204.8</b>	<b>212.1</b>	<b>217.2</b>	<b>218.4</b>	<b>216.5</b>	<b>217.5</b>
Managed Soils	153.0	161.1	168.6	175.0	177.1	175.5	177.1
Pasture, Range, & Paddock Livestock	40.4	43.6	43.6	42.2	41.4	41.0	40.5
<b>Indirect</b>	<b>73.6</b>	<b>78.6</b>	<b>80.3</b>	<b>80.0</b>	<b>79.8</b>	<b>79.8</b>	<b>79.8</b>
<b>Total</b>	<b>267.1</b>	<b>283.4</b>	<b>292.4</b>	<b>297.2</b>	<b>298.3</b>	<b>296.3</b>	<b>297.4</b>

Note: Totals may not sum due to independent rounding.

Table 5-11: N<sub>2</sub>O Emissions from Agricultural Soil Management (Gg)

Activity	1990	1995	1996	1997	1998	1999	2000
<b>Direct</b>	<b>624</b>	<b>661</b>	<b>684</b>	<b>701</b>	<b>705</b>	<b>698</b>	<b>702</b>
Managed Soils	494	520	544	564	571	566	571
Pasture, Range, & Paddock Livestock	130	141	140	136	133	132	131
<b>Indirect</b>	<b>237</b>	<b>254</b>	<b>259</b>	<b>258</b>	<b>257</b>	<b>257</b>	<b>257</b>
<b>Total</b>	<b>862</b>	<b>914</b>	<b>943</b>	<b>959</b>	<b>962</b>	<b>956</b>	<b>959</b>

Note: Totals may not sum due to independent rounding.

**Table 5-12: Direct N<sub>2</sub>O Emissions from Managed Soils (Tg CO<sub>2</sub> Eq.)**

Activity	1990	1995	1996	1997	1998	1999	2000
Commercial Fertilizers*	55.4	59.2	61.2	61.3	61.4	61.6	61.7
Livestock Manure	12.7	13.2	13.2	13.4	13.6	13.8	13.8
Sewage Sludge	0.4	0.6	0.6	0.7	0.7	0.7	0.7
N Fixation	58.5	61.8	63.9	68.2	69.2	68.2	69.0
Crop Residue	23.2	23.4	26.8	28.7	29.3	28.3	29.1
Histosol Cultivation	2.8	2.8	2.8	2.9	2.9	2.9	2.9
<b>Total</b>	<b>153.0</b>	<b>161.1</b>	<b>168.6</b>	<b>175.0</b>	<b>177.1</b>	<b>175.5</b>	<b>177.1</b>

Note: Totals may not sum due to independent rounding.

\* Excludes sewage sludge and livestock manure used as commercial fertilizers.

**Table 5-13: Direct N<sub>2</sub>O Emissions from Pasture, Range, and Paddock Livestock Manure (Tg CO<sub>2</sub> Eq.)**

Animal Type	1990	1995	1996	1997	1998	1999	2000
Beef Cattle	35.2	38.9	39.0	37.8	37.0	36.7	36.2
Dairy Cows	1.7	1.5	1.4	1.3	1.3	1.2	1.2
Swine	0.5	0.3	0.3	0.2	0.2	0.2	0.2
Sheep	0.4	0.3	0.3	0.3	0.3	0.3	0.3
Goats	0.2	0.2	0.2	0.2	0.2	0.2	0.2
Poultry	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Horses	2.3	2.3	2.3	2.3	2.3	2.3	2.3
<b>Total</b>	<b>40.4</b>	<b>43.6</b>	<b>43.6</b>	<b>42.2</b>	<b>41.4</b>	<b>41.0</b>	<b>40.5</b>

Note: Totals may not sum due to independent rounding.

**Table 5-14: Indirect N<sub>2</sub>O Emissions (Tg CO<sub>2</sub> Eq.)**

Activity	1990	1995	1996	1997	1998	1999	2000
<b>Volatilization &amp; Atm. Deposition</b>	<b>11.6</b>	<b>12.4</b>	<b>12.6</b>	<b>12.6</b>	<b>12.5</b>	<b>12.5</b>	<b>12.5</b>
Commercial Fertilizers*	4.9	5.3	5.4	5.5	5.5	5.5	5.5
Livestock Manure	6.6	7.0	7.1	7.0	6.9	6.9	6.9
Sewage Sludge	0.1	0.1	0.1	0.1	0.1	0.1	0.1
<b>Surface Leaching &amp; Runoff</b>	<b>62.0</b>	<b>66.2</b>	<b>67.6</b>	<b>67.4</b>	<b>67.3</b>	<b>67.3</b>	<b>67.3</b>
Commercial Fertilizers*	36.9	39.5	40.8	40.9	40.9	41.1	41.1
Livestock Manure	24.7	26.3	26.4	26.1	25.8	25.7	25.7
Sewage Sludge	0.3	0.5	0.5	0.5	0.5	0.5	0.5
<b>Total</b>	<b>73.6</b>	<b>78.6</b>	<b>80.3</b>	<b>80.0</b>	<b>79.8</b>	<b>79.8</b>	<b>79.8</b>

Note: Totals may not sum due to independent rounding.

\* Excludes sewage sludge and livestock manure used as commercial fertilizers.

Annex M provides more detailed information on the methodologies and data used to calculate N<sub>2</sub>O emissions from each of these three components.

### **Direct N<sub>2</sub>O Emissions from Managed Soils**

Direct N<sub>2</sub>O emissions from managed soils are composed of two parts, which are estimated separately and then summed. These two parts are 1) emissions due to nitrogen applications, and 2) emissions from histosol cultivation.

Estimates of direct N<sub>2</sub>O emissions from nitrogen applications were based on the total amount of nitrogen that is applied to soils annually through the following practices: (a) the application of synthetic and organic commercial fertilizers, (b) the application of livestock manure through both daily spread operations and through the eventual application of manure that had been stored in manure management systems, (c) the application of sewage sludge, (d) the production of nitrogen-fixing crops, and (e) the retention of crop residues (i.e., leaving residues in the field after harvest). For each of these practices, the annual amounts of nitrogen applied were estimated as follows:

a) Synthetic and organic commercial fertilizer nitrogen applications were derived from annual fertilizer consumption data and the nitrogen content of the fertilizers.

b) Livestock manure nitrogen applications were based on the assumption that all livestock manure is applied to soils except for two components: 1) a small portion of poultry manure that is used as a livestock feed supplement, and 2) the manure from pasture, range, and paddock livestock. The manure nitrogen data were derived from animal population and weight statistics, information on manure management system usage, annual nitrogen excretion rates for each animal type, and information on the fraction of poultry litter that is used as a livestock feed supplement.

c) Sewage sludge nitrogen applications were derived from estimates of annual U.S. sludge production, the nitrogen content of the sludge, and periodic surveys of sludge disposal methods.

d) The amounts of nitrogen made available to soils through the cultivation of nitrogen-fixing crops were based on estimates of the amount of nitrogen in aboveground plant biomass, which were derived from annual crop production statistics, mass ratios of aboveground residue to crop product, dry matter fractions, and nitrogen contents of the plant biomass.

e) Crop residue nitrogen retention data were derived from information about which residues are typically left on the field, the fractions of residues left on the field, annual crop production statistics, mass ratios of aboveground residue to crop product, and dry matter fractions and nitrogen contents of the residues.

After the annual amounts of nitrogen applied were estimated for each practice, each amount of nitrogen was reduced by the fraction that is assumed to volatilize according to the *Revised 1996 IPCC Guidelines* and the *IPCC Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories*. The net amounts left on the soil from each practice were then summed to yield total unvolatilized applied nitrogen, which was multiplied by the IPCC default emission factor for nitrogen applications.

Estimates of annual N<sub>2</sub>O emissions from histosol cultivation were based on estimates of the total U.S. acreage of histosols cultivated annually for each of two climatic zones: 1) temperate, and 2) sub-tropical. To estimate annual emissions, the total temperate area was multiplied by the IPCC default emission factor for temperate regions, and the total sub-tropical area was multiplied by the average of the IPCC default emission factors for temperate and tropical regions.<sup>8</sup>

Total annual emissions from nitrogen applications, and annual emissions from histosol cultivation, were then summed to estimate total direct emissions from managed soils.

### **Direct N<sub>2</sub>O Emissions from Pasture, Range, and Paddock Livestock Manure**

Estimates of N<sub>2</sub>O emissions from this component are based on the amount of nitrogen in the manure that is deposited annually on soils by livestock in pasture, range, and paddock. Estimates of annual manure nitrogen from

<sup>8</sup> Note that the IPCC default emission factors for histosols have been revised in the *IPCC Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories* (IPCC 2000). These revised default emission factors (IPCC 2000) were used in these calculations.

these livestock were derived from animal population and weight statistics; information on the fraction of the total population of each animal type that is on pasture, range, or paddock; and annual nitrogen excretion rates for each animal type. The annual amounts of manure nitrogen from each animal type were summed over all animal types to yield total pasture, range, and paddock manure nitrogen, which was then multiplied by the IPCC default emission factor for pasture, range, and paddock nitrogen to estimate N<sub>2</sub>O emissions.

### Indirect N<sub>2</sub>O Emissions from Soils

Indirect emissions of N<sub>2</sub>O are composed of two parts, which are estimated separately and then summed. These two parts are 1) emissions resulting from volatilization and subsequent deposition of the nitrogen in applied fertilizers, applied sewage sludge, and all livestock manure,<sup>9</sup> and 2) leaching and runoff of nitrogen in applied fertilizers, applied sewage sludge, and applied plus deposited livestock manure. The activity data (i.e., nitrogen in applied fertilizers, applied sewage sludge, all livestock manure, and applied plus deposited livestock manure) were estimated in the same way as for the direct emission estimates.

To estimate the annual amount of applied nitrogen that volatilizes, the annual amounts of applied synthetic fertilizer nitrogen, applied sewage sludge nitrogen, and all livestock manure nitrogen, were each multiplied by the appropriate IPCC default volatilization fraction. The three amounts of volatilized nitrogen were then summed, and the sum was multiplied by the IPCC default emission factor for volatilized/deposited nitrogen.

To estimate the annual amount of nitrogen that leaches or runs off, the annual amounts of applied synthetic fertilizer nitrogen, applied sewage sludge nitrogen, and applied plus deposited livestock manure nitrogen were each multiplied by the IPCC default leached/runoff fraction. The three amounts of leached/runoff nitrogen were then summed, and the sum was multiplied by the IPCC default emission factor for leached/runoff nitrogen.

Total annual indirect emissions from volatilization, and annual indirect emissions from leaching and runoff, were then summed to estimate total indirect emissions of N<sub>2</sub>O from managed soils.

## Data Sources

The activity data used in these calculations were obtained from numerous sources. Annual synthetic and organic fertilizer consumption data for the United States were obtained from annual publications on commercial fertilizer statistics (TVA 1991, 1992a, 1993, 1994; AAPFCO 1995, 1996, 1997, 1998, 1999, 2000b). Fertilizer nitrogen contents were taken from these same publications and AAPFCO (2000a). Livestock population data were obtained from USDA publications (USDA 1994b,c; 1995a,b; 1998a,c; 1999a-e; 2000a-g; 2001b-g), the FAOSTAT database (FAO 2001), and Lange (2000). Manure management information was obtained from Poe et al. (1999), Safley et al. (1992), and personal communications with agricultural experts (Anderson 2000, Deal 2000, Johnson 2000, Miller 2000, Milton 2000, Stettler 2000, Sweeten 2000, Wright 2000). Livestock weight data were obtained from Safley (2000), USDA (1996, 1998d), and ASAE (1999); daily rates of nitrogen excretion from ASAE (1999) and USDA (1996); and information about the fraction of poultry litter used as a feed supplement from Carpenter (1992). Data collected by the EPA were used to derive annual estimates of land application of sewage sludge (EPA 1993, 1999). The nitrogen content of sewage sludge was taken from Metcalf and Eddy, Inc. (1991). Annual production statistics for nitrogen-fixing crops were obtained from USDA reports (USDA 1994a, 1998b, 2000i, 2001a), a book on forage crops (Taylor and Smith 1995, Pederson 1995, Beuselinck and Grant 1995, Hoveland and Evers 1995), and personal communications with forage experts (Cropper 2000, Gerrish 2000, Hoveland 2000, Evers 2000, and Pederson 2000). Mass ratios of aboveground residue to crop product, dry matter fractions, and nitrogen contents for nitrogen-fixing crops were obtained from Stüttele (1987), Barnard and Kristoferson (1985), Karkosh (2000), Ketzis (1999), and IPCC/UNEP/OECD/IEA (1997).

<sup>9</sup> Total livestock manure nitrogen is used in the calculation of indirect N<sub>2</sub>O emissions from volatilization because all manure nitrogen, regardless of how the manure is managed or used, is assumed to be subject to volatilization.

Annual production statistics for crops whose residues are left on the field, except for rice in Florida, were obtained from USDA reports (USDA 1994a, 1998b, 2000i, 2001a). Production statistics for rice in Florida are not recorded by USDA, so these were derived from Smith (2001) and Schueneman (1999, 2001). Aboveground residue to crop mass ratios, residue dry matter fractions, and residue nitrogen contents were obtained from Strehler and Stützel (1987), Turn et al. (1997), Ketzis (1999), and Barnard and Kristoferson (1985). Estimates of the fractions of residues left on the field were based on information provided by Karkosh (2000), and on information about rice residue burning (see the Agricultural Residue Burning section). The annual areas of cultivated histosols were estimated from 1982, 1992, and 1997 statistics in USDA's 1997 *National Resources Inventory* (USDA 2000h, as extracted by Eve 2001, and revised by Ogle 2002).

All emission factors, volatilization fractions, and the leaching/runoff fraction were taken from the *Revised 1996 IPCC Guidelines* (IPCC/UNEP/OECD/IEA 1997), as amended by the IPCC *Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories* (IPCC 2000).

## Uncertainty

The amount of N<sub>2</sub>O emitted from managed soils depends not only on N inputs, but also on a large number of variables, including organic carbon availability, O<sub>2</sub> partial pressure, soil moisture content, pH, soil temperature, and soil amendment management practices. However, the effect of the combined interaction of these other variables on N<sub>2</sub>O flux is complex and highly uncertain. Therefore, the IPCC default methodology, which is used here, is based only on N inputs and does not utilize these other variables. As noted in the *Revised 1996 IPCC Guidelines* (IPCC/UNEP/OECD/IEA 1997), this is a generalized approach that treats all soils, except cultivated histosols, as being under the same conditions. The estimated ranges around the IPCC default emission factors provide an indication of the uncertainty in the emission estimates due to this simplified methodology. Most of the emission factor ranges are about an order of magnitude, or larger. Developing an emission estimation

methodology that explicitly utilizes these other variables will require more scientific research and much more detailed databases, and will likely involve the use of process models.

Uncertainties also exist in the activity data used to derive emission estimates. In particular, the fertilizer statistics include only those organic fertilizers that enter the commercial market, so non-commercial fertilizers (other than the estimated manure and crop residues) have not been captured. The livestock excretion values, while based on detailed population and weight statistics, were derived using simplifying assumptions concerning the types of management systems employed. Statistics on sewage sludge applied to soils were not available on an annual basis; annual production and application estimates were based on figures and projections that were calculated from surveys that yielded uncertainty levels as high as 14 percent (Bastian 1999). Annual data were obtained by interpolating and extrapolating at constant rates from these uncertain figures, though change between the years was unlikely to be constant (Bastian 2001). The production statistics for the nitrogen-fixing crops that are forage legumes are highly uncertain because statistics are not compiled for any of these crops except alfalfa, and the alfalfa statistics include alfalfa mixtures. Conversion factors for the nitrogen-fixing crops were based on a limited number of studies, and may not be representative of all conditions in the United States. Data on crop residues left on the field are not available, so expert judgment was used to estimate the amount of residues left on soils. And finally, the estimates of cultivated histosol areas are uncertain because they are from a natural resource inventory that was not explicitly designed as a soil survey, and this natural resource inventory contains data for only three years (1982, 1992, and 1997). Annual histosol areas were estimated by linear interpolation and extrapolation.

## Agricultural Residue Burning

Large quantities of agricultural crop residues are produced by farming activities. There are a variety of ways to dispose of these residues. For example, agricultural residues can be left on or plowed back into the field, composted and then applied to soils, landfilled, or burned

in the field. Alternatively, they can be collected and used as a fuel or sold in supplemental feed markets. Field burning of crop residues is not considered a net source of carbon dioxide (CO<sub>2</sub>) because the carbon released to the atmosphere as CO<sub>2</sub> during burning is assumed to be reabsorbed during the next growing season. Crop residue burning is, however, a net source of methane (CH<sub>4</sub>), nitrous oxide (N<sub>2</sub>O), carbon monoxide (CO), and nitrogen oxides (NO<sub>x</sub>), which are released during combustion.

Field burning is not a common method of agricultural residue disposal in the United States; therefore, emissions from this source are minor. The primary crop types whose residues are typically burned in the United States are wheat, rice, sugarcane, corn, barley, soybeans, and peanuts. Of these residues, less than 5 percent is burned each year, except for rice.<sup>10</sup> Annual emissions from this source over the period 1990 through 2000 averaged approximately 0.7 Tg CO<sub>2</sub> Eq. (35 Gg) of CH<sub>4</sub>, 0.4 Tg CO<sub>2</sub> Eq. (1 Gg) of N<sub>2</sub>O, 725 Gg of CO, and 31 Gg of NO<sub>x</sub> (see Table 5-15 and Table 5-16).

## Methodology

The methodology for estimating greenhouse gas emissions from field burning of agricultural residues is consistent with the *Revised 1996 IPCC Guidelines* (IPCC/UNEP/OECD/IEA 1997). In order to estimate the amounts of carbon and nitrogen released during burning, the following equations were used:<sup>11</sup>

$$\text{Carbon Released} = (\text{Annual Crop Production}) \times (\text{Residue/Crop Product Ratio}) \times (\text{Fraction of Residues Burned } in \text{ situ}) \times (\text{Dry Matter Content of the Residue}) \times (\text{Burning Efficiency}) \times (\text{Carbon Content of the Residue}) \times (\text{Combustion Efficiency})^{12}$$

$$\text{Nitrogen Released} = (\text{Annual Crop Production}) \times (\text{Residue/Crop Product Ratio}) \times (\text{Fraction of Residues Burned } in \text{ situ}) \times (\text{Dry Matter Content of the Residue}) \times (\text{Burning Efficiency}) \times (\text{Nitrogen Content of the Residue}) \times (\text{Combustion Efficiency})$$

**Table 5-15: Emissions from Agricultural Residue Burning (Tg CO<sub>2</sub> Eq.)**

Gas/Crop Type	1990	1995	1996	1997	1998	1999	2000
<b>CH<sub>4</sub></b>	<b>0.7</b>	<b>0.7</b>	<b>0.7</b>	<b>0.8</b>	<b>0.8</b>	<b>0.8</b>	<b>0.8</b>
Wheat	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Rice	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Sugarcane	+	+	+	+	+	+	+
Corn	0.3	0.3	0.3	0.3	0.3	0.3	0.4
Barley	+	+	+	+	+	+	+
Soybeans	0.1	0.2	0.2	0.2	0.2	0.2	0.2
Peanuts	+	+	+	+	+	+	+
<b>N<sub>2</sub>O</b>	<b>0.4</b>	<b>0.4</b>	<b>0.4</b>	<b>0.4</b>	<b>0.5</b>	<b>0.4</b>	<b>0.5</b>
Wheat	+	+	+	+	+	+	+
Rice	+	+	+	+	+	+	+
Sugarcane	+	+	+	+	+	+	+
Corn	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Barley	+	+	+	+	+	+	+
Soybeans	0.2	0.2	0.2	0.3	0.3	0.3	0.3
Peanuts	+	+	+	+	+	+	+
<b>Total</b>	<b>1.1</b>	<b>1.0</b>	<b>1.2</b>	<b>1.2</b>	<b>1.2</b>	<b>1.2</b>	<b>1.2</b>

+ Does not exceed 0.05 Tg CO<sub>2</sub> Eq.

Note: Totals may not sum due to independent rounding.

<sup>10</sup> The fraction of rice straw burned each year is significantly higher than that for other crops (see “Data Sources” discussion below).

<sup>11</sup> Note: As is explained below, the fraction of residues burned varies among states for rice, so these equations were applied at the state level for rice. These equations were applied at the national level for all other crop types.

<sup>12</sup> Burning Efficiency is defined as the fraction of dry biomass exposed to burning that actually burns. Combustion Efficiency is defined as the fraction of carbon in the fire that is oxidized completely to CO<sub>2</sub>. In the methodology recommended by the IPCC, the “burning efficiency” is assumed to be contained in the “fraction of residues burned” factor. However, the number used here to estimate the “fraction of residues burned” does not account for the fraction of exposed residue that does not burn. Therefore, a “burning efficiency factor” was added to the calculations.

**Table 5-16: Emissions from Agricultural Residue Burning (Gg)\***

Gas/Crop Type	1990	1995	1996	1997	1998	1999	2000
<b>CH<sub>4</sub></b>	<b>33</b>	<b>31</b>	<b>36</b>	<b>36</b>	<b>37</b>	<b>36</b>	<b>37</b>
Wheat	7	5	5	6	6	5	5
Rice	4	4	4	3	3	3	3
Sugarcane	1	1	1	1	1	1	1
Corn	13	13	16	16	17	16	17
Barley	1	1	1	1	1	+	1
Soybeans	7	8	9	10	10	10	10
Peanuts	+	+	+	+	+	+	+
<b>N<sub>2</sub>O</b>	<b>1</b>						
Wheat	+	+	+	+	+	+	+
Rice	+	+	+	+	+	+	+
Sugarcane	+	+	+	+	+	+	+
Corn	+	+	+	+	+	+	+
Barley	+	+	+	+	+	+	+
Soybeans	1	1	1	1	1	1	1
Peanuts	+	+	+	+	+	+	+
<b>CO</b>	<b>685</b>	<b>656</b>	<b>747</b>	<b>761</b>	<b>781</b>	<b>760</b>	<b>786</b>
Wheat	137	109	114	124	128	115	111
Rice	81	80	85	66	58	69	69
Sugarcane	18	20	19	21	22	23	24
Corn	282	263	328	328	347	336	355
Barley	16	13	15	13	13	10	12
Soybeans	148	167	183	207	211	204	213
Peanuts	2	2	2	2	2	2	2
<b>NO<sub>x</sub></b>	<b>28</b>	<b>29</b>	<b>32</b>	<b>34</b>	<b>35</b>	<b>34</b>	<b>35</b>
Wheat	4	3	3	3	3	3	3
Rice	3	3	3	2	2	2	2
Sugarcane	+	+	+	+	+	+	+
Corn	7	6	8	8	8	8	9
Barley	1	+	+	+	+	+	+
Soybeans	14	16	17	20	20	19	20
Peanuts	+	+	+	+	+	+	+

\* Full molecular weight basis.

+ Does not exceed 0.5 Gg

Note: Totals may not sum due to independent rounding.

Emissions of CH<sub>4</sub> and CO were calculated by multiplying the amount of carbon released by the appropriate IPCC default emission ratio (i.e., CH<sub>4</sub>-C/C or CO-C/C). Similarly, N<sub>2</sub>O and NO<sub>x</sub> emissions were calculated by multiplying the amount of nitrogen released by the appropriate IPCC default emission ratio (i.e., N<sub>2</sub>O-N/N or NO<sub>x</sub>-N/N).

## Data Sources

The crop residues that are burned in the United States were determined from various state level greenhouse gas emission inventories (ILENR 1993, Oregon Department of Energy 1995, Wisconsin Department of Natural Resources 1993) and publications on agricultural burning in the United States (Jenkins et al. 1992, Turn et al. 1997, EPA 1992).

Crop production data for all crops except rice in Florida were taken from the USDA's *Field Crops, Final Estimates 1987-1992, 1992-1997* (USDA 1994, 1998), *Crop Production 1999 Summary* (USDA 2000), and *Crop Production 2000 Summary* (USDA 2001). Crop production data for Florida, which are not collected by USDA, were estimated by applying average primary and ratoon crop yields for Florida (Smith 2001) to Florida acreages (Schueneman 1999b, 2001). The production data for the crop types whose residues are burned are presented in Table 5-17.

The percentage of crop residue burned was assumed to be 3 percent for all crops in all years, except rice, based on state inventory data (ILENR 1993, Oregon Department of

**Table 5-17: Agricultural Crop Production (Thousand Metric Tons of Product)**

Crop	1990	1995	1996	1997	1998	1999	2000
Wheat	74,292	59,404	61,980	67,534	69,327	62,569	60,512
Rice	7,105	7,935	7,828	8,339	8,570	9,381	8,708
Sugarcane	25,525	27,922	26,729	28,766	30,896	32,023	32,973
Corn*	201,534	187,970	234,518	233,864	247,882	239,549	253,208
Barley	9,192	7,824	8,544	7,835	7,667	6,103	6,921
Soybeans	52,416	59,174	64,780	73,176	74,598	72,223	75,338
Peanuts	1,635	1,570	1,661	1,605	1,798	1,737	1,491

\*Corn for grain (i.e., excludes corn for silage).

**Table 5-18: Percentage of Rice Area Burned by State**

State	Percent Burned 1990-1998	Percent Burned 1999	Percent Burned 2000
Arkansas	10	10	10
California	variable <sup>a</sup>	27	27
Florida <sup>b</sup>	0	0	0
Louisiana	6	0	5
Mississippi	10	40	40
Missouri	5	5	8
Texas	1	2	0

<sup>a</sup> Values provided in Table 5-19.

<sup>b</sup> Burning of crop residues is illegal in Florida.

**Table 5-19: Percentage of Rice Area Burned in California**

Year	California
1990	75
1995	59
1996	63
1997	34
1998	33
1999	27
2000	27

Energy 1995, Noller 1996, Wisconsin Department of Natural Resources 1993, and Cibrowski 1996). Estimates of the percentage of rice acreage on which residue burning took place were obtained on a state-by-state basis from agricultural extension agents in each of the seven rice-producing states (Bollich 2000; Guethle 1999, 2000, 2001; Fife 1999; California Air Resources Board 1999; Klosterboer 1999a, 1999b, 2000, 2001; Linscombe 1999a, 1999b, 2001; Najita 2000, 2001; Schueneman 1999a, 1999b, 2001; Slaton 1999a, 1999b, 2000; Street 1999a, 1999b, 2000, 2001; Wilson 2001) (see Table 5-18 and Table 5-19).

The estimates provided for Arkansas and Florida remained constant over the entire 1990 through 2000 period, while the estimates for all other states varied over the time series. For California, it was assumed that the annual percents of rice acreage burned in Sacramento Valley are representative of burning in the entire state, because the Sacramento Valley accounts for over 95 percent of the rice acreage in California (Fife 1999). The annual percents of rice acreage burned in Sacramento Valley were obtained from a report of the California Air Resources Board (2001). These values declined over the 1990 through 2000 period because of a legislated reduction in rice straw burning (see Table 5-19).

All residue/crop product mass ratios except sugarcane were obtained from Strehler and Stützel (1987). The datum for sugarcane is from University of California (1977). Residue dry matter contents for all crops except soybeans and peanuts were obtained from Turn et al. (1997). Soybean dry matter content was obtained from Strehler and Stützel (1987). Peanut dry matter content was obtained through personal communications with Jen Ketzis (1999), who accessed Cornell University's Department of Animal Science's computer model, Cornell Net Carbohydrate and

Protein System. The residue carbon contents and nitrogen contents for all crops except soybeans and peanuts are from Turn et al. (1997). The residue carbon content for soybeans and peanuts is the IPCC default (IPCC/UNEP/OECD/IEA 1997). The nitrogen content of soybeans is from Barnard and Kristoferson (1985). The nitrogen content of peanuts is from Ketzis (1999). These data are listed in Table 5-20. The burning efficiency was assumed to be 93 percent, and the combustion efficiency was assumed to be 88 percent, for all crop types (EPA 1994). Emission ratios for all gases (see Table 5-21) were taken from the *Revised 1996 IPCC Guidelines* (IPCC/UNEP/OECD/IEA 1997).

## Uncertainty

The largest source of uncertainty in the calculation of non-CO<sub>2</sub> emissions from field burning of agricultural residues is in the estimates of the fraction of residue of each crop type burned each year. Data on the fraction burned, as well as the gross amount of residue burned each year, are

not collected at either the national or State level. In addition, burning practices are highly variable among crops, as well as among States. The fractions of residue burned used in these calculations were based upon information collected by State agencies and in published literature. It is likely that these emission estimates will continue to change as more information becomes available in the future.

Other sources of uncertainty include the residue/crop product mass ratios, residue dry matter contents, burning and combustion efficiencies, and emission ratios. A residue/crop product ratio for a specific crop can vary among cultivars, and for all crops except sugarcane, generic residue/crop product ratios, rather than ratios specific to the United States, have been used. Residue dry matter contents, burning and combustion efficiencies, and emission ratios, all can vary due to weather and other combustion conditions, such as fuel geometry. Values for these variables were taken from literature on agricultural biomass burning.

**Table 5-20: Key Assumptions for Estimating Emissions from Agricultural Residue Burning\***

Crop	Residue/ Crop Ratio	Fraction of Residue Burned	Dry Matter Fraction	Carbon Fraction	Nitrogen Fraction
Wheat	1.3	0.03	0.93	0.4428	0.0062
Rice	1.4	variable	0.91	0.3806	0.0072
Sugarcane	0.8	0.03	0.62	0.4235	0.0040
Corn	1.0	0.03	0.91	0.4478	0.0058
Barley	1.2	0.03	0.93	0.4485	0.0077
Soybeans	2.1	0.03	0.87	0.4500	0.0230
Peanuts	1.0	0.03	0.86	0.4500	0.0106

\* The burning efficiency and combustion efficiency for all crops were assumed to be 0.93 and 0.88, respectively.

**Table 5-21: Greenhouse Gas Emission Ratios**

Gas	Emission Ratio
CH <sub>4</sub> <sup>a</sup>	0.005
CO <sup>a</sup>	0.060
N <sub>2</sub> O <sup>b</sup>	0.007
NO <sub>x</sub> <sup>b</sup>	0.121

<sup>a</sup> Mass of carbon compound released (units of C) relative to mass of total carbon released from burning (units of C).  
<sup>b</sup> Mass of nitrogen compound released (units of N) relative to mass of total nitrogen released from burning (units of N).

