



FIGURE 3-1
MAP OF COMMUNITIES IN OU1
FIVE-YEAR REVIEW

N





Legend

- 0 - 349.9 mg/kg
- 350 - 999.9 mg/kg
- 1,000 - 4,999.9 mg/kg
- 5,000 and greater mg/kg

N

FIGURE 3-9a

**CURRENT SUBSURFACE SOIL
(BELOW BARRIER)
LEAD CONCENTRATIONS
FOR SMELTERVILLE, IDAHO**

BUNKER HILL SUPERFUND SITE FIVE-YEAR REVIEW

Figure 3-6
Children's Blood Lead Levels by Year, 1974-2002




Figure 3-5
House Dust Vacuum Bag Lead Concentration by City, 1988-2004






Figure 3-4
Percentage of Yard Remediations Completed by City, 1989-2004^a



^aThe number of residential properties was originally estimated from tax assessor parcel maps. The number of remediated residential yards is based on the PRP soil database, which is not based on tax assessor parcel maps. Therefore, 100% agreement between counts is not expected.

Note: Kellogg includes outlying communities such as Elizabeth Park and Montgomery Gulch.

Figure 3-3
Community Geometric Mean Soil Lead Concentration and Progress Towards Remedial Action Objective (RAO), 1989-2004






FIGURE 3-2

OU1 TIMELINE BUNKER HILL SUPERFUND SITE

FIVE-YEAR REVIEW

