

 <p style="text-align: center;">U.S. ENVIRONMENTAL PROTECTION AGENCY Office of Pesticide Programs Registration Division (7505C) 1200 Pennsylvania Ave., N.W. Washington, D.C. 20460</p> <p style="text-align: center;">NOTICE OF PESTICIDE: <input checked="" type="checkbox"/> Registration <input type="checkbox"/> Reregistration (under FIFRA, as amended)</p>	EPA Reg. Number:	Date of Issuance:
	71711-26	AUG 01 2008
	Term of Issuance:	Conditional
		Name of Pesticide Product:
		NNI-0001 Technical
Name and Address of Registrant (include ZIP Code):		
Nichino America, Inc. c/o Bayer CropScience LP 2 T.W. Alexander Drive Research Triangle Park, NC 27709-2014		
Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.		
On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.		
Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.		
This product is conditionally registered in accordance with FIFRA section 3(c)(7) provided that you:		
1. Make the following change to the label:		
a. Change the product registration number to "EPA Reg. No. 71711-26"		
(continued on page 2)		
Signature of Approving Official:	Date:	
Refer to Page #2.		
Richard J. Gebken, Product Manager (10) Insecticide Branch, Registration Division (7505P)	AUG 01 2008	

EPA Form 8570-6

2. Submit two (2) copies of the final printed labeling before releasing the product for shipment.

2/4

Your release for shipment of these products constitutes acceptance of the conditions of registration as outlined in the preliminary acceptance letter for flubendiamide, dated July 31, 2008. If these conditions are not complied with, the registration will be subject to cancellation in accordance with section 6(e) of FIFRA.

A stamped "Accepted" copy of the label for this product is enclosed for your records.

Sincerely yours,



Richard J. Gebken,
Product Manager (10)
Insecticide Branch,
Registration Division (7505P)

Enclosures: Copy of label for NNI-0001 Technical stamped "Accepted," dated August 1, 2008

071711-00264 D366875

3/4

ACCEPTED
August 1, 2008
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 71711-26

NNI-0001 Technical

For Use in the Manufacture of Insecticides

ACTIVE INGREDIENT:

Flubendiamide* (N-[1,1-dimethyl-2-(methylsulfonyl)ethyl]-3-iodo-N-[2-methyl-4-[1,2,2,2-tetrafluoro-1-(trifluoromethyl)ethyl]phenyl]-1,2-benzenedicarboxamide)..... 97.76%

OTHER INGREDIENTS:..... 2.24%

Total:..... 100.00%

*CAS Number: 272451-65-7

EPA Reg. No. 71711-26

EPA Est. No.

KEEP OUT OF REACH OF CHILDREN CAUTION

For **MEDICAL** And **TRANSPORTATION** Emergencies **ONLY** Call 24 Hours A Day 1-800-334-7577

For **PRODUCT USE** Information Call 1-866-99BAYER (1-866-992-2937)

FIRST AID

IF INHALED	<ul style="list-style-type: none"> • Move the person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice.
IF SWALLOWED:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Have person sip a glass of water if able to swallow. • Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

For medical emergencies, health concerns, or pesticide incidents, you may call the Bayer CropScience Emergency Response toll free number 24 hours a day at 1-800-334-7577.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if inhaled, swallowed or absorbed through skin. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Wear long-sleeved shirt and long pants, socks, shoes and chemical-resistant gloves. Remove and wash contaminated clothing before reuse.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to aquatic invertebrates. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans, or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance, contact your State Water Board or Regional Office of the EPA.

4/4

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This product may be used only for formulation into an insecticide for:

1. the following uses:
Terrestrial Food and Feed Crops: Brassica (Cole) Leafy Vegetables, Corn (Field Corn, Pop Corn, Sweet Corn, Silage, and Corn Grown for Seed), Cotton, Cucurbit Vegetables, Fruiting Vegetables, Grapes, Leafy Vegetables (except Brassica), Okra, Pome Fruit, Stone Fruit, Tobacco, and Tree Nuts.
2. uses for which the U.S. EPA has accepted the required data and/or citations of data that the formulator has submitted in support of registration and
3. uses for experimental purposes that are in compliance with U.S. EPA requirements.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE

Do not store for more than 30 consecutive days at an average daily temperature exceeding 100° F. Keep container tightly closed when not in use. Avoid cross contamination with other pesticides.

PESTICIDE DISPOSAL

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL

Completely empty bag by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into the processing equipment. Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available, or dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Nichino America, Inc. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, NICHINO AMERICA, INC. MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Nichino America, Inc. is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, NICHINO AMERICA, INC. DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

LIMITATIONS OF LIABILITY: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT NICHINO AMERICA, INC.'S ELECTION, THE REPLACEMENT OF PRODUCT.

NET CONTENTS:

Nichino America, Inc.
4550 New Linden Hill Road
Suite 501
Wilmington, DE 19808

264-1025

8/01/2008

1/12



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs
Registration Division (7505C)
1200 Pennsylvania Ave., N.W.
Washington, D.C. 20460

EPA Reg. Number:

264-1025

Date of Issuance:

AUG 01 2008

NOTICE OF PESTICIDE:

Registration
 Reregistration
(under FIFRA, as amended)

Term of Issuance:

Conditional

Name of Pesticide Product:

NNI-0001 480 SC

Name and Address of Registrant (include ZIP Code):

Bayer CropScience LP
2 T.W. Alexander Drive
Research Triangle Park, NC 27709-2014

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA section 3(c)(7) provided that you:

1. Make the following change to the label:
 - a. Change the product registration number to "EPA Reg. No. 264-1025"

(continued on page 2)

Signature of Approving Official:

Refer to Page #2.

Richard J. Gebken, Product Manager (10)
Insecticide Branch, Registration Division (7505P)

Date:

AUG 01 2008

EPA Form 8570-6

2/
12

2. Submit two (2) copies of the final printed labeling before releasing the product for shipment.

Your release for shipment of these products constitutes acceptance of the conditions of registration as outlined in the preliminary acceptance letter for flubendiamide, dated July 31, 2008. If these conditions are not complied with, the registration will be subject to cancellation in accordance with section 6(e) of FIFRA.

A stamped "Accepted" copy of the label for this product is enclosed for your records.

Sincerely yours,



Richard J. Gebken,
Product Manager (10)
Insecticide Branch,
Registration Division (7505P)

Enclosures: Copy of label for NNI-0001 480 SC stamped "Accepted," dated August 1, 2008

000264-01025 D366878

3/12

GROUP	28	INSECTICIDE
-------	----	-------------

NNI-0001 480 SC

ACTIVE INGREDIENT:

Flubendiamide (*N*²-[1,1-dimethyl-2-(methylsulfonyl)ethyl]-3-iodo-*N*¹-[2-methyl-4-[1,2,2,2-tetrafluoro-1-(trifluoromethyl)ethyl]phenyl]-1,2-benzenedicarboxamide)..... 39%

OTHER INGREDIENTS: **61%**

NNI-0001 480 SC contains 4 pounds of flubendiamide per US gallon (480 grams per liter). **TOTAL:** **100%**

EPA Reg. No. 264-1025

EPA Est. No.

STOP - Read the label before use
KEEP OUT OF REACH OF CHILDREN
CAUTION

For **MEDICAL** And **TRANSPORTATION** Emergencies **ONLY** Call 24 Hours A Day 1-800-334-7577
For **PRODUCT USE** Information Call 1-866-99BAYER (1-866-992-2937)

FIRST AID

IF ON SKIN OR CLOTHING:

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

IF SWALLOWED:

- Call a poison control center or doctor immediately for treatment advice.
- Do not induce vomiting unless told to do so by a poison control center or doctor.
- Have person sip a glass of water if able to swallow.
- Do not give anything by mouth to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

For medical emergencies, health concerns, or pesticide incidents, you may call the Bayer CropScience Emergency Response toll free number 24 hours a day at 1-800-334-7577.

NOTE TO PHYSICIAN: No specific antidote is known. Treat symptomatically.

PRECAUTIONARY STATEMENTS

HAZARD TO HUMANS AND DOMESTIC ANIMALS**CAUTION**

Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves (such as Natural Rubber). If you want more options, follow the instructions for Category A on the EPA chemical-resistance category selection chart.
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

ACCEPTED
August 1, 2008
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
Registered under
EPA Reg. No. 264-1025

4
12

ENGINEERING CONTROLS STATEMENT

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands thoroughly before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove Personal Protective Equipment immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to aquatic invertebrates. For terrestrial uses: Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

Ground Water Advisory

Flubendiamide and its degradate NNI-0001-des-iodo have properties and characteristics associated with chemicals detected in ground water. This chemical may leach into ground water if used in areas where soils are permeable, particularly where the water table is shallow.

Surface Water Advisory

Flubendiamide and its degradate NNI-0001-des-iodo may also impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. These chemicals are classified as having a medium potential for reaching both surface water and aquatic sediment via runoff several months or more after application. A vegetative buffer strip as required under the Directions for Use will reduce the potential for loading of flubendiamide and its degradate NNI-0001-des-iodo from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
Read entire label before using this product.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the same area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

BUFFER ZONES

Vegetative Buffer Strip

Construct and maintain a minimum 15-foot wide vegetative filter strip of grass or other permanent vegetation between field edge and down gradient aquatic habitat (such as, but not limited to, lakes; reservoirs; rivers; permanent streams; marshes or natural ponds; estuaries; and commercial fish farm ponds).

Only apply products containing flubendiamide onto fields where a maintained vegetative buffer strip of at least 15 feet exists between the field edge and down gradient aquatic habitat.

For guidance, refer to the following publication for information on constructing and maintaining effective buffers: *Conservation Buffers to Reduce Pesticide Losses*. Natural Resources Conservation Services, USDA, 2000, Fort Worth, Texas, 21 pp. <http://www.in.ces.usda/v/technical/agronom/newconbuf.pdf>

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours following application.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated such as plants, soil or water, is: coveralls, chemical-resistant gloves such as barrier laminate, butyl rubber, nitrile rubber, or viton, and shoes plus socks.

5
12

GENERAL INFORMATION

NNI-0001 480 SC is a Suspension Concentrate formulation. The active ingredient contained in NNI-0001 480 SC is active by insect larval ingestion leading to a rapid cessation of feeding followed by death of the insect. Application should be timed to coincide with early threshold level in a developing larval population. Thorough coverage of all plant parts is required for optimum performance.

RESISTANCE MANAGEMENT

NNI-0001 480 SC contains an active ingredient with a novel mode of action. Studies to determine cross-resistance with NNI-0001 480 SC linked to other commercial insecticide have demonstrated no cross-resistance. However, repeated use of any crop protection product may increase the development of resistant strains of pests, including insects and mites. Rotation to another product with a different mode of action is recommended.

APPLICATION GUIDELINES

For all insects, timing of application should be based on careful scouting and local thresholds.

Foliar Spray Applications

Ground applications: A minimum of 10 gallons of diluted product/A.

Aerial applications: A minimum of 5 gallons of diluted product/A. Aerial applications made to dense canopies may not provide sufficient coverage of lower leaves to provide acceptable pest control. Under these conditions, the higher rate of NNI-0001 480 SC specified in the crop/pest specific tables within the Directions for Use section of this label may be necessary for optimum pest control.

Chemigation applications (see use in Chemigation Systems directions below) should be made as concentrated as possible. For best results apply at 100% input/travel speed, for center pivots or 0.10 inch (2,716 gallons) up to 0.15 inch (4,073 gallons) of water/A, for other systems. Higher labeled rates of NNI-0001 480 SC may be necessary for chemigation applications.

CHEMIGATION SYSTEMS

NNI-0001 480 SC may be applied through irrigation systems only on those crops listed under Recommended Applications where application through irrigation systems is recommended.

Types of Irrigation Systems: Apply NNI-0001 480 SC only through sprinkler, including center pivot, lateral move, side roll, or overhead solid set irrigation systems. Do not apply NNI-0001 480 SC through any other type of irrigation system.

GENERAL DIRECTIONS FOR ALL RECOMMENDED TYPES OF IRRIGATION SYSTEMS

Uniform Water Distribution and System Calibration: The irrigation system must provide uniform distribution of treated water. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water. The system must be calibrated to uniformly apply the rates specified. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

Chemigation Monitoring: A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Drift: Do not apply when wind speed favors drift beyond the area intended for treatment.

Required System Safety Devices: The system must contain a functional check valve, vacuum relief valve, and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. The irrigation line or water pump must include a functional pressure switch that will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Using Water from Public Water Systems: Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone (RPZ), back flow preventer or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the flow outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection. The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Cleaning the Chemical Injection System: In order to accurately apply pesticides, the chemical injection system must be kept clean; free of chemical or fertilizer residues and sediments. Refer to your owner's manual or ask your equipment supplier for the cleaning procedure for your injection system.

Flushing the Irrigation System: At the end of the application period, allow time for all lines to flush the pesticide through all nozzles before turning off irrigation water. To ensure the lines are flushed and free of pesticides, a dye indicator may be injected into the lines to mark the end of the application period.

Equipment Area Contamination Prevention: It is recommended that nozzles in the immediate area of control panels, chemical supply tanks, pumps and system safety devices be plugged to prevent chemical contamination of these areas

Center-Pivot and Automatic-Move Linear Systems: Inject the specified dosage per acre continuously for one complete revolution or move of the system. DO NOT USE END GUNS. The system should be run at maximum speed.

Solid Set and Manually Controlled Linear Systems: Injection should be during the last 30 to 60 minutes of regular irrigation period or as a separate 30 to 60 minute application not associated with a regular irrigation. Adjust end guns to keep treated water on the treated area in a uniform manner

SPRAY DRIFT REDUCTION MANAGEMENT

Do not apply when wind speed favors drift beyond the area intended for treatment. The interaction of many equipment and weather related factors determine the potential for spray drift. The applicator is responsible for considering all of these factors when making application decisions. Avoiding spray drift is the responsibility of the applicator.

Importance of Droplet Size:

An important factor influencing drift is droplet size. Small droplets (<150 - 200 microns) drift to a greater extent than large droplets. Within typical equipment specifications, applications should be made to deliver the largest droplet spectrum that provides sufficient control and coverage. Use only Medium or coarser spray nozzles (for ground and non-ULV aerial application) according to ASAE (S572) definition for standard nozzles. In conditions of low humidity and high temperatures, applicators should use a coarser droplet size.

Ground Applications:

Wind speed must be measured adjacent to the application site on the upwind side, immediately prior to application. For ground boom applications, apply using a nozzle height of no more than 4 feet above the ground or crop canopy. For airblast applications, turn off outward pointing nozzles at row ends and when spraying the outer two (2) rows. To minimize spray loss over the top in orchard applications, spray must be directed into the canopy.

Aerial Applications:

The spray boom should be mounted on the aircraft so as to minimize drift caused by wing tip vortices. The minimum practical boom length should be used, and must not exceed 75% of the wing span or 80% rotor diameter. Flight speed and nozzle orientation must be considered in determining droplet size. Spray must be released at the lowest height consistent with pest control and flight safety. Do not release spray at a height greater than 10 feet above the crop canopy unless a greater height is required for aircraft safety. When applications are made with a cross-wind, the swath will be displaced downwind. The applicator must compensate for this displacement at the downwind edge of the application area by adjusting the path of the aircraft upwind. Making applications at the lowest height that is safe reduces the exposure of the droplets to evaporation and wind.

Wind Speed Restrictions:

Drift potential increases at wind velocities of less than 3 mph (due to inversion potential) or more than 10 mph. However, many factors, including droplet size, canopy and equipment specifications determine drift potential at any given wind speed. Only apply this product if the wind direction favors on-target deposition. Do not apply when wind velocity exceeds 15 mph and avoid gusty and windless conditions. Risk of exposure to sensitive aquatic areas can be reduced by avoiding applications when wind direction is toward the aquatic area.

Restrictions During Temperature Inversions:

Do not make ground applications during temperature inversions. Drift potential is high during temperature inversions. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain close to the ground and move laterally in a concentrated cloud. Temperature inversions are characterized by stable air and increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by mist or ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source. Smoke that layers and moves laterally near the ground surface in a concentrated cloud (under low wind conditions) indicated an inversion, while smoke that moves upward and rapidly dissipates indicated good vertical mixing.

MIXING INSTRUCTIONS

COMPATIBILITY

NNI-0001 480 SC is physically and biologically compatible with many registered pesticides and fertilizers or micronutrients. When considering mixing NNI-0001 480 SC with other pesticides, or other additives, first contact your supplier for advice. For further information, contact your local Bayer Representative. If you have no experience with the combination you are considering, you should conduct a test to determine physical compatibility. To determine physical compatibility, add the recommended proportions of each chemical with the same proportion of water, as will be present in the chemical supply tank, into a suitable container, mix thoroughly and allow to stand for five minutes. If the combination remains mixed, or can be readily re-mixed, the mixture is considered physically compatible.

ORDER-OF-MIXING

NNI-0001 480 SC may be used with other recommended pesticides, fertilizers and micronutrients. The proper mixing procedure for NNI-0001 480 SC alone or in tank mix combinations with other pesticides is:

- 1) Fill the spray tank 1/4 to 1/3 full with clean water;
- 2) While recirculating and with the agitator running, add any products in PVA bags (**See Note**). Allow time for thorough mixing;
- 3) Continue to fill spray tank with water until 1/2 full;
- 4) Add any other wettable powder (WP) or water dispersible granule (WG) products;
- 5) Add the required amount of NNI-0001 480 SC, and any other "flowable" (FL or SC) type products;
- 6) Allow enough time for thorough mixing of each product added to tank;
- 7) If applicable, add any remaining tank mix components: emulsifiable concentrates (EC), fertilizers and micronutrients.
- 8) Fill spray tank to desired level and maintain constant agitation to ensure uniformity of spray mixture.

NOTE: Do not use PVA packets in a tank mix with products that contain boron or release free chlorine. The resultant reaction of PVA and boron or free chlorine is a plastic that is not soluble in water or solvents.

ROTATIONAL CROP STATEMENT

Treated areas may be replanted with any crop specified on this label as soon as practical following the last application.

ROTATIONAL PLANT-BACK INTERVALS¹

Immediate plant-back: Brassica (Cole) Leafy Vegetables, Corn (Field, Pop, and Sweet), Cotton, Cucurbit Vegetables, Fruiting Vegetables, Leafy Vegetables (except Brassica), Okra, Tobacco

30-Day plant-back: Alfalfa, Barley, Buckwheat, Clover, Grasses, Millet (pearl), Millet (proso), Oats, Root Crops (Root, Tuber, and Bulb Vegetables), Rye, Sorghum, Soybeans, Teosinte, Triticale, Wheat

9-Month plant-back: All other crops

¹Cover Crops for soil building or erosion control may be planted at any time, but do not graze or harvest for food or feed.

FIELD CROPS

Recommended Applications: Apply specified dosage of NNI-0001 480 SC as needed for control. For best results, treatment should be made when insect populations begin to build and before a damaging population becomes established. Rate selected for use should depend on stage of pest development at application, pest infestation level, plant size and density of plant foliage. Thorough coverage of plant foliage is recommended for optimum product performance. NNI-0001 480 SC may be applied by air, ground equipment or through overhead irrigation systems as designated in the CHEMIGATION statement in the Application Recommendations section of this label. Please contact your local Bayer CropScience representative or Pest Control Advisor for specific recommendations by crop.

CORN (FIELD CORN, POP CORN, SWEET CORN, and CORN GROWN FOR SEED)

PESTS CONTROLLED	RATE PER APPLICATION fluid ounces/Acre
Armyworms (including beet, fall, yellowstriped, and true)	1.0 - 3.0
Black cutworm	
Corn earworm	
European corn borer	
Southwestern corn borer	
Western bean cutworm	

Notes

- Do not enter or allow entry into treated areas during the restricted entry interval (REI) of 12 hours.
- Pre-harvest Interval (PHI): Green forage and silage - **1 day**; Sweet corn - **1 day**; Grain or stover - **28 days**.
- Do not apply more than **3 fl oz per acre (0.094 lb ai/A) per 3-day interval**.
- Do not apply more than **12.0 fl oz per acre (0.375 lb ai/A) per crop season**.
- Do not apply more than **4 times per crop season**.
- Minimum application volume: 10.0 GPA - ground, 5.0 GPA - aerial application.
- Application should be timed to coincide with early threshold level in a developing larval population.
- See CHEMIGATION statement in *Application Guidelines* section of this label.

COTTON

PESTS CONTROLLED	RATE PER APPLICATION fluid ounces/Acre
Armyworms (including beet, fall, yellowstriped, and true)	
Cotton leafworm	1.0 - 2.0
Cotton leaf perforator	
Loopers (including cabbage and soybean)	
Saltmarsh caterpillar	
Cotton bollworm	2.0 - 3.0
Tobacco budworm	

Notes

Do not enter or allow entry into treated areas during the restricted entry interval (REI) of 12 hours.

Pre-harvest Interval (PHI): **28 days.**

Do not apply more than **3.0 fl oz per acre (0.094 lb ai/A) per 5-day interval.**

Do not apply more than **9.0 fl oz per acre (0.262 lb ai/A) per crop season.**

Do not apply more than 3 times per crop season.

Minimum application volume: 10.0 GPA – ground; 5.0 GPA – aerial application.

Application should be timed to coincide with early threshold level in a developing larval population.

See CHEMIGATION statement in *Application Guidelines* section of this label.

TOBACCO

PESTS CONTROLLED	RATE PER APPLICATION fluid ounces/Acre
Tobacco budworm	1.0 - 3.0
Tobacco hornworm	

Notes

Do not enter or allow entry into treated areas during the restricted entry interval (REI) of 12 hours.

Pre-harvest Interval (PHI): **14 days.**

Do not apply more than **3 fl oz per acre (0.094 lb ai/A) per 5-day interval.**

Do not apply more than **12.0 fl oz per acre (0.375 lb ai/A) per crop season.**

Do not apply more than 4 times per crop season.

Minimum application volume: 10.0 GPA – ground; 5.0 GPA – aerial application

Application should be timed to coincide with early threshold level in a developing larval population.

See CHEMIGATION statement in *Application Guidelines* section of this label.

TREE FRUIT, NUT, AND VINE CROPS

Recommended Applications: Apply specified dosage of NNI-0001 480 SC as needed for control. For best results, treatment should be made when insect populations begin to build and before a damaging population becomes established. Recommended application rates within this label are based on full-size mature trees and vines. Thorough coverage of plant foliage and fruit is recommended for optimum product performance. Please contact your local Bayer CropScience representative or Pest Control Advisor for specific recommendations by crop.

POME FRUIT

Crops of Crop Groups 11 including: Apple, Crabapple, Loquat, Mayhaw, Pear, Oriental pear, Quince

PESTS CONTROLLED	RATE PER APPLICATION
	fluid ounces/Acre
Codling moth (West of the Rockies)	5.0
<i>For use against low to moderate infestations in conjunction with alternate control measures such as in established mating disruption blocks.</i>	
Codling moth (East of the Rockies)	3.0 - 5.0
Eyespotted bud moth	
Green fruitworm	
Lacanobia fruitworm	
Leafrollers (including obliquebanded, pandemic, redbanded, and variegated)	
Lesser appleworm	

Notes

- Do not enter or allow entry into treated areas during the restricted entry interval (REI) of 12 hours.
- Pre-Harvest Interval (PHI): **14 days.**
- Do not apply more than **5.0 fl oz per acre (0.156 lb ai/A) per 7-day interval.**
- Do not apply more than **15.0 fl oz per acre (0.468 lb ai/A) per crop season.**
- Do not apply more than 3 times per crop season.
- Minimum application volumes: 100 GPA – ground application. Aerial application is prohibited.
- Application should be timed to coincide with early threshold level in a developing larval population.

STONE FRUIT

Crops of Crop Group 12 including: Apricot, Cherry (sweet and tart), Nectarine, Peach, Plum (includes Chickasaw plum, Damson plum, and Japanese plum), Plumcot, Prune (fresh and dried).

PESTS CONTROLLED

RATE PER APPLICATION

fluid ounces/Acre

Green fruitworm

2.0 - 4.0

Leafrollers (including obliquebanded, pandemic, redbanded, and variegated)

Notes

Do not enter or allow entry into treated areas during the restricted entry interval (REI) of 12 hours.

Pre-Harvest Interval (PHI): 7 days.

Do not apply more than 4.0 fl oz per acre (0.125 lb ai/A) per 7-day interval.

Do not apply more than 12.0 fl oz per acre (0.375 lb ai/A) per crop season.

Do not apply more than 3 times per crop season.

Minimum application volumes: 50 GPA – ground application. Aerial application is prohibited.

Application should be timed to coincide with early threshold level in a developing larval population.

TREE NUT CROPS

Crops of Crop Group 14 including: Almond, Beech Nut, Brazil Nut, Butternut, Cashew, Chestnut, Chinquapin, Filbert, Hickory Nut, Macadamia Nut, Pecan, Walnut (black and English)

PESTS CONTROLLED

RATE PER APPLICATION

fluid ounces/Acre

Fall webworm

2.0 - 4.0

Hickory shuckworm

Naval orangeworm

Peach twig borer

Pecan nut casebearer

Walnut caterpillar

Notes

Do not enter or allow entry into treated areas during the restricted entry interval (REI) of 12 hours.

Pre-Harvest Interval (PHI): 14 days.

Do not apply more than 4.0 fl oz per acre (0.125 lb ai/A) per 7-day interval.

Do not apply more than 12.0 fl oz per acre (0.375 lb ai/A) per crop season.

Do not apply more than 3 times per crop season.

Minimum application volumes: 50 GPA – ground application. Aerial application is prohibited.

Application should be timed to coincide with early threshold level in a developing larval population.

11/12

GRAPE

Including American bunch grape, Muscadine grape, and Vinifera grape

PESTS CONTROLLED	RATE PER APPLICATION
	fluid ounces/Acre
Cutworm	2.0 - 4.0
Grape Leafroller	
Grape leaf skeletonizer	
Omnivorous leafroller	
Orange tortrix	

Notes

Do not enter or allow entry into treated areas during the restricted entry interval (REI) of 12 hours.

Pre-Harvest Interval (PHI): **7 days.**

Do not apply more than **4.0 fl oz per acre (0.125 lb ai/A) per 5-day interval.**

Do not apply more than **12.0 fl oz per acre (0.375 lb ai/A) per crop season.**

Do not apply more than 3 times per crop season.

Minimum application volumes: 50 GPA – ground application. Aerial application is prohibited.

Application should be timed to coincide with early threshold level in a developing larval population.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal

PESTICIDE STORAGE

Do not store for more than 30 consecutive days at an average daily temperature exceeding 100° F. If allowed to freeze, shake well to ensure the product is homogenous before use. Store in original container and out of the reach of children, preferable in a locked storage area. Avoid cross contamination with other pesticides.

PESTICIDE DISPOSAL

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling, if available, or puncture and dispose of in a sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

12
12

IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Bayer CropScience. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Bayer CropScience is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

LIMITATIONS OF LIABILITY: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER CROPSCIENCE'S ELECTION, THE REPLACEMENT OF PRODUCT.

NET CONTENTS:

[----- is a registered trademark of Bayer.]

PRODUCED FOR

Bayer CropScience LP
P.O. Box 12014, 2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709
1-866-99BAYER (1-866-992-2937)
<http://www.bayercropscience.us>

NNI-0001 480 SC (PENDING) Changes Made 07-24-08