Exhibit 49



FIFRA 2(ee) RECOMMENDATION

For Alternative Application Methods – **Split Applications**

FIFRA 2ee RECOMMENDATION

DACTHAL® Flowable Herbicide

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATES OF GEORGIA and NORTH CAROLINA

EPA Reg. No.: 5481-487

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

THIS BULLETIN MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF APPLICATION.

Before applying **DACTHAL Flowable Herbicide** please read the complete label affixed to the product container. All applicable directions, restrictions and precautions on the EPA-registered label are to be followed. Use of **DACTHAL Flowable Herbicide** according to this supplemental labeling is subject to the use precautions and limitations imposed by the label on the container.

IMPORTANT: Read the entire Directions for Use and the Limited Warranty and Disclaimer on the EPA-registered **DACTHAL Flowable Herbicide** label before using this product.

Use Directions

Crop	When to Apply	Rate per acre	Remarks
Onion [dry bulb onions, dry bulb shallots and green onions (green onions, leeks, spring onions or scallions, Japanese bunching onions, green shallots or green eschalots) (dry bulb onions)]	At seeding or transplanting and/or at layby.	Maximum cumulative rate of 10 pints/Acre	Apply DACTHAL Flowable at 3-4 pints/A before onion emergence. A sequential application of DACTHAL Flowable at 3-4 pints/A may be applied 10-14 days later. Apply uniformly to the soil as a spray. DACTHAL Flowable can be sprayed directly over transplants without injury. A layby application can be made on onions either alone or in addition to a DACTHAL Flowable preemergence treatment (with a minimum of 14 days between treatments) up to 14 weeks after planting at rates up to 14 pints/A on any soil type. If weeds emerge prior to layby, the onions should be cultivated or weeded prior to application. Use ½ inch of overhead irrigation to incorporate DACTHAL Flowable into the soil. Where onion emergence is expected to be slow due to cold and wet soil conditions, delay application until seeds begin to germinate to reduce the risk of crop injury. Preplant incorporation is not recommended for onions.

©2020 AMVAC Chemical Corporation. A wholly owned subsidiary of American Vanguard Corporation. All Rights Reserved. American Vanguard, AMVAC, the AMVAC logo, and DACTHAL are trademarks owned by AMVAC Chemical Corporation.

Registrant:

AMVAC Chemical Corporation 4695 MacArthur Court, Suite 1200 Newport Beach, CA U.S.A. 1-888-462-6822