

1
2
3
4 Dear Administrator Jackson,
5

6 The Clean Air Scientific Advisory Committee (CASAC) Particulate Matter (PM)
7 Review Panel met on March 10 – 11, 2010 to review the *Quantitative Health Risk*
8 *Assessment for Particulate Matter – Second External Review Draft* (February 2010) and
9 *Particulate Matter Urban-Focused Visibility Assessment – Second External Review Draft*
10 (January 2010). This letter provides our main comments concerning the *Quantitative*
11 *Health Risk Assessment for Particulate Matter* (RA). CASAC’s responses to the charge
12 questions, along with comments from individual Panel members follow.
13

14 The Second Draft RA presents quantitative assessments of PM-associated risks to
15 support judgments that will be made in the Policy Assessment (PA) with regard to
16 recommendations for the PM National Ambient Air Quality Standard (NAAQS). The
17 RA provides estimates of the degree and nature of risk reductions associated with various
18 population exposure scenarios, corresponding to alternative suites of annual and 24-hour
19 PM_{2.5} standards for the 15 selected urban case study areas. Toward that end, the Second
20 Draft has largely accomplished its objective of providing input to the PA. The Second
21 Draft has been substantially improved in response to comments provided by CASAC
22 following our review of the first draft. The rationale, methods and results for the
23 assessment are well described. The discussion and clarifications regarding uncertainty
24 and variability and the characterization of confidence in the core risk estimates markedly
25 enhance the document. The enhanced discussion of the interplay of annual and 24-hour
26 design values together with patterns in PM_{2.5} monitoring helps to interpret patterns of risk
27 reduction in study areas. With some additional changes recommended here, the RA will
28 be a solid foundation for the PA.
29

30 While the Second Draft has appropriately justified EPA’s decision not to estimate
31 risks for thoracic coarse PM (PM_{10-2.5}) based on limitations in the health effects data and
32 the PM_{10-2.5} monitoring network, we encourage EPA to remain focused on pursuing the
33 research needed for a quantitative risk assessment of thoracic coarse PM. As mentioned
34 in our letter of November 24, 2009 reviewing the first draft RA, we also encourage the
35 Agency to develop the capacity to conduct a formal quantitative exposure assessment for
36 future revisions of the NAAQS.
37

38 As part of simulating air quality standards in the chapter on Urban Case Study
39 Analysis Methods (Chapter 3), EPA uses three different approaches for simulating just
40 meeting the current and alternative suites of PM_{2.5} standards: the proportional approach,
41 the hybrid approach, and the peak-shaving approach. CASAC recommends sharpening
42 the descriptions of these three alternatives. We suggest a graphical and/or mathematical
43 approach, along with the inclusion of examples based on the 15 urban study areas. EPA
44 might also consider discussing how these three approaches relate to potential control
45 scenarios. The hybrid approach merits the greatest emphasis, while the proportional and
46 peak-shaving approaches represent bounding scenarios. As a suggestion on

1 nomenclature, an alternative label for the "peak shaving" approach would be preferable,
2 since this term does not adequately describe the actual method.
3

4 The Second Draft RA includes a new chapter which provides a national scale
5 assessment of long-term mortality related to PM_{2.5} exposure (Chapter 5). In addition to
6 providing a national estimate, this chapter identifies where the urban case study areas fall
7 along the distribution of national risk. While this chapter is useful for assessing the
8 generalizability of the findings in the 15 urban study areas, its findings are otherwise not
9 central to the risk assessment. Consequently, CASAC recommends that the chapter be
10 moved to an appendix and that discussion of the key findings, summarized in Figure 5-4,
11 be placed at the appropriate points in Chapter 4 (Urban Case Study Results) and in the
12 current Chapter 6 (Integrative Discussion of Urban Case Study Analysis of PM_{2.5}-related
13 Risks).
14

15 The Second Draft estimates risk reductions for different air quality scenarios
16 involving specified values of 24-hr and annual standards. Five alternative sets of
17 standards are considered, with the lowest scenario being an annual standard of 12 µg/m³
18 combined with a 24-hour standard of 25 µg/m³. Two additional scenarios were presented
19 at the March 2010 CASAC meeting with pairings of 10/35 and 10/25 µg/m³. The
20 reduction of the annual standard to 10 µg/m³ showed additional benefits beyond those
21 estimated for the scenarios in the RA. However, it was not clear whether these additional
22 scenarios represented potential revisions to the NAAQS and will be added to the final
23 RA. With the assumption of a linear, no-threshold risk model, further reductions would
24 be anticipated as the annual standard is progressively lowered. We recommend that EPA
25 develop and apply specific criteria for determining the lower-bound exposure
26 concentrations to be considered in the risk assessment. Mounting uncertainty at lower
27 concentrations would be one such reasonable basis. Other relevant considerations
28 include the range of concentrations at which the epidemiological studies have been
29 carried out and the need for consideration of the degree of protection afforded to
30 susceptible populations under various scenarios. The interpretation of the findings should
31 also acknowledge the controlling influence of the annual average concentration as the
32 proposed scenario values are progressively lowered.
33

34 On a related matter, CASAC recommends greater clarity as to the basis for
35 selecting the exposure scenarios included in the RA. Are the potential suites of standards
36 or exposure concentration scenarios intended to guide selection among alternative
37 revisions to the NAAQS? As the new overall approach to NAAQS revision is
38 implemented, CASAC and the EPA need to address the extent to which the RA and the
39 PA should be linked.
40

41 In response to CASAC's previous advice, EPA provided an integrated discussion
42 of risk-related analyses that draws on uncertainty and variability analyses as well as the
43 national-scale analyses completed as part of this assessment. This integrated discussion
44 is presented in Chapter 6, arguably the most critical chapter of the document. EPA has
45 captured the key policy-relevant questions and appropriately characterized the
46 uncertainty and variability associated with its core risk estimates. While we are in

3-30-10 Deliberative Draft Letter on the Quantitative Health Risk Assessment for Particulate Matter (Second External Review Draft). DO NOT CITE OR QUOTE.

1 general agreement with EPA’s interpretations of urban study area results as well as its
2 observations on the degree of risk reduction likely to be associated with alternative suites
3 of standards, given its importance, we recommend a careful rewriting and editing of
4 Chapter 6 to ensure that its findings are clearly presented and that unneeded detail is
5 removed.

6
7 CASAC thanks the Agency for the opportunity to provide advice on the Risk
8 Assessment. We look forward to continued discussions with the Agency as we review
9 the *Policy Assessment for the Review of the PM NAAQS - First External Review Draft*
10 (March 2010).

11
12
13
14 Sincerely,

15
16
17
18 Dr. Jonathan M. Samet
19 Chair
20 Clean Air Scientific Advisory Committee
21
22
23

- 24 Enclosure A: CASAC Particulate Matter Review Panel Roster
25 Enclosure B: CASAC Responses to Charge Questions
26 Enclosure C: Individual Panelists’ Responses to Charge Questions

1 **Enclosure A**

2 **Clean Air Scientific Advisory Committee**
3 **Particulate Matter Review Panel**

4
5
6 **CHAIR**

7 Dr. Jonathan M. Samet, Professor and Chair, Department of Preventive Medicine,
8 University of Southern California, Los Angeles, CA

9
10 **CASAC MEMBERS**

11 Dr. Joseph Brain, Philip Drinker Professor of Environmental Physiology, Department of
12 Environmental Health, Harvard School of Public Health, Harvard University, Boston,
13 MA

14
15 Dr. Ellis B. Cowling, University Distinguished Professor At-Large Emeritus, Colleges of
16 Natural Resources and Agriculture and Life Sciences, North Carolina State University,
17 Raleigh, NC

18
19 Dr. James Crapo, Professor of Medicine, Department of Medicine , National Jewish
20 Medical and Research Center, Denver, CO

21
22 Dr. H. Christopher Frey, Professor, Department of Civil, Construction and Environmental
23 Engineering, College of Engineering, North Carolina State University, Raleigh, NC

24
25 Dr. Donna Kenski, Data Analysis Director, Lake Michigan Air Directors Consortium,
26 Rosemont, IL

27
28 Dr. Armistead (Ted) Russell, Professor, Department of Civil and Environmental
29 Engineering , Georgia Institute of Technology, Atlanta, GA

30
31 **CONSULTANTS**

32 Dr. Lowell Ashbaugh, Associate Research Ecologist, Crocker Nuclear Lab, University of
33 California, Davis, Davis, CA

34
35 Prof. Ed Avol, Professor, Preventive Medicine, Keck School of Medicine, University of
36 Southern California, Los Angeles, CA

37
38 Dr. Wayne Cascio, Professor, Medicine, Cardiology, Brody School of Medicine at East
39 Carolina University, Greenville, NC

40
41 Dr. David Grantz, Director, Botany and Plant Sciences and Air Pollution Research
42 Center, Riverside Campus and Kearney Agricultural Center, University of California,
43 Parlier, CA

44
45 Dr. Joseph Helble, Dean and Professor, Thayer School of Engineering, Dartmouth
46 College, Hanover, NH

3-30-10 Deliberative Draft Letter on the Quantitative Health Risk Assessment for Particulate Matter (Second External Review Draft). DO NOT CITE OR QUOTE.

1 Dr. Rogene Henderson, Senior Scientist Emeritus, Lovelace Respiratory Research
2 Institute, Albuquerque, NM

3
4 Dr. Philip Hopke, Bayard D. Clarkson Distinguished Professor, Department of Chemical
5 Engineering, Clarkson University, Potsdam, NY

6
7 Dr. Morton Lippmann, Professor, Nelson Institute of Environmental Medicine, New
8 York University School of Medicine, Tuxedo, NY

9
10 Dr. Helen Suh MacIntosh, Associate Professor, Environmental Health, School of Public
11 Health, Harvard University, Boston, MA

12
13 Dr. William Malm, Research Physicist, National Park Service Air Resources Division,
14 Cooperative Institute for Research in the Atmosphere, Colorado State University, Fort
15 Collins, CO

16
17 Mr. Charles Thomas (Tom) Moore, Jr., Air Quality Program Manager, Western
18 Governors' Association, Cooperative Institute for Research in the Atmosphere, Colorado
19 State University, Fort Collins, CO

20
21 Dr. Robert F. Phalen, Professor, Department of Community & Environmental Medicine;
22 Director, Air Pollution Health Effects Laboratory; Professor of Occupational &
23 Environmental Health, Center for Occupation & Environment Health, College of
24 Medicine, University of California Irvine, Irvine, CA

25
26 Dr. Kent Pinkerton, Professor, Regents of the University of California, Center for Health
27 and the Environment, University of California, Davis, CA

28
29 Mr. Richard L. Poirot, Environmental Analyst, Air Pollution Control Division,
30 Department of Environmental Conservation, Vermont Agency of Natural Resources,
31 Waterbury, VT

32
33 Dr. Frank Speizer, Edward Kass Professor of Medicine, Channing Laboratory, Harvard
34 Medical School, Boston, MA

35
36 Dr. Sverre Vedal, Professor, Department of Environmental and Occupational Health
37 Sciences, School of Public Health and Community Medicine, University of Washington,
38 Seattle, WA

39
40 **SCIENCE ADVISORY BOARD STAFF**

41 Dr. Holly Stallworth, Designated Federal Officer, EPA Science Advisory Board Staff
42 Office

1
2
3
4
5
6
7
8
9
10
11
12
13

NOTICE

This report has been written as part of the activities of the EPA's Clean Air Scientific Advisory Committee (CASAC), a federal advisory committee independently chartered to provide extramural scientific information and advice to the Administrator and other officials of the EPA. CASAC provides balanced, expert assessment of scientific matters related to issues and problems facing the Agency. This report has not been reviewed for approval by the Agency and, hence, the contents of this report do not necessarily represent the views and policies of the EPA, nor of other agencies within the Executive Branch of the federal government. In addition, any mention of trade names or commercial products does not constitute a recommendation for use. CASAC reports are posted on the EPA Web site at: <http://www.epa.gov/casac>.

1 **Enclosure B**

2
3 **CASAC Responses to Charge Questions**

4
5 Chapter 3 – Urban Case Study Analysis Methods

6 1) Air quality inputs (section 3.2): We have expanded the consideration of alternative
7 approaches to simulating just meeting the current and alternative suites of PM_{2.5}
8 standards (i.e., rollback approaches) to include a peak shaving approach, in addition to
9 the hybrid and proportional approaches considered in the first draft assessment. This peak
10 shaving approach is intended to represent more localized, rather than regional, patterns of
11 PM_{2.5} reductions (discussed in section 3.2.3.3).

12
13 a) To what extent does the Panel believe that the use of the peak shaving approach
14 provides useful additional exploration of variability associated with how ambient PM_{2.5}
15 concentrations are simulated to change upon just meeting the current and alternative
16 suites of standards?

17
18 **As noted in our overall comments, CASAC recommends that the "peak shaving"**
19 **approach be more clearly defined in a graphical and/or mathematical fashion. We**
20 **also recommend that it be renamed, given the potential to misinterpret its actual**
21 **application. The "peak shaving" approach does provide useful information, but**
22 **specific examples are needed as to the situation(s) and potential control**
23 **approach(es) it is meant to simulate. Appendix B should also provide the specific**
24 **mathematical formulation. It would be instructive to provide an example of the**
25 **three rollback approaches as applied to data for several cities in section 3.2, or in**
26 **Appendix B.**

27
28 b) We have used comparisons of composite monitor annual averages generated using the
29 different rollback approaches as a surrogate for differences in long-term exposure-related
30 mortality in looking across all three rollback approaches. To what extent does the Panel
31 believe that this is a reasonable approach for assessing the impact of variability
32 associated with simulating changes in air quality patterns on estimates of long-term
33 exposure-related mortality?

34
35 **A very brief description of how annual design values are currently calculated should**
36 **be provided to help motivate the procedure developed here. CASAC is**
37 **uncomfortable with the approach used for imputing missing values, and has**
38 **discussed alternatives in the members' individual comments. Whatever method is**
39 **chosen, the impact of the approach taken can be characterized using data from**
40 **monitors where there is little missing data, then removing some data, applying the**
41 **imputation approach, and testing to see how closely the method reproduces the**
42 **original annual average. The approaches to compositing and imputing missing**
43 **values would be more easily followed if equations were provided.**

44
45 2) Selection of model inputs (section 3.3): We have expanded and clarified the discussion
46 of our rationale for identifying modeling choices comprising the core risk model,

1 focusing in particular on selection of C-R functions (section 3.3.3). To what extent does
2 the Panel consider this discussion to be clear and the model selections appropriate?
3

4 **CASAC commends the authors for expanding and clarifying the rationale for**
5 **identifying modeling choices comprising the core risk model in a logical and**
6 **satisfactory manner. Their model selections were appropriate for this review cycle,**
7 **which is focused on PM_{2.5} exposure and cardiovascular responses. The stated**
8 **rationale provides a solid foundation for the selection of the epidemiological studies**
9 **that were utilized to establish C-R functions. The expansion of the discussion and**
10 **integration of the ISA was useful in reinforcing gaps in knowledge, as in the text on**
11 **p. 3-20 stating that there were no multi-city studies for the category of short-term**
12 **exposure to PM_{2.5} and emergency department visits for cardiovascular and/or**
13 **respiratory illnesses. The summary tables (Tables 3-5 through 3-8) provide a useful**
14 **synopsis of the model inputs for the core risk models and sensitivity analyses.**
15

16 Charge Question 3: Addressing uncertainty and variability (section 3.5): We have
17 clarified the process used to evaluate sources of variability and added coverage for
18 specific sources of variability (section 3.5.2); expanded our discussion of the qualitative
19 analysis of uncertainty (section 3.5.3); and included analyses of pair-wise interactions of
20 sources of uncertainty (section 3.5.4). To what extent does the Panel consider these
21 discussions to be clear and appropriate?
22

23 **In general, the second draft RA appropriately identifies and discusses key sources of**
24 **variability and uncertainty, and includes sensitivity analyses that provide insight**
25 **regarding the impact of some sources of uncertainty on the core risk estimates. The**
26 **authors have provided a footnote explaining the rationale for identifying “key”**
27 **sources of variability. The document should indicate if the same process was used to**
28 **identify “key” sources of uncertainty. We note that EPA added material regarding**
29 **co-pollutant concentrations and demographic and socioeconomic status, as**
30 **requested in CASAC comments on the first draft of the RA.**
31

32 **EPA did not address CASAC's recommendation on the first draft RA suggesting**
33 **that “exposure modeling should be included in the REA. A probabilistic Tier 3**
34 **approach should be used for the exposure assessment.” While we understand that**
35 **timing may have precluded adequate treatment of this topic, we urge EPA to**
36 **develop the capacity to do so for future revisions of the NAAQS. CASAC made the**
37 **same recommendation at the time of the last review of the PM NAAQS. With**
38 **regard to uncertainties, in response to CASAC comments on the first draft of the**
39 **REA, EPA has included uncertainty in the C-R function itself, which was developed**
40 **from single studies. EPA has appropriately taken into account differences in C-R**
41 **functional form associated with studies that addressed long-term or short-term**
42 **effects for single or multi-city studies even if they were not the basis for the final set**
43 **of C-R functions used in the RA.**
44

45 **The RA now provides adequate explanation of the uncertainty categories of “low”,**
46 **“medium”, and “high.” However, there is a confusing statement to the effect that**

1 **“high” sources of uncertainty “are likely to influence the interpretation of risk...”**
2 **“if those sources of uncertainty are reduced or more fully characterized.” The**
3 **parenthetical “if” clause seems to confuse the issue, and should be deleted. The RA**
4 **needs to describe the process by which "staff consensus" was achieved (see lines 6-**
5 **20 of p. 3-63). If staff consensus is an element of decision-making in carrying out the**
6 **RA, then this aspect of the process needs to be described.**

7
8 **EPA has adequately commented on the extent to which there are dependencies**
9 **among pairwise combinations of sources of uncertainty, and whether these**
10 **dependencies would tend to offset or to increase the overall range and direction of**
11 **uncertainty in the assessment results. For example, the statistical fit of the C-R**
12 **functions, and the shape of the functions, are inter-related.**

13
14 **Based on quantifiable sensitivity analysis, the report generally clearly conveys that**
15 **the “core” estimates appear to be at the low end of alternative “plausible” estimates.**
16 **However, particularly in Chapter 6, the role of sources of uncertainty treated**
17 **qualitatively should also be addressed. In particular, given exposure**
18 **misclassification, it is likely that the core estimates are biased low and hence**
19 **conservative, an important point to convey consistently.**

20
21 **Charge Question 4: Sensitivity analysis results (section 4.3): We have included a**
22 **discussion of how the results of the sensitivity analysis can be used as an additional set of**
23 **reasonable risk estimates to inform consideration of uncertainty in the core risk estimates**
24 **(see section 4.3.2). What are the Panel’s views on how we have used the sensitivity**
25 **analysis results to support consideration of uncertainty in the core risk estimates?**

26
27 **Overall, the sensitivity analysis section 4.3 is very good and nicely covers a complex**
28 **topic. Table 4-3 is a useful summary. The classification of descriptive categories for**
29 **small, moderate, moderate-large, and large contributions is useful. However, it**
30 **would be more appropriate to refer to these as contributions to “sensitivity” rather**
31 **than “uncertainty.” As noted in several place, the sensitivity analyses represent**
32 **plausible alternatives to the core estimate, but are not probabilistic. The language**
33 **of the RA is appropriate on this point and has clearly articulated that the sensitivity**
34 **analyses results represent plausible and scientifically defensible estimates. The**
35 **range of these estimates provides an indication of the implications of uncertainty.**

36
37 **The evaluation of alternative model structure is critically important, because model**
38 **structure can potentially be a larger source of uncertainty than the range of values**
39 **for an input to a given model. The results in Table 4-3 indicate, for example, that**
40 **the random effects log-log model provides larger risk estimates than the fixed effects**
41 **log-linear model used for the core estimates. This information is very useful and is**
42 **an excellent addition to the RA. The more thorough treatment of model choices and**
43 **alternative C-R functions provides plausible alternative estimates to the core**
44 **estimate.**

45

1 **Per CASAC’s comments on the first draft RA, EPA indicates the direction of the**
2 **percent changes in risk. In addition to the percent difference, the actual difference**
3 **in risk should be reported to provide further context. The second draft RA seems to**
4 **put emphasis on relative changes in risk. However, the NAAQS are intended to be**
5 **protective of public health, and therefore the magnitude of the risk estimates is**
6 **ultimately a more useful policy-relevant metric.**

7
8 **The sensitivity analysis related to peak shaving and “peakiness” was not very clear**
9 **in its interpretation. What are the main points to take away from these analyses?**

10
11 5) Consideration of design values and patterns of PM_{2.5} monitoring data in interpreting
12 core risk estimates (section 4.5): To enhance our interpretation of the patterns of core risk
13 estimates generated for both the current and alternative suites of standards, we have
14 included analyses of 24-hour and annual design values together with patterns of PM_{2.5}
15 monitoring data for the 15 urban study areas. This reflects the fact that these two factors
16 play a key role in determining the degree of risk reduction estimated upon just meeting
17 the current and alternative suites of standards under alternative rollback approaches. As
18 part of the consideration of design values, we have also contrasted the 15 urban study
19 areas with patterns of design values seen for the broader set of urban areas in the U.S. in
20 order to help place the urban study area in a broader national context.

21
22 a) To what extent is the Panel supportive of these additional assessments?

23
24 b) Does the Panel have any recommendations for additional insights based on
25 consideration of patterns in design values and PM_{2.5} monitoring data across the 15 urban
26 study areas and at the national level?

27
28 **The graphical presentations depicting the 24-hour and annual average design values**
29 **for US urban areas and the 15 urban study areas used in the RA were very helpful**
30 **for understanding the concept of the controlling standard, the implications of**
31 **reducing either the annual or the 24-hour standard, or both, and the**
32 **representativeness of the 15 urban study areas. To further enhance the value of**
33 **these graphical presentations, more complex color coding could be used to provide**
34 **information on the US region of each urban study area. In these plots, unsupported**
35 **conclusions were drawn for cities lying on the border between zones. Graphical**
36 **presentation of the design values by monitoring site for the 15 urban study areas**
37 **provides valuable insights into the role of patterns of PM monitoring data in**
38 **different cities in determining consequences of various control strategies. We**
39 **recommend that the term “peaky” as used to describe PM patterns be better**
40 **defined and applied consistently. The main observations from both sets of plots**
41 **should be summarized, especially as regards impacts of the alternative rollback**
42 **procedures.**

43
44 **Chapter 5 of the revised RA provides estimates of the numbers of deaths**
45 **attributable to long-term PM_{2.5} exposure, based on air quality estimates from the**
46 **Community Model for Air Quality (CMAQ) and the environmental Benefits**

1 **Mapping and Analysis Program (BenMap), and uses the risk estimates derived for**
2 **the Krewski, 2009 assessment of the ACS data with a LML of 5.8 ug/m³. A**
3 **principal purpose for inclusion of this chapter is to place the PM_{2.5}-associated risks**
4 **for the 15 urban study areas within the distribution of risks nationally. Figure 5-4**
5 **provides the key findings in regard to this purpose and indicates that the selected**
6 **urban study areas in large part fall in the highest 20% of the distribution of sites. .**
7 **We recommend that Chapter 5 be moved to an Appendix with inclusion of Figure 5-**
8 **4 at appropriate points within Chapter 4 and the current Chapter 6. The figure**
9 **provides information relevant to the generalizability of findings from the 15 areas to**
10 **the entire United States. However, the estimates themselves are not directly**
11 **relevant to the overall purpose of the RA; the estimation approach differs from that**
12 **used for the 15 urban study areas; and the chapter is brief and does not adequately**
13 **set out sources of uncertainty and variability. By placing the chapter's contents into**
14 **an appendix and specifically acknowledging its purpose, it will not distract from the**
15 **flow of the RA and the major objective of the analysis will be met by inclusion of**
16 **Figure 5-4 in Chapter 4 and the revised Chapter 5.**

17
18 Chapter 6 – Integrative Discussion of PM_{2.5}-related Risks

19 6) We have developed an integrated discussion of the PM_{2.5}-related risk estimates which
20 considers the results of the qualitative and quantitative treatment of uncertainty and
21 variability together with the various national-scale assessments completed for the analysis
22 to support interpretation of the core risk estimates. As part of the integrative discussion,
23 we also provide key observations that bear on policy-relevant risk-based questions.
24

25 **CASAC was unanimously pleased by the addition of this chapter to the document.**
26 **The chapter summarizes the many detailed analyses carried out in Chapter 4 and**
27 **extended in the Appendices. However, the results presented could be summarized**
28 **more effectively and rather than being presented in great detail, there should be a**
29 **more integrative discussion. The choice of the 15 urban study areas was previously**
30 **discussed in Chapter 4. Similarly, the choices of endpoints were already**
31 **documented. Clearly, the use of IHD mortality (as opposed to all- causes of**
32 **cardiovascular mortality) represents an upper bound of effects. This outcome was**
33 **apparently selected because of its availability in the Krewski et al. 2009 analysis.**
34 **CASAC discussed the selection of alternative scenarios under the assumption of a**
35 **linear, no-threshold C-R function. In the current draft RA, the lowest levels**
36 **assessed were 12/25 µg/m³; however, preliminary analyses at 10/25 µg/m³ were**
37 **presented, indicating a variable effect across the 15 urban study areas, resulting**
38 **from different characteristics of some of the areas. As mentioned previously,**
39 **specific criteria need to be developed and applied in selecting the lower bound**
40 **scenarios. These scenarios will be particularly informative for considering**
41 **protection of susceptible populations as uncertainty increases at lower bound**
42 **concentrations. There are a number of potential bases for the specification of these**
43 **scenarios including the concentrations at which observations have been made, the**
44 **level of uncertainty, and the extent of protection to be achieved.**
45

1 a) To what extent does the Panel believe that we have captured the key policy-relevant
2 questions that can be addressed by this risk assessment?
3

4 **EPA has presented and captured the key-policy relevant questions. However,**
5 **CASAC continues to recommend a qualitative discussion on PM_{10-2.5} and on those**
6 **effects of PM_{2.5} for which the evidence was found to be “suggestive”.**
7

8 b) We provide a set of key observations related to estimates of risk associated with
9 simulations of just meeting the current and alternative suites of standards. These
10 observations are based not only on consideration of trends in risk reduction across
11 alternative suites of standards and residual risk remaining after simulation of just meeting
12 specific suites of standards, but also on additional factors that can impact risk (e.g., the
13 role of annual and 24-hour design values, the peakiness of PM_{2.5} distributions within a
14 study area, and application of different rollback approaches). To what extent do the Panel
15 members believe that the observations presented in section 6.2 are well supported by the
16 results of the analyses? Are there other observations that might be made that would help
17 to address the policy-relevant questions identified at the beginning of the chapter?
18

19 **Section 6.2 summarizes the findings of key analyses. The sensitivity analyses**
20 **presented explore multiple factors including the roll back approach, concentration**
21 **scenarios, and location, the latter influencing findings through the the PM_{2.5}**
22 **concentration profile and mortality rate. The text describing these key observations**
23 **is too long and not sufficiently structured. While the analyses do support the**
24 **observations made, the text does not make these linkages with sufficient clarity. In**
25 **fact, much of the discussion becomes too anecdotal as individual cities and scenarios**
26 **are reviewed.**
27

28 **Key observations are presented and adequately discussed with regard to the roles of**
29 **annual and 24-hour design values in determining population risk and of the role of**
30 **“peakiness” of distributions. The text should more sharply characterize the critical**
31 **finding of the cross-area variation in the extent of risk reduction under the various**
32 **scenarios. This variation needs emphasis as extension of the findings from the 15**
33 **urban study areas to the entire country is considered.**
34

35 c) Part of our interpretation of the core risk estimates presented in section 6.2 is our
36 characterization of confidence in the core risk estimates and in observations made based
37 on those estimates. These assessments of confidence are based on consideration of the
38 results of the sensitivity analysis as well as on the qualitative assessment of uncertainty
39 and variability. To what extent does the Panel believe that the characterizations of
40 confidence in the core risk estimates and associated policy-related observations are
41 reasonable given available information?
42

43 **See above. The RA reasonably and appropriately describes the level of confidence**
44 **that can be given to its analyses, given sources of uncertainty and variability.**
45

1 d) As part of the integrative discussion, we use the results of several national-scale
2 analyses (i.e., the national scale PM_{2.5} mortality analysis, the representativeness analysis,
3 and the new exploration of design values and patterns of PM_{2.5} monitoring data
4 presented in section 4.5) to place the results of the risk assessment in a broader national-
5 context. What are the Panel members' views on appropriateness of this effort to place
6 results of the analysis in a national context?
7

8 **The RA presents multiple analyses that explore issues related to the extension of the**
9 **findings from the 15 urban study areas to a national-scale. These areas were**
10 **admittedly not chosen as a representative sample per se, but to be illustrative of**
11 **locations with different patterns of PM_{2.5} concentrations. The analyses presented in**
12 **sections 4.4 and 4.5 and in Chapter 5 satisfactorily set out a basis for placing the RA**
13 **results in a national context. Figure 5.4 is also valuable, and should be included in**
14 **the revised Chapter 6.**
15

16 e) We conclude chapter 6 with a list of key observations. Does the Panel believe that we
17 have appropriately highlighted key findings of the risk assessment in these observations?
18 Of particular note is the observation that, while alternative 24-hour standard levels can be
19 used to reduce annual-average PM_{2.5} concentrations and thus to reduce estimated risk,
20 the results are likely to be highly variable across urban areas. More consistent lowering of
21 annual-average PM_{2.5} concentrations across study areas, and thus more consistent
22 reductions in estimated risk, may result from application of alternative annual standard
23 levels. We also note that simulation of the alternative 24-hour standard level of 25 µg/m³
24 resulted in reductions in annual-average PM_{2.5} levels for some study areas that were well
25 below the lowest annual standard level assessed (i.e., below 12 µg/m³). As a
26 consequence, we observed risk reductions reflecting these changes in annual-average
27 PM_{2.5} levels below 12 µg/m³. Given these results, does the Panel believe that there is
28 utility in estimating risks for alternative annual standard levels below 12 µg/m³?
29

30 **Key observations are presented in a balanced and fair way. Staff acknowledges that**
31 **the range of effects are in two categories: 3-9% and 0-3% in two halves of the**
32 **country. This may be less precise than what the data indicate in that it would**
33 **appear from their own estimates that the bulk of the effect comes from the upper**
34 **end of the exposure in the counties (pg 5.8, line 12-15).**
35

1 **Enclosure C**

2
3
4
5
6
7
8

*Compendium of Individual Preliminary Panelists Comments
Quantative Health Risk Assessment for Particulate Matter (February 2010)*

9	Comments from Mr. Ed Avol	15
10	Comments from Dr. Joe Brain	22
11	Comments from Dr. Wayne Cascio	25
12	Comments from Dr. Christopher Frey	26
13	Comments from Dr. Philip Hopke	34
14	Comments from Dr. Morton Lippmann	36
15	Comments from Dr. Robert Phalen	41
16	Comments from Dr. Ted Russell	42
17	Comments from Dr. Frank E. Speizer	45
18	Comments from Dr. Helen Suh:	49
19	Comments from Dr. Sverre Vedal:	50

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45

Comments from Mr. Ed Avol

General:

The second external review draft of the Quantitative Health Risk Assessment for Particulate Matter is a dramatic and substantial improvement over previous versions, and sets a new thoughtful foundation and prototype for subsequent risk assessments to generally follow. Improved presentations, superior figures, and useful tables and footnotes are present to guide the reader through the process by which EPA staff and consultants have reviewed available data, performed additional analyses, and interpreted the results of those analyses. The NAAQS health risk assessment process is becoming increasingly transparent and trackable (to the benefit of the reader and the public), but some convolutions and meanderings still persist to prevent complete clarity of presentation and focused presentations of conclusions, decisions, and positions. Chapter 4 provided a detailed step-by-step examination of the approach, the analyses, and the results in appropriate depth. Chapter 5 seemed minimal and uncertain as to whether the authors wanted to present it in the main body of the risk assessment or in an appendix. The addition of Chapter 6 as an opportunity to present an integrated view of the work performed is much-appreciated and was generally well-presented.

Charge Questions to CASAC:

1a) Utility of the peak-shaving approach:

The peak shaving approach seems useful and warranted, and aids in understanding the impact and effects of various reduction approaches for a range of possible 24hr/annual standards.

1b) Use of the composite monitor approach:

This may well be the appropriate approach to take, but it would be more convincing if some supporting data for this approach were provided in the appendix or main body of the document. This could readily be done using sites in several of the urban study areas with complete or near-complete data, and dropping out one or more data points, creating the “need” for fill-in strategies and a ready comparison with the actual (observed) data.

2) Selection of model inputs:

This seemed generally clear and supported. The justified emphasis on PM_{2.5} and cardiovascular endpoints may be appropriate at this point in the NAAQS review cycle, but ongoing concern about other particle sizes and constituents, as well other health endpoints, makes the current specific approach time-limited and in need of likely review in successive cycles.

3) Addressing uncertainty and variability:

The discussion and clarifications regarding uncertainty and variability are much-improved, much-appreciated, and markedly enhance the credibility and stature of the entire document. The presentations are definitely appropriate and mostly clear.

3-30-10 Deliberative Draft Letter on the Quantitative Health Risk Assessment for Particulate Matter (Second External Review Draft). DO NOT CITE OR QUOTE.

1 4) In general, the approach seemed appropriate and well-documented. Some assumptions
2 appear to have been made (or at least, not justified in the main body of the presentation),
3 such as the consideration of Philadelphia and Los Angeles as “representative” East and
4 West Coast urban areas, respectively, for sensitivity analyses.

5
6 5a) The approach to include both 24hr and annual design values in assessing core risk
7 estimates is a reasonable one that helps to clarify differences between reduction strategies
8 that may not otherwise have been apparent. Expansion of the discussion to 15 urban
9 study areas is a worthwhile endeavor that yields a number of insights and benefits (not
10 the least of which is added credibility to the calim of national representativeness). The
11 approach was generally well-done, added to the discussion, and enhanced the validity of
12 the presentation. Several specific questions remain (see detailed queries below) but in
13 general, I am wholly supportive of the presented approach and applaud the staff and
14 consultants for their work.

15
16 5b) (No additional insights or recommendations at this time)

17
18 6a) The key policy-relevant questions have been captured in the presentation.

19
20 6b) The observations in Section 6.2 are well-supported but not always clearly articulated.

21
22 6c) The characterization of confidence in the core risk estimates is a marked
23 improvement over previous versions of the document, and helps to establish the
24 credibility of the overall approach. Given the available information, the presentation
25 appears appropriate.

26
27 6d) The presentation of the placement of the risk assessment results in to a national
28 context appeared a bit tentative and unsure. A more forceful presentation and assertion
29 about representativeness could and should be made.

30
31 6e) The key observations listed at the end of Chapter 6 are generally present, but not
32 presented in as clear or concise a manner as could be done. I believe it is useful to
33 present estimated risks for a range of standards and levels, so that the Administrator has a
34 more complete perspective of the possible implications of the various reduction
35 strategies. However, I don't think it is worth a great deal of effort or time to estimate risk
36 reduction at 6 or 8 or 10 ug/m³ annual average. Practically speaking, there are US urban
37 areas unlikely to ever get those low levels, so with a uniform national standard, it seems
38 more of an academic exercise.

39
40
41 Questions:

42 P3-11, lines 30 on (the composite monitor approach) – The procedure is described for
43 filling in missing data, but there is not presentation or reference to any validation of how
44 precise, accurate, or appropriate the fill-in approach is, based on more complete data at
45 sites within the respective urban study areas. Was any assessment of the efficacy of this
46 approach done? The comment in the text (p3-13, lines 7-11 that in New York, for

3-30-10 Deliberative Draft Letter on the Quantitative Health Risk Assessment for Particulate Matter (Second External Review Draft). DO NOT CITE OR QUOTE.

1 example, 2/3 of the data was interpolated may raise some cause for concern that could be
2 allayed by providing some sample analyses with existing and retrievable data.

3
4 P4-28, line 18, sensitivity analysis modeling using Philadelphia and Los Angeles as
5 “representative” East and West Coast urban areas...how was this decision made? On
6 what basis or comparison criteria? How “representative” are either of these urban areas of
7 their respective coastal counterparts? Some explanation or justification should be
8 provided for these selections, even if it is only to refer to an Appendix or footnote.

9
10 P4-49, line2 – the explanation behind this discussion of determinants of PM_{2.5} health
11 estimates is confusing. On p4-48, lines 21-24, the estimates of risk in the risk assessment
12 framework are claimed to be based on four elements: “...population, baseline incidence
13 rates, air quality, and the coefficient rating air quality and the health outcome (i.e, the
14 PM_{2.5} effect estimates)”. On the next page, the determinants of the PM_{2.5} effect estimates
15 are claimed to”... be grouped into three areas: demographics, baseline health conditions,
16 and climate and air quality” (which are actually *four* areas). However, it would appear
17 that these statements argue that the fourth element of the basis for the risk estimates is
18 composed of the first three elements (if population = demographics, baseline incidence
19 rates = baseline health conditions, and air quality = air quality), which seems a bit
20 circular in description and confusing.

21
22 P4-49, line 14 bullet – It is not clear why climate and air quality are included in the same
23 bullet, since the references and variables listed seem to cover each of the climate and air
24 quality categories separately and without overlap.

25
26 P4-51, Table 4-5, Co-Pollutant Levels row – in the text discussion of PM_{2.5} and relevant
27 risk estimates, SO₂ is specifically identified and included in the figures and tables, yet in
28 the table presented national-level data for use in developing risk determinants, ozone is
29 the only co-pollutant presented. It would therefore seem appropriate to note why Ozone
30 is listed (cardiovascular mortality reference?) and why SO₂ is not.

31
32 P4-54, line27 – It is a bit confusing for the reader to follow the discussion from text to
33 table to figures, since the number of urban case study areas plotted vary in the table (4-6)
34 by category, are not presented in any key with the appropriate figures, and don’t appear
35 to be consistent between the table listing and the related figure.

36
37 P4-57, Figure 4-9 through 4-16: (a) It would make it easier for the reader to follow the
38 discussion and the illustrated material in the figures if some additional information were
39 provided in a key or caption with the figures, such as the number (n) for each of the three
40 data categories plotted (all counties CDF, Case study counties CDF, case study counties)
41 and some comment about the splines or plots and inflection points displayed; (b) there is
42 an inconsistency in the presentation of these figures, since some specifically comment on
43 the number of urban case study areas above or below a given percentile (Figures 4-9, 4-
44 11,4-14, 4-15, 4-16), while others have no comment (Figure 4-10, 4-12, 4-13); A
45 summary comment for these three figures would help guide the reader as to the key
46 implications of the provided figure; (c) Would there be additional clarity or power of

3-30-10 Deliberative Draft Letter on the Quantitative Health Risk Assessment for Particulate Matter (Second External Review Draft). DO NOT CITE OR QUOTE.

1 argument gained by somehow highlighting or identifying the plotted values of the 15
2 urban study areas from amongst the larger number shown?

3
4 P4-66, lines 10-12 and P4-67 (Figure 4-17) – The text and figure are inconsistent, since
5 the text says (with respect to the 35/15 plot in Figure 4-17) “...Atlanta, Birmingham, and
6 Houston fall into this zone...” (Zone C), but Figure 4-17 shows Birmingham in Zone B2,
7 Atlanta arguably on the cusp between C and D, and Houston well into Zone D...and in
8 fact, the next bullet in the text (P4-66, lines 13-16) discusses Houston’s presence in Zone
9 D.

10
11 Chapter 5 (National-Scale Assessment of Long-Term Mortality) – This chapter seems a
12 bit tentative. The data presented is critical to the overall discussion, but the manner in
13 which it is presented doesn’t fit with the previous chapter or document flow. The
14 introductory sections are fine and lay the ground for presentation of the results, but the
15 methods section seemed spotty and uncertain in tone as to whether this should be better
16 placed entirely in Appendix G with more details, or presented more conceptually here,
17 with reference to the appropriate appendix for details. The two figures presented are the
18 key information for the chapter. The text provides a description of what is literally
19 presented in the figures, but provides little in the way of interpretation (see text related to
20 Figure 5-3, for example). What is the intended message associated with Figure 5-3? A
21 sentence or two would help to direct readers’ interpretation or consideration of this.

22
23 P5-2, lines 14 thru 21 (interplay between regional-scale reductions in adjacent urban
24 areas) – the issue under discussion and the key problem being identified still is not
25 clear...

26
27 P5-5, line 15 - The reference to “...7 Health Effects Institute PM regions...” seems
28 unnecessary and distracting; this discussion of health impacts relates to geographic
29 regions of the country (southwest, southeast, etc, as presented in Table 3.2, p3-14), and
30 how the Health Effects Institute identifies these areas does not seem germane to the
31 central argument.

32
33 Chapter 6 (Integrative Discussion) offers the promise of pulling together the previous
34 chapters’ presentations into one focused summary for reader review, but up through
35 Section 6.3, too often self-destructs by using meandering phrasing and including
36 extraneous comments in conclusionary statements. The net result of the writing approach
37 employed is to obscure the message and frustrate the reader. There are some excellent,
38 insightful, and important points to be made in this presentation, but they are often hidden
39 in multiple qualifying commentaries when the main point is being presented. Use of
40 bullets in the commentary can be an effective way of focusing the reader on the main
41 persuasive points of the presentation, but in this chapter, their use suffers from a lack of
42 clarity. With each bullet, make the statement, and then provide a terse justification in the
43 ensuing paragraph. For example, the bullet might be “Dramatically different rates of
44 mortality risk reduction are observed in areas with peaky PM_{2.5} distributions, when peak-
45 shaving, compared to proportional rollback , approaches are used.” A brief supporting
46 paragraph could use Salt Lake City or Los Angeles data as an example to make the point.

3-30-10 Deliberative Draft Letter on the Quantitative Health Risk Assessment for Particulate Matter (Second External Review Draft). DO NOT CITE OR QUOTE.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46

P6-7, line 21-22 – This comment is self-obvious and not necessary; of course the variability in incidence estimates is driven by differences in the study population (if the analyses are correctly performed and identifiable confounders are adjusted for). Remove this opening sentence and begin this paragraph with something like, “Substantial variability in incidence estimates was observed for the 15 urban study cases. Substantially less variability would be expected in estimates...”

P6-17, lines 26 thru 29 – This seems like a very odd disclosure comment to make as a closing comment. Under what set of circumstances does EPA believe someone reading this document would interpret the 15 urban areas under discussion being the ONLY 15 areas in the country having health risks associated with PM2.5? This last sentence in the paragraph seems entirely unnecessary, but if there is a sense that this point is worthy of a comment, I would suggest that the last two sentences in this section of the text (lines 24-29) be deleted and replaced with something like the following: “The results of the national-scale mortality analysis suggests that the 15 urban study areas selected for presentation in this document are representative of a continuum of urban area results towards the upper end of the cumulative mortality distribution.”

P6-19, lines 16 thru 23 (the final bullet on the page) – This comment, which establishes the credibility for evaluating the 15 urban study areas chosen, does not directly address the policy –relevant question posed under which it is listed, but does go to a central issue in the considering the relevance of the risk assessments undertaken. Accordingly, this bullet should be moved up to become the first bullet in Section 6.4, establishing the credibility of the study areas used in the analyses, *before* the results of those analyses and their relevance to policy are presented.

Editing Details:

- P2-6, line 28 – delete one of the periods at the end of the sentence
- P2-8, line 27 – something is missing where the “(chapter 5)” placeholder is in the sentence; perhaps “long-term mortality”?
- P3-7, line 34 – need hyphen between “empirically estimated”
- P3-13, line 6 – “synced” is mis-spelled; shouldn’t this be “synched” or “synchronized”?
- P4-17, line 22 – remove underline for “of total incidence”
- P4-17, line 28 – change “these head negative point estimates” to “these had negative...”
- P.4-23, line 13 – change “...2-4hour average risk...” to “24-hour average risk”
- P4-27, line 26 – capitalize “eleven” to become “Eleven”
- P4-38, line 2 – “...morality...” should be “...mortality...”
- P4-40, line 26 - “...at the same urban study are...” should be “...at the same urban study area...”
- P4-41, line 25 - This first sentence anticipates a listing of several factors that are then never presented, so the sentence should be re-structured, perhaps as, “There are several observations regarding key sources of ... that can be made.”
- P4-41, line 28 – change “factor” to “factors”
- P4-42, line 1 – what does “...the *highest* sensitivity analysis results...” mean? Does it mean the biggest changes, or the best outcome? A little re-phrasing would help.

3-30-10 Deliberative Draft Letter on the Quantitative Health Risk Assessment for Particulate Matter (Second External Review Draft). DO NOT CITE OR QUOTE.

- 1 P4-42, line 25 – for consistency with previous line, probably better to say “...log-linear
- 2 model with fixed effects...”
- 3 P4-42, line 28 – remove hyphen from local-sources
- 4 P4-42, line 34 – should be re-phrased to read “...in the area of the curve between the
- 5 LML and the PRB” rather than “...as you move below the LML”
- 6 P4-42, lines 36 to 38 – this is an important comment that, as written, is difficult to
- 7 understand. I would offer the following: “ Due to large uncertainties associated with C-R
- 8 functions in the range between the LML and PRB, such estimates should be excluded
- 9 when considering *reasonable* alternative risks associated with core risk estimates.”
- 10 P4-45, footnote refers to open circles in Figures 4-6 and 4-7, but neither has open circles
- 11 nor displays SO₂ risk estimates; I believe Figure 4-8 is the correct reference.
- 12 P4-46, Figure 4-8 key – add a “g” to “modelin”
- 13 P4-47, line 13 – change phrasing from “...attributable to secondarily formed PM_{2.5}” to
- 14 “...partially attributable to secondary formation of PM_{2.5}”
- 15 P4-47, line 18 – this statement can be more definitive, based on Figure 4-8; Instead of
- 16 “Most if not all”, change to read “Virtually all of the alternative model specifications...”
- 17 P4-47, lines 29, 31, and 32 – Figure references should be to Figures 4-7 and 4-8, not 4-6
- 18 and 4-7.
- 19 P4-49, line2 – shouldn’t this be four areas, not three (demographics, baseline health,
- 20 climate, and air quality)?
- 21 P4-49, line3 – change “..., and climate and air quality...” to “..., climate, and air
- 22 quality...”
- 23 P4-49, line31 – Incorrect table referenced; should be Table 4-5 in current draft version.
- 24 P4-54, line 5 –Kolmogorov- Smirnov test results appear in Table 4-7, not in Table 4-4.
- 25 P4-54, line21 – Figure numbers are incorrect in text.
- 26 P4-70, Table 4-8 – the entry for Urban study area Pittsburgh PA has a superscript “5”
- 27 following it, but no footnote or key to what this refers to.
- 28 P4-71, line 5 – should be Figure 4-19, not F-19.
- 29 P4-75, footnote 61 – second line should read “...related mortality – see section...”
- 30 P4-75, lines 22 thru 27 – this sentence can and should be divided into several shorter
- 31 sentences.
- 32 P5-5, lines 27 thru 29 – this looks like the title of a table or figure, not part of the
- 33 text...???
- 34 P5-6, footnote, second-to-last line – should read “...are subject to greater uncertainty...”
- 35 P6-1, line 31 – remove one of the two periods at the end of the sentence.
- 36 P6-4, line 23 – “INTERPRETATION” is mis-spelled (missing a T).
- 37 P6-6, line 2 – it would be clearer to re-phrase this to read: “...relatively high 24-hr design
- 38 values compared to the respective area’s corresponding annual average design values...”
- 39 P6-6, line 6 – the term “composite monitor” should be defined or explained in a
- 40 parenthetical comment or footnote in this integrative summary chapter, to avoid having
- 41 readers backtrack to previous chapters to try and find the intended meaning.
- 42 P6-6, line 18 – the correct figure reference (for the current standard values) is Figure 4-
- 43 17.
- 44 P6-7, lines 17 & 18 – Actual IHD mortality values in Table 4-1 are 15-19 for Salt Lake
- 45 City and 1755-2222 for New York City.
- 46 P6-7, line 25 – should read “...current suite of standards...”, not “...if standards...”

3-30-10 Deliberative Draft Letter on the Quantitative Health Risk Assessment for Particulate Matter (Second External Review Draft). DO NOT CITE OR QUOTE.

- 1 P6-9, line 27 – change “that” to “than”
- 2 P6-11, line 1 – change “...just meeting the a given...” to “...just meeting a given...”
- 3 P6-14, footnotes 78 and 79 – references to Figure 4-20 should be 4-19
- 4 P6-15, lines 7-9 – This sentence is garbled and confusing, and needs re-writing and
- 5 clarification.
- 6 P6-15, footnote 81 – reference Figure 4-18, not 4-20.
- 7 P6-18, lines 11-12 – delete the rest of the sentence and end after “...2,000 cases per
- 8 year.” (This is repetitive and not needed in this summary statement).
- 9 P6-18, lines 14-15 – delete the rest of the sentence after “...in a given year for the urban
- 10 study areas.”
- 11 P6-18, line 17 – Delete “Generally comparable” and begin the sentence “Estimates of
- 12 CV-related mortality...”
- 13 P6-18, lines 24-25 – Correct first sentence to read, “A broader array of health effects has
- 14 also been associated with PM_{2.5} exposures, including reproductive effects and
- 15 developmental effects in children.”
- 16 P6-18, lines 29-32 – Change first sentence to read: “Given the quantitative and qualitative
- 17 assessments of uncertainty and variability performed as part of the risk assessment, it is
- 18 unlikely that the degree of risk remaining upon the simulated meeting of the current
- 19 (15/35) suite of standards has been over-stated.”
- 20 P6-18, line 41- Re-write to read: “This variability in annual-average PM_{2.5} concentrations
- 21 is most prominent in study areas where the 24-hour standard is “controlling”.”
- 22 P6-19, line 14 – Delete the word “well”; not clear what the definition of “well below a
- 23 value” is, if some of the values are said to be approximately equal to it). In this context,
- 24 it seems sufficient to state that the annual average concentrations under discussion are
- 25 below the level of the current annual standard.
- 26 P6-19, line 19 – delete the phrase “in most such areas”; it is redundant, given the
- 27 beginning of the sentence and the sentence structure.
- 28 P6-19, line 21 – replace “reflective of” with “as do”
- 29 P6-19, line 36 – add “%” after “11”
- 30 P6-21, line 2 – delete comma after “... with a”
- 31 P6-21, line 3 – insert “U.S.” before “...counties”
- 32 P6-21, line 5 – replace the phrase “...likely capture well...” with “do represent”
- 33 P6-21, lines 7-9 – why is this comment here? Isn’t it self-evident in selecting a
- 34 representative sub-sample? As I have commented previously, this is unnecessary and can
- 35 be deleted.
- 36
- 37

1 **Comments from Dr. Joe Brain**

2
3 **Charge Question #6a-e**

4
5 *6) We have developed an integrated discussion of the PM_{2.5}-related risk estimates which*
6 *considers the results of the qualitative and quantitative treatment of uncertainty and*
7 *variability together with the various national-scale assessments completed for the*
8 *analysis to support interpretation of the core risk estimates. As part of the integrative*
9 *discussion, we also provide key observations that bear on policy-relevant risk-based*
10 *questions.*

11
12 *a) To what extent does the Panel believe that we have captured the key policy-relevant*
13 *questions that can be addressed by this risk assessment?*

14
15 Response: Chapter 6 is the culmination a rationale developed throughout the document.
16 It needs to pull all the pieces together and clearly integrate them in a form that provides
17 the basis for decisions. The panel responds positively to Question 6a. We believe that
18 this second draft does effectively capture the key policy-relevant questions that can be
19 addressed by the preceding risk assessment.

20
21 *b) We provide a set of key observations related to estimates of risk associated with*
22 *simulations of just meeting the current and alternative suites of standards. These*
23 *observations are based not only on consideration of trends in risk reduction across*
24 *alternative suites of standards and residual risk remaining after simulation of just*
25 *meeting specific suites of standards, but also on additional factors that can impact risk*
26 *(e.g., the role of annual and 24-hour design values, the peakiness of PM_{2.5} distributions*
27 *within a study area, and application of different rollback approaches). To what extent do*
28 *the Panel members believe that the observations presented in section 6.2 are well*
29 *supported by the results of the analyses? Are there other observations that might be made*
30 *that would help to address the policy-relevant questions identified at the beginning of the*
31 *chapter?*

32
33 Response: The panel believes that Section 6.2, Interpretation of Urban Study Area
34 Results, is well supported by the results of the previous analyses. We are not aware of
35 other observations that might affect the answers to the policy-relevant questions
36 identified at the beginning of Chapter 6.

37
38 *c) Part of our interpretation of the core risk estimates presented in section 6.2 is our*
39 *characterization of confidence in the core risk estimates and in observations made based*
40 *on those estimates. These assessments of confidence are based on consideration of the*
41 *results of the sensitivity analysis as well as on the qualitative assessment of uncertainty*
42 *and variability. To what extent does the Panel believe that the characterizations of*
43 *confidence in the core risk estimates and associated policy-related observations are*
44 *reasonable given available information?*

45

1 Response: The authors have done an appropriate sensitivity analysis and have adequately
2 characterized both uncertainty and variability. The panel believes that the degree of
3 confidence expressed in the core risk estimates is appropriate. We believe that the
4 adjacent policy-related observations are reasonable given the information available to the
5 EPA and summarized in the final version of the PM ISA.

6
7 *d) As part of the integrative discussion, we use the results of several national-scale*
8 *analyses (i.e., the national scale PM_{2.5} mortality analysis, the representativeness analysis,*
9 *and the new exploration of design values and patterns of PM_{2.5} monitoring data presented*
10 *in section 4.5) to place the results of the risk assessment in a broader national-context.*
11 *What are the Panel members' views on appropriateness of this effort to place results of*
12 *the analysis in a national context?*

13
14 Response: One could always suggest additional cities or parts of the country where PM
15 data could be applied. The panel believes that the national-scale analyses which were
16 carried out are appropriate and are reasonably varied and thus they do provide the context
17 for placing the results of the policy analysis in a reasonably representative national
18 context.

19
20 *e) We conclude chapter 6 with a list of key observations. Does the Panel believe that we*
21 *have appropriately highlighted key findings of the risk assessment in these observations?*
22 *Of particular note is the observation that, while alternative 24-hour standard levels can*
23 *be used to reduce annual-average PM_{2.5} concentrations and thus to reduce estimated risk,*
24 *the results are likely to be highly variable across urban areas. More consistent lowering*
25 *of annual-average PM_{2.5} concentrations across study areas, and thus more consistent*
26 *reductions in estimated risk, may result from application of alternative annual standard*
27 *levels. We also note that simulation of the alternative 24-hour standard level of 25 µg/m³*
28 *resulted in reductions in annual-average PM_{2.5} levels for some study areas that were well*
29 *below the lowest annual standard level assessed (i.e., below 12 µg/m³). As a*
30 *consequence, we observed risk reductions reflecting these changes in annual-average*
31 *PM_{2.5} levels below 12 µg/m³. Given these results, does the Panel believe that there is*
32 *utility in estimating risks for alternative annual standard levels below 12 µg/m³?*

33
34 Response: Section 6.4, Key Observations, is a valuable part of this chapter and of the
35 document as a whole. The three questions asked are important and the answers provided
36 are well founded and useful. They get to the heart of the matter. What are the
37 consequences of maintaining the current standard? What are the likely benefits of
38 lowering the standards? To what extent are policies about PM_{2.5} applicable throughout
39 the country? The bullets following the three questions succinctly address these questions,
40 and will be useful to the administrator in forming the basis for her decision.

41
42 Consistent with discussions at the last meeting of the PM advisory committee, we are
43 comfortable with not estimating risks for alternative annual standard levels below 12
44 µg/meter³. At these concentrations, data is increasingly less common and less reliable.
45 Therefore, estimates of risk will be subject to greater and greater uncertainty. We believe

3-30-10 Deliberative Draft Letter on the Quantitative Health Risk Assessment for Particulate Matter (Second External Review Draft). DO NOT CITE OR QUOTE.

- 1 that the risk estimates at levels higher than this provide an adequate basis for decision
- 2 making.

1 **Comments from Dr. Wayne Cascio**

2
3 General Comments:

4 With the second draft assessment document the staff of the EPA’s Office of Air Quality
5 Planning and Standards (OAQPS) has produces an excellent document that largely
6 addresses concerns raised by members of CASAC and the public. Specifically, the
7 second external review draft now contains a more comprehensive discussion of the core
8 risk estimate through the integration of uncertainty and variability analyses from 15 urban
9 centers, and national-scale long-term exposure PM2.5 mortality assessment. The rationale
10 for not including a quantitative assessment of risk associated with PM10-2.5 presented in
11 section 2.3 and appendix H is justified based on the limitations in the data available for
12 characterizing PM10-2.5 exposure and risk. The overall presentation and discussion of
13 the consideration of variability and uncertainty is much improved.

14
15 Charge Question 2. Selection of model inputs (section 3.3): We have expanded and
16 clarified the discussion of our rationale for identifying modeling choices comprising the
17 core risk model, focusing in particular on selection of C-R functions (section 3.3.3). To
18 what extent does the Panel consider this discussion to be clear and the model selections
19 appropriate?

20
21 Response. The expanded discussion provided in Section 3.3.3 Selection of
22 Epidemiological Studies and Concentration-response (C-R) Function within those Studies
23 conveys the rationale for identifying modeling choices and adequately justifies the
24 selection of specific epidemiological studies utilized to establish the C-R functions. The
25 rules employed for study selection as described on 3.27 and 3.28 provided a strong
26 foundation to identify studies that provide the most accurate data to derive the C-R
27 relationship. The expansion of the discussion and integration of the ISA was also very
28 useful in that it provides an opportunity to reinforce the gaps in knowledge, for example
29 on 3-30 where it is stated that, “There were no multi-city studies for this [short-term
30 exposure to PM2.5 and emergency department visits for cardiovascular and/or respiratory
31 illnesses] category of health endpoint.”

1
2 **Comments from Dr. Christopher Frey**

3
4 Charge Question 3:

5
6 In general, the second draft REA appropriately identifies and discusses key sources of
7 variability and uncertainty, and includes sensitivity analysis that provides insight
8 regarding the impact of some sources of uncertainty on the core risk estimates.

9
10 EPA provided a footnote explaining the rationale for identifying “key” sources of
11 variability. The document should indicate if the same process was used to identify “key”
12 sources of uncertainty.

13
14 In the discussion of key sources of variability, EPA added material regarding copollutant
15 concentrations and on demographic and socioeconomic status, as requested in CASAC
16 comments on the first draft of the REA. As a minor comment, it is not entirely clear that
17 age of housing only affects air exchange rate because of air conditioning use. Newer
18 homes are typically “tighter” than older homes, and thus have lower infiltration rates.
19 Climate zones are another factor in infiltration. For example, northeastern homes do not
20 have as high a proportion of central air conditioning as southeastern homes. Given its
21 effect on particle composition, concentrations, ventilation and activity patterns, it would
22 make sense to also include seasonality in the list of variability sources.

23
24 We note that EPA did not address a comment on the first draft REA to the effect that
25 “exposure modeling should be included in the REA. A probabilistic Tier 3 approach
26 should be used for the exposure assessment.” While we understand that timing may have
27 precluded adequate treatment of this topic, we expect that EPA will develop this capacity
28 for future revisions of the standard. CASAC asked for this five years ago, and would
29 like to see this in the next revision.

30
31 With regard to uncertainties, in response to CASAC comments on the first draft of the
32 REA, EPA has included uncertainty in the C-R function itself, which was developed from
33 single studies. EPA has appropriately taken into account differences in C-R functional
34 form associated with studies that addressed long-term or short-term effects for single or
35 multi-city studies even if they were not the basis for the final set of C-R functions used in
36 the REA.

37
38 EPA has provide some explanation of the meaning of uncertainty categories of “low”,
39 “medium”, and “high.” This discussion is adequate. However, there is a confusing
40 statement to the effect that “high” sources of uncertainty “are likely to influence the
41 interpretation of risk...” “if those sources of uncertainty are reduced or more fully
42 characterized.” The parenthetical “if” clause seems to confuse the issue, and should be
43 deleted. A reader of the paragraph on lines 6-20 of p. 3-63 might wonder how “staff
44 consensus” was achieved, and whether consensus is an appropriate goal when
45 characterizing uncertainty. A potential concern is that achievement of “consensus” might
46 mean that some opinions over-ride others and that the resulting characterization of

1 uncertainty might be biased. There are short-comings of group-based elicitation
2 processes, such as dominance by strong personalities or a tendency to provide opinions
3 about goals rather than state of knowledge. It would be useful to explain the process by
4 which “consensus” was achieved.

5
6 EPA has done a nice job on commenting on the extent to which there are dependencies
7 among pairwise combinations of sources of uncertainty, and whether these dependencies
8 would tend to offset or to increase the overall range and direction of uncertainty in the
9 assessment results. For example, the statistical fit of the C-R functions, and the shape of
10 the functions, are inter-related. EPA has provided a nice treatment of this on page 3-71.

11
12 Based on quantifiable sensitivity analysis, the report generally clearly conveys that the
13 “core” estimates appear to be at the low end of alternative “plausible” estimates.
14 However, particularly in Chapter 6, the role of sources of uncertainty treated qualitatively
15 should also be addressed. In particular, given exposure misclassification, it is likely that
16 the core estimates are biased low. This is an important point to convey consistently. The
17 core estimates seem to be conservative in the sense of being underestimated, which is not
18 typical practice for public health endpoints, given that virtually all of the sensitivity
19 analyses result in higher risk estimates as compared to the core.

20
21 Charge Question 4:

22
23 Overall, the sensitivity analysis section 4.3 is very good and nicely covers a complex
24 topic. Table 4-3 is a useful summary. The classification of descriptive categories for
25 small, moderate, moderate-large, and large contributions is useful. However, it would be
26 more appropriate to refer to these as contributions to “sensitivity” rather than
27 “uncertainty.” As noted in several place, the sensitivity analyses represent plausible
28 alternatives to the core estimate, but are not a probability sample. Thus, there is not a
29 probabilistic interpretation to the sensitivity analysis results. EPA has appropriately
30 addressed this point and has clearly articulated, quite reasonably, that the sensitivity
31 analysis results represent plausible and scientifically defensible estimates. The range of
32 these estimates provides an indication of the implications of uncertainty.

33
34 The evaluation of alternative model structure is critically important, because model
35 structure can potentially be a larger source of uncertainty than the range of values for an
36 input to a given model. The results in Table 4-3 indicate, for example, that the random
37 effects log-log model provides larger risk estimates than the fixed effects log-liner model
38 used for the core estimates. This information is very useful and is an excellent addition
39 to the REA. The more thorough treatment of model choices and alternative C-R
40 functions provides plausible alternative estimates to the core estimate.

41
42 It was not apparent that EPA responded to this comment on the first draft REA: “The
43 range of uncertainty associated with confidence intervals for a given C-R function (which
44 is an example of a Tier 3 assessment, which should be mentioned) should be compared to
45 the range of estimates obtained by comparing alternative functional forms. This would
46 provide insight as to whether model structure, or random error for a given model, is a

3-30-10 Deliberative Draft Letter on the Quantitative Health Risk Assessment for Particulate Matter (Second External Review Draft). DO NOT CITE OR QUOTE.

1 more important source of uncertainty.” It would be useful to make this comparison,
2 which can be discussed qualitatively.

3
4 Per CASAC’s comments on the first draft REA, EPA indicates the direction of the
5 percent changes in risk. In addition to the percent difference, the actual difference in risk
6 should be reported to provide further context. The second draft REA seems to put
7 emphasis on relative changes in risk. However, the NAAQS are intended to be protective
8 of public health, and therefore the magnitude of the risk estimates is ultimately a more
9 useful policy-relevant metric.

10
11 The sensitivity analysis related to peak shaving and “peakiness” was not very clear.
12 What are the main points to take away from these analysis could be clarified. Is there an
13 implication of some sort of risk trade-off between the cities with and without
14 “peakiness”, as shown in the comparison of results for the different roll-back approaches?

15
16 What is the premise of the overarching conclusions – e.g., that there are strong regional
17 or inter-city effects? Would this carry forth to the policy analysis in some way?

1 **Comments from Dr. Joseph Helble**

2
3 *1) Air quality inputs (section 3.2): We have expanded the consideration of alternative*
4 *approaches to simulating just meeting the current and alternative suites of PM2.5*
5 *standards (i.e., rollback approaches) to include a peak shaving approach, in addition to*
6 *the hybrid and proportional approaches considered in the first draft assessment. This*
7 *peak shaving approach is intended to represent more localized, rather than regional,*
8 *patterns of PM2.5 reductions (discussed in section 3.2.3.3).*

9
10 *a) To what extent does the Panel believe that the use of the peak shaving approach*
11 *provides useful additional exploration of variability associated with how ambient PM2.5*
12 *concentrations are simulated to change upon just meeting the current and alternative*
13 *suites of standards?*

14
15 It is certainly reasonable to explore alternative approaches to just-meeting standards. The
16 traditional proportional rollback approach was augmented in an earlier draft of the current
17 PM risk assessment by including “hybrid rollback,” in which localized monitor-specific
18 reductions are imposed and then allowed to propagate through an air quality study area
19 through imposition of a distance-decay function. Area-wide proportional rollback is then
20 applied as needed. As noted previously, this seems a reasonable approach to estimating
21 the potential effect of local controls on PM levels for comparison with the region-wide
22 effects approximated by proportional rollback.

23
24 The current question regards the benefits of adding a third approach that involves peak-
25 shaving of concentrations at specific monitors within an air quality study area in an effort
26 to assess the potential effects of highly localized controls. In this approach, only those
27 monitors exceeding the 24 hour standard are rolled back. The use of this alternative to
28 examine hypothetical “what-if” scenarios seems reasonable if the goal is to understand
29 whether reducing concentrations at only the peak monitoring locations would be
30 sufficient to meet overall air quality targets in a given study area. Beyond this, however,
31 the value of the exercise is not clear. It is difficult to imagine a scenario in which such
32 targeted localized controls could be effectively deployed for PM2.5. Given this, the
33 hybrid approach seems more appropriate.

34
35 *b) We have used comparisons of composite monitor annual averages generated using the*
36 *different rollback approaches as a surrogate for differences in long-term exposure-*
37 *related mortality in looking across all three rollback approaches. To what extent does the*
38 *Panel believe that this is a reasonable approach for assessing the impact of variability*
39 *associated with simulating changes in air quality patterns on estimates of long-term*
40 *exposure-related mortality?*

41
42 Given the uncertainty in the data, particularly due to incomplete monitoring datasets or
43 differences in data collection patterns from region to region, this approach is reasonable.

44
45 Related to this, regarding the approach to generating composite data sets in locations
46 where there are gaps in individual monitor datasets, was the accuracy of the interpolation

3-30-10 Deliberative Draft Letter on the Quantitative Health Risk Assessment for Particulate Matter (Second External Review Draft). DO NOT CITE OR QUOTE.

1 routines tested? If at least one urban study area has a complete data set, the accuracy of
2 the interpolation approach could be assessed by artificially (randomly) deleting specific
3 values from the monitoring dataset, interpolating as before, and then comparing the
4 estimated values with actual.

5
6
7

1 **Comments from Dr. Rogene Henderson**

2
3 Comments on Assigned Charge Questions for 2nd Draft of PM-RA

4
5 Charge Question 5: Consideration of design values and patterns of PM_{2.5} monitoring
6 data in interpreting core risk estimates (section 4.5): To enhance our interpretation of the
7 patterns of core risk estimates generated for both the current and alternative suites of
8 standards, we have included analyses of 24-hour and annual design values together with
9 patterns of PM_{2.5} monitoring data for the 15 urban study areas. This reflects the fact that
10 these two factors play a key role in determining the degree of risk reduction estimated
11 upon just meeting the current and alternative suites of standards under alternative
12 rollback approaches. As part of the consideration of design values, we have also
13 contrasted the 15 urban study areas with patterns of design values seen for the broader set
14 of urban areas in the U.S. in order to help place the urban study area in a broader national
15 context

16
17 a) To what extent is the Panel supportive of these additional assessments?

18 I found Figures 17-21 quite helpful in putting the 15 urban areas into context with the
19 monitoring values from other cities in the US. The figures also helped me to visualize
20 what the overall picture looked like and what the controlling values are.

21 I would like to have more discussion about how the peak shaving rollback method
22 was used. When the short-term values are controlling, it does not make sense to me to
23 use the same percentage rollback for the annual values as is required by the short-term
24 values. This results in non-feasible annual values (e.g., 7 ug/m³) for some cities such as
25 SLC. Do we have actual evidence to indicate that if the short-term value were reduced
26 55% that the annual value would also be reduced 55%? I do not see the logic to that
27 assumption.

28 Why not consider the risk of mortality from short-term exposures separately from the risk
29 of mortality from the average annual exposures. It is stated in the beginning of Chapter
30 6 (page 6-1) that the primary focus will be based on risk associated with long-term
31 exposure to PM, because long-term exposure to PM_{2.5} has been shown to produce
32 substantially larger mortality risk compared to short-term PM_{2.5} exposure. Based on
33 that, one might expect the calculated risks from the annual exposures to be the most
34 critical consideration to protect public health. I do not think it is appropriate to mix the
35 two design values in the rollback procedures.

36 But perhaps I do not understand how the peak shaving rollback method was used.
37 It states on page 3-15, lines 20-23, that the proportional rollback method was the only one
38 used to generate core risk estimates and the other two rollback methods were only used
39 for sensitivity analysis. I would appreciate more discussion of this at our meeting.

40 I suggest adding the term "design value" to the list of terms on page viii, with a
41 reference to page 3-16 for a definition.

42
43 b) Does the Panel have any recommendations for additional insights based on
44 consideration of patterns in design values and PM_{2.5} monitoring data across the 15 urban
45 study areas and at the national level?

1 It would be nice to have Figures 17-19 in three dimensions with some measure of health
2 effects for the 15 cities on the third axis. Can that be done? It stated that the C-R
3 functions are fairly linear and it would be nice to see that.

4
5 Chapter 6 – Integrative Discussion of PM2.5-related Risks

6
7 6) We have developed an integrated discussion of the PM2.5-related risk estimates which
8 considers the results of the qualitative and quantitative treatment of uncertainty and
9 variability together with the various national-scale assessments completed for the analysis
10 to

11 support interpretation of the core risk estimates. As part of the integrative discussion, we
12 also provide key observations that bear on policy-relevant risk-based questions.

13
14 a) To what extent does the Panel believe that we have captured the key policy-relevant
15 questions that can be addressed by this risk assessment?

16 I thought the policy-relevant questions were well stated.

17
18 b) We provide a set of key observations related to estimates of risk associated with
19 simulations of just meeting the current and alternative suites of standards. These
20 observations are based not only on consideration of trends in risk reduction across
21 alternative suites of standards and residual risk remaining after simulation of just meeting
22 specific suites of standards, but also on additional factors that can impact risk (e.g., the
23 role of annual and 24-hour design values, the peakiness of PM2.5 distributions within a
24 study area, and application of different rollback approaches). To what extent do the Panel
25 members believe that the observations presented in section 6.2 are well supported by the
26 results of the analyses? Are there other observations that might be made that would help
27 to address the policy-relevant questions identified at the beginning of the chapter?

28 I thought the list of observations was appropriate.

29
30 c) Part of our interpretation of the core risk estimates presented in section 6.2 is our
31 characterization of confidence in the core risk estimates and in observations made based
32 on those estimates. These assessments of confidence are based on consideration of the
33 results of the sensitivity analysis as well as on the qualitative assessment of uncertainty
34 and variability. To what extent does the Panel believe that the characterizations of
35 confidence in the core risk estimates and associated policy-related observations are
36 reasonable given available information?

37 The confidence statements were well presented.

38
39 d) As part of the integrative discussion, we use the results of several national-scale
40 analyses (i.e., the national scale PM2.5 mortality analysis, the representativeness analysis,
41 and the new exploration of design values and patterns of PM2.5 monitoring data
42 presented in
43 section 4.5) to place the results of the risk assessment in a broader national-context. What
44 are the Panel members' views on appropriateness of this effort to place results of the
45 analysis in a national context?

46 I thought it was a good idea.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16

e) We conclude chapter 6 with a list of key observations. Does the Panel believe that we have appropriately highlighted key findings of the risk assessment in these observations? Of particular note is the observation that, while alternative 24-hour standard levels can be used to reduce annual-average PM_{2.5} concentrations and thus to reduce estimated risk, the results are likely to be highly variable across urban areas. More consistent lowering of annual-average PM_{2.5} concentrations across study areas, and thus more consistent reductions in estimated risk, may result from application of alternative annual standard levels. We also note that simulation of the alternative 24-hour standard level of 25 µg/m³ resulted in reductions in annual-average PM_{2.5} levels for some study areas that were well below the lowest annual standard level assessed (i.e., below 12 µg/m³). As a consequence, we observed risk reductions reflecting these changes in annual-average PM_{2.5} levels below 12 µg/m³. Given these results, does the Panel believe that there is utility in estimating risks for alternative annual standard levels below 12 µg/m³?

Definitely not. I think there is too much uncertainty to do that with much confidence.

1
2 **Comments from Dr. Philip Hopke**
3

4 In general, there are no major problems with this assessment. It is generally following
5 the approaches that we have now seen in multiple rounds of review.
6

7 1) Air quality inputs (section 3.2): We have expanded the consideration of alternative
8 approaches to simulating just meeting the current and alternative suites of PM_{2.5} standards
9 (i.e., rollback approaches) to include a peak shaving approach, in addition to the hybrid
10 and proportional approaches considered in the first draft assessment. This peak shaving
11 approach is intended to represent more localized, rather than regional, patterns of PM_{2.5}
12 reductions (discussed in section 3.2.3.3).

13 a) To what extent does the Panel believe that the use of the peak shaving approach
14 provides useful additional exploration of variability associated with how ambient PM_{2.5}
15 concentrations are simulated to change upon just meeting the current and alternative
16 suites of standards?
17

18 It does not seem useful to me since it is unlikely that there are obvious sources that could
19 be controlled that would let one shave peaks in practice. Thus, it provides unrealistic
20 scenarios that really do not contribute useful information to the assessment. The other
21 rollback approaches seem fine and it is hard to see a role for the peak shaving approach.
22

23 b) We have used comparisons of composite monitor annual averages generated using the
24 different rollback approaches as a surrogate for differences in long-term exposure-related
25 mortality in looking across all three rollback approaches. To what extent does the Panel
26 believe that this is a reasonable approach for assessing the impact of variability
27 associated with simulating changes in air quality patterns on estimates of long-term
28 exposure-related mortality?
29

30 It would make the compositing easier to follow if equations were provided. If you do not
31 wish to put them into the main chapter, then an appendix can be added to guide the
32 interested reader through exactly how all of the calculations were performed.
33

34 I agree with the approach to removing quarters in which too few values are reported to
35 provide a reliable estimate of the average concentration during that period. However, it
36 makes little sense to replace missing values with a mean value. If you are going to
37 attempt to impute missing values, then a much more sophisticated approach should be
38 employed using other monitors in the area as well as historic data for similar
39 meteorological conditions. I would suggest that for those quarters where there are
40 missing values, but a sufficient number of values to provide a valid mean value, then that
41 should be the value used in the health analyses.
42

43 2) Selection of model inputs (section 3.3): We have expanded and clarified the discussion
44 of our rationale for identifying modeling choices comprising the core risk model,
45 focusing in particular on selection of C-R functions (section 3.3.3). To what extent does
46 the Panel consider this discussion to be clear and the model selections appropriate?

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15

3) Addressing uncertainty and variability (section 3.5): We have clarified the process used to evaluate sources of variability and added coverage for specific sources of variability (section 3.5.2); expanded our discussion of the qualitative analysis of uncertainty (section 3.5.3); and included analyses of pair-wise interactions of sources of uncertainty (section 3.5.4). To what extent does the Panel consider these discussions to be clear and appropriate?

It is time to move toward stochastic uncertainty analysis. Every time we get a risk assessment, we continue to get the qualitative review of uncertainties and some hand waving. If there are insufficient resources available to do the full stochastic risk assessment, then the Agency should state this clearly so that it can be made clear why they have chosen not to move to a more complete analysis. Otherwise, we should not see them continue to duck this approach.

1 **Comments from Dr. Morton Lippmann**

2
3 **Overall Comments:**

4 The authors have been very responsive to the prior CASAC PM Panel comments and
5 suggestions, and the 2nd draft provides a thorough and well-prepared presentation of
6 realistic quantitative estimates of the effects of long-term ambient air PM_{2.5} exposures on
7 premature mortality in 15 US urban areas, their variability and uncertainties, and their
8 representativeness of the US population as a whole. The alternate risk assessments (RAs)
9 for recent years' concentrations, meeting the current suite of PM_{2.5} NAAQS, and alternate
10 NAAQS (14, 13, and 12 ug/m³ annual and 30 and 25 ug/m³ daily, and combinations
11 thereof) is an appropriate way to tee up the selection options for the Policy Assessment
12 (PA).

13
14 Having offered a strong endorsement of the overall work of the OAQPS in preparing the
15 RA document, I need to raise two issues that I have raised before that have not been
16 properly addressed. These are:

- 17
18 1) The fact that the ACS cohort is not a representative US population, but rather is of
19 higher SES. Within this cohort, the mortality risk is lower for those of higher
20 SES. Thus, the risk coefficient for a representative population would be higher, as
21 it is for the 6-cities cohort that was selected to be more representative (at least for
22 cities in the eastern half of the US).
- 23 2) The description and use of the data from the Ito et al. (2007) study has not been
24 corrected. As noted below, this was a study covering all of New York City
25 (Kings, Queens, New York, Bronx, and Richmond Counties), not just New York
26 County (Borough of Manhattan).

27
28 **Specific Comments on Text Entries:**

29
30 Page Line Comment

31
32 3-13 Table 3.1. The entry for New York City is incorrect. Change New York City
33 (Manhattan)" to "New York County (Manhattan)".

34
35 3-13 6 Change "New York (Manhattan)" to "New York City".

36
37 3-30 11 Delete "(Manhattan)".

38
39 3-31 7 The number of members of the ACS cohort in the 156 MSAs with air
40 quality data is much smaller than 1.2 million.

41
42 3-40 Counties Column for "New York". Change New York City (Manhattan)" to "New
43 York County (Manhattan)", and delete "New York City (Manhattan)" where it precedes
44 "Ito et al. (2007)". The Ito et al. (2007) study covered all of NYC, not just Manhattan!

3-30-10 Deliberative Draft Letter on the Quantitative Health Risk Assessment for Particulate Matter (Second External Review Draft). DO NOT CITE OR QUOTE.

1 3-50 Table 3-10. There should not be two entries for New York, NY. As noted above,
2 the Ito et al. (2007) study covered all of NYC, not just Manhattan! The second New
3 York entry, which shows only the population of New York County (Manhattan), should
4 be deleted, and the calculations for the Ito et al. study need to be revised to reflect the
5 whole NYC population and all of the NYC PM_{2.5} monitors.
6

7 3-77 Section 3.54.3. This section refers to single- and multiple-elements, as well as to
8 single- and multiple-factors, in terms of sensitivity analyses. There is no definition
9 provided as to what constitutes an element or a factor. I suggest that element is a poor
10 choice in terms of terminology, since it implies, at least to me, a chemical element. This
11 will become important in the next round of PM NAAQS review. [On page 4-42, there is a
12 clarification indicating that elements refers to “modeling elements”]
13

14 4-19 10 Change “fig” to “fit”.

15
16 4-19 11 Change “greater” to “great”.

17
18 4-36 12 Change “CPD” to “CVD”.

19
20 4-45 Note at bottom of the page: Change “Figures 4-6 and 4-7” to “Figure 4-8”.

21
22 4-47 29 Change “Figures 4-6 and 4-7” to “Figures 4-7 and 4-8”.

23
24 4-48 para. 2 : Once again, there are elements and factors with no definitions. Here the
25 elements are not modeling elements, but something else. [On page 4-54, line 21, I learned
26 that these were “critical risk function elements”]. Please reword to avoid confusion!
27

28 4-55 23 Change “population” to “populated”.

29
30 6-1 31 Delete extra period at the end.

31
32 6-14 25 Change “Pitts” to “Pittsburgh”.

33
34 6-18 15,16 Where can we find these estimates?

35
36 6-18 21-23 Where can we find these estimates?
37

38 6-21 8 Change “PM2.5” to “PM_{2.5}”.

39
40
41 **Charge Questions to the CASAC PM Review Panel** - focus on the charge questions
42 listed below in review of the *Quantitative Health Risk Assessment for Particulate Matter*
43

44 Chapter 3 – Urban Case Study Analysis Methods
45

46 **Charge Question #2: Selection of model inputs** (section 3.3):

1
2 We have expanded and clarified the discussion of our rationale for identifying modeling
3 choices comprising the core risk model, focusing in particular on selection of C-R
4 functions (section 3.3.3). To what extent does the Panel consider this discussion to be
5 clear and the model selections appropriate?
6

7 **Response:**

8 The Panel commends the authors for expanding and clarifying their rationale for
9 identifying modeling choices comprising the core risk model in a logical and satisfactory
10 manner. Their model selections were appropriate
11

12 Chapter 6 – Integrative Discussion of PM_{2.5}-related Risks
13

14 **Charge Questions #6a-e:**

15 We have developed an integrated discussion of the PM_{2.5}-related risk estimates which
16 considers the results of the qualitative and quantitative treatment of uncertainty and
17 variability together with the various national-scale assessments completed for the analysis
18 to support interpretation of the core risk estimates. As part of the integrative discussion,
19 we also provide key observations that bear on policy-relevant risk-based questions.
20

21 **Charge Questions #6a:** To what extent does the Panel believe that we have captured the
22 key policy-relevant questions that can be addressed by this risk assessment?
23

24 **Response:**

25 The Panel considers that the authors have captured the key policy-relevant questions that
26 can be addressed by this risk assessment that is focused solely on PM_{2.5}.
27

28 **Charge Questions #6b:**

29 We provide a set of key observations related to estimates of risk associated with
30 simulations of just meeting the current and alternative suites of standards. These
31 observations are based not only on consideration of trends in risk reduction across
32 alternative suites of standards and residual risk remaining after simulation of just meeting
33 specific suites of standards, but also on additional factors that can impact risk (e.g., the
34 role of annual and 24-hour design values, the peakiness of PM_{2.5} distributions within a
35 study area, and application of different rollback approaches).
36

37 To what extent do the Panel members believe that the observations presented in section
38 6.2 are well supported by the results of the analyses? Are there other observations that
39 might be made that would help to address the policy-relevant questions identified at the
40 beginning of the chapter?
41

42 **Response:**

43 The Panel considers that the observations presented in section 6.2 are consistent with the
44 analytical results, and provide all of the information needed in the development of the PA
45 document.
46

1 **Charge Questions #6c:** Part of our interpretation of the core risk estimates presented in
2 section 6.2 is our characterization of confidence in the core risk estimates and in
3 observations made based on those estimates. These assessments of confidence are based
4 on consideration of the results of the sensitivity analysis as well as on the qualitative
5 assessment of uncertainty and variability.
6

7 To what extent does the Panel believe that the characterizations of confidence in the core
8 risk estimates and associated policy-related observations are reasonable given available
9 information?
10

11 **Response:**

12 The Panel considers that the characterizations of confidence in the core risk estimates and
13 associated policy-related observations are reasonable.
14

15 **Charge Questions #6d:** As part of the integrative discussion, we use the results of
16 several national-scale analyses (i.e., the national scale PM_{2.5} mortality analysis, the
17 representativeness analysis, and the new exploration of design values and patterns of
18 PM_{2.5} monitoring data presented in section 4.5) to place the results of the risk assessment
19 in a broader national-context.
20

21 What are the Panel members' views on appropriateness of this effort to place results of
22 the analysis in a national context?
23

24 **Response:**

25 The Panel considers that the effort to place results of the analysis in a national context
26 was appropriate.
27

28 **Charge Questions #6e:** We conclude chapter 6 with a list of key observations.
29 Does the Panel believe that we have appropriately highlighted key findings of the risk
30 assessment in these observations?
31

32 **Response:**

33 The Panel considers that Staff has appropriately highlighted key findings of the risk
34 assessment in these observations
35

36 Of particular note is the observation that, while alternative 24-hour standard levels can be
37 used to reduce annual-average PM_{2.5} concentrations and thus to reduce estimated risk, the
38 results are likely to be highly variable across urban areas. More consistent lowering of
39 annual-average PM_{2.5} concentrations across study areas, and thus more consistent
40 reductions in estimated risk, may result from application of alternative annual standard
41 levels. We also note that simulation of the alternative 24-hour standard level of 25 µg/m³
42 resulted in reductions in annual-average PM_{2.5} levels for some study areas that were well
43 below the lowest annual standard level assessed (i.e., below 12 µg/m³). As a
44 consequence, we observed risk reductions reflecting these changes in annual-average
45 PM_{2.5} levels below 12 µg/m³.
46

3-30-10 Deliberative Draft Letter on the Quantitative Health Risk Assessment for Particulate Matter (Second External Review Draft). DO NOT CITE OR QUOTE.

1 Given these results, does the Panel believe that there is utility in estimating risks for
2 alternative annual standard levels below 12 $\mu\text{g}/\text{m}^3$?

3

4 **Response:**

5 The Panel does not consider that estimating risks for alternative annual standard levels
6 below 12 $\mu\text{g}/\text{m}^3$ is a worthwhile endeavor on several grounds. One, the extrapolation of
7 the risk coefficients at such concentrations becomes increasingly uncertain; and two, the
8 prospects of achieving compliance with such levels in the foreseeable future is
9 vanishingly small.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42

Comments from Dr. Robert Phalen

General Comments:

The second draft of “Quantitative Health Risk Assessment for Particulate Matter” is clearly-written and the logical development is well-described. The staff has done an excellent job. My enthusiasm for the entire document is diminished by factors that are apparently outside of policy-driven limitations imposed on EPA staff. These limitations include:

- PM2.5 is a mass-based metric (indicator) for assessing health-effects. It is likely that specific components, such as vanadium, nickel, and elemental carbon, are actually driving the health effects. The use of a mass-based indicator can lead to air-quality standards that do not permit appropriate abatements.
- Secondary health effects, e.g. as generated by abatements that adversely affect the economy (and the many associated health effects), are not considered. Such indirect adverse health effects are real, and the affected populations must face them along with the direct effects.
- The current risk assessment does not conform to the recommendations made by the National Research Council of the National Academies (Science and Decisions: Advancing Risk Assessment, the National Academies Press, Washington D.C., 2008). Specifically, ... “that risk assessment should be viewed as a method for evaluating the relative merits of various options for managing risk rather than as an end in itself.” The concept of managing risk must include all of the significant risks associated with a particular decision, not just some selected direct effects associated with the decision.
- National air quality standards have the disadvantage of forcing abatements on regions of the U.S. for which the abatements may harm health to a greater extent than they improve health. As an example consider restrictions on diesel-emissions to try to meet standards in areas where a significant portion of PM2.5 can be found in soil fine-particle mass. Control of soil aerosols is not practical, so the diesels must be targeted.

Specific Comments on Section 3.3

- This reviewer found only minor errors.
- Table 3-4, pg. 3-24: Replace “LA” with “Los Angeles”, as LA is also the designator for Louisiana.
 - Line 34, pg. 3-31: Drop either “the” or “our” for clarity.

1
2 **Comments from Dr. Ted Russell**
3

4 Overall, I continue to be pleased with the amount of analysis conducted in this PM Risk
5 Assessment (RA), though I am still disappointed that a more comprehensive exposure
6 modeling effort was not made. The document provides the type of information needed to
7 inform the review of the primary NAAQS, and does a good job of providing a
8 quantitative assessment of the potential risks, and has done a commendable sensitivity
9 analysis.

10
11 While I appreciate the addition of Chapter 6, it was not as effective as it might be. First, I
12 was looking for Chapter 6 to be more integrative of the results from Chapter 4 and 5 as it
13 is now more focused on Chapter 4. Second, I was looking for it to also integrate
14 uncertainties in to the discussion to a greater degree. This is the point that they could
15 bring in the issue of using the LML as the zero risk level, and how that might influence
16 the overall interpretation of the results. Third, it was a bit repetitive, bringing up the
17 issue of “peaky nature” and explaining it more often than necessary. It is an important
18 concept, but it was overdone in Chapter 6. Still, Chapter 6 is a good addition, but it could
19 be made stronger in response to the above comments.
20

21 *Chapter 3 Charge Questions:*
22

23 *1) Air quality inputs (section 3.2): We have expanded the consideration of alternative*
24 *approaches to simulating just meeting the current and alternative suites of PM_{2.5}*
25 *standards 5 (i.e., rollback approaches) to include a peak shaving approach, in addition*
26 *to the hybrid and proportional approaches considered in the first draft assessment. This*
27 *peak shaving approach is intended to represent more localized, rather than regional,*
28 *patterns of PM_{2.5} reductions (discussed in section 3.2.3.3).*
29

30
31 *a) To what extent does the Panel believe that the use of the peak shaving approach*
32 *provides useful additional exploration of variability associated with how ambient PM_{2.5}*
33 *concentrations are simulated to change upon just meeting the current and alternative*
34 *suites of standards?*
35

36 Response: This approach, along with the hybrid approach, provide a reasonable method
37 to get an estimate of the lower bound impact of what would happen when an urban area
38 attains a specific standard. This approach is specifically of interest in areas where the 24-
39 hour standard will be the driving standard, and the proportional roll-back would lead to
40 reductions in the annual level beyond that which might be viewed as likely. The method,
41 either in the report or in the appendices, should be specified mathematically, as well as in
42 words. They might present a specific set of example applications of the approaches, e.g.,
43 for three different cities, at this point.
44

45 *b) We have used comparisons of composite monitor annual averages generated using the*
46 *different rollback approaches as a surrogate for differences in long-term exposure-*

1 *related mortality in looking across all three rollback approaches. To what extent does the*
2 *Panel believe that this is a reasonable approach for assessing the impact of variability*
3 *associated with simulating changes in air quality patterns on estimates of long-term*
4 *exposure-related mortality?*

5
6 Response: First, the approach to compositing should be better explained. A critical
7 question is how well it aligns with what is actually done in reporting for the calculation of
8 design values. Indeed, the approach they used should be motivated by first explaining
9 how the annual average PM is found (provide specific equations as needed), and then
10 showing that the approach taken is in line with how the annual average PM is now found.
11 Also, they need to explain how and when compositing is used in calculating design
12 values. This will explain why quarterly averages are first calculated, and may impact my
13 thoughts on how the current approach to compositing should be altered.

14
15 I do not like how they replace missing values, i.e., using the average value to replace
16 missing values when a certain number of samples are missing. It seems to be a bit
17 arbitrary, and could lead to a bias. Is this what is done in practice (i.e., specified)? In
18 terms of compositing, the values could each be adjusted using a centering approach. In
19 this case, the annual average from each monitor being used in the composite is subtracted
20 from the daily value from that monitor, leading to a string of values that have a mean of
21 zero. These annual averages are also used to calculate the composite annual average of
22 the stations being used in the composite. The daily values for each monitor (after
23 subtracting the mean of that station) are then averaged as available. This leads to the
24 average variation from the mean for that day. The annual average composite value is
25 then added back to get the daily composite value. This is relatively insensitive to stations
26 dropping out. Further, it should exactly give the observed annual mean at each station,
27 and well as the composite mean. The current approach for imputing missing days can
28 lead to an average that would not agree with the reported value. The same approach for
29 calculating quarterly averages, that are then used to calculate the annual average and
30 design value, can be used.

31
32 Like my response to part (a), the mathematical equations should be provided here or in
33 the appendix.

34
35 *5) Consideration of design values and patterns of PM_{2.5} monitoring data in interpreting*
36 *core risk estimates (section 4.5): To enhance our interpretation of the patterns of core*
37 *risk estimates generated for both the current and alternative suites of standards, we have*
38 *included analyses of 24-hour and annual design values together with patterns of PM_{2.5}*
39 *monitoring data for the 15 urban study areas. This reflects the fact that these two factors*
40 *play a key role in determining the degree of risk reduction estimated upon just meeting*
41 *the current and alternative suites of standards under alternative rollback approaches. As*
42 *part of the consideration of design values, we have also contrasted the 15 urban study*
43 *areas with patterns of design values seen for the broader set of urban areas in the U.S. in*
44 *order to help place the urban study area in a broader national context*

45
46 *a) To what extent is the Panel supportive of these additional assessments?*

1
2 Response: These are beneficial. Figures 4-17 to 4-19, and Table 4-8 do a good job of
3 identifying which are the controlling standards for each of the 15 cities, and also show
4 that the cities examined do a good job of spanning the space of conditions for cities
5 around the US. It might have been good to also identify some of the major outliers on
6 Fig. 4-17 (there really is only one to identify). Also, you could consider color coding the
7 dots on the figure, such that each portion of the US is assigned a color, so it would be
8 apparent if there were specific areas where an issue might identified. This would also
9 help address why no Upper Midwest city is targeted. Without this, I don't think this
10 section really does as good job of contrasting the 15 urban study areas with other areas,
11 that they are supposed to represent, particularly at a regional level. Section 4.4 presents
12 additional information putting other areas in to perspective, but what would be valuable is
13 to show that all regions and sizes of cities are addressed adequately.

14
15 *b) Does the Panel have any recommendations for additional insights based on*
16 *consideration of patterns in design values and PM_{2.5} monitoring data across the 15 urban*
17 *study areas and at the national level?*

18
19 Response: See above. I think this section could be strengthened by being able to have a
20 concluding statement somewhat along the lines of “We have captured the range of
21 conditions found for cities that would be in non-attainment of the various combinations
22 of standards in each of the six of the seven (I would have preferred all seven) regions of
23 the country, and the 15 cities capture X% of the population. The analysis has identified
24 cities that span the range of non-attainment levels, and which standard would require the
25 greatest level of control based on this analysis.” It is close to being able to say this.

26
27 Other details:

28
29 Page 3-8, Footnote: This footnote is not needed and I find it confusing. Also, while it
30 need not be added at this point, unless $\beta\Delta x$ is above about 0.2, the error in linearizing (3)
31 and (4) is rather small. This might simplify interpretation of later analyses since it
32 makes the response to concentration changes linear.

33
34 In Chapter 3, no city in the Upper Midwest was identified. Might this be alleviated?
35
36

1 **Comments from Dr. Frank E. Speizer**

2
3 Pre-meeting Comments: 3/5/2010

4 General Comment:

5 Staff has done an excellent job of using the available literature and assessments from the
6 ISA to generate a series of outcome assessments over an appropriate range of alternative
7 levels of both 24 hour and annual averages of PM2.5 that are justified by the available
8 data. They appear to have been responsive to our previous comments on the first draft, and
9 where they have not followed our suggestions have presented evidence that they
10 considered our requests and accepted or rejected or modified with appropriate
11 justification. The one area where I would have liked to have seen more analyses is with
12 regard to PM10-2.5, simply to have demonstrated the “inappropriateness” of trying to
13 rely on such an analysis would have emphasized the need for more data related to the
14 coarse fraction effects.

15
16 Specific Comments and Charge Questions

17 Chapter 2

18 Page 2.4, line 14. Typo: PM2.5

19
20 Page 2.6, line 2-4 Query the continuing developmental work on population exposure
21 analysis methodology. It would have been useful to read or hear further what this plan
22 would be. One would have thought that over the years much of this would have already
23 been worked out.

24
25 Page 2.6, line 22-24. Appendix H provides a well reasoned argument for Staff concluding
26 that they could not do a quantitative risk assessment for PM10-2.5. From my perspective
27 it would seem to me that some estimate of risk should be made if for no other reason than
28 to document wide confidence intervals and thus reason for not including it in the
29 quantitative assessment.

30
31 Page 2.7, line 10-11. Perhaps at the end of this process a note could be sent to CASAC to
32 indicate what the plan might be for the future to prepare for the next round by carrying
33 out the methods development necessary to use “specialized analysis of risk...” Is this
34 important enough for CASAC to include such a request in our letter to the administrator?
35

36 Page 2.12, line 13-20. The full set of model choices offered seems appropriate, given the
37 plan is to maintain an annual as well as a 24 hour standard..

38
39 Chapter 3

40 Page 3.19-3.20 and Charge Question 1a and 1b. Use of peak shaving rollback approach
41 as discussed is confusing. Staff indicates (at top of page 3.20) that because of time
42 constraints they did not calculate health risks using this method. They go on to indicate
43 what the method does. The appendix that discusses the method more fully refers to
44 Tables F-49 and F-50. These tables present the curious finding that except for Fresno
45 (where there is virtually no change) and Tacoma, all of the other cities show a
46 progressive increase in maximum values going from the “proportional” to “hybrid” to

3-30-10 Deliberative Draft Letter on the Quantitative Health Risk Assessment for Particulate Matter (Second External Review Draft). DO NOT CITE OR QUOTE.

1 “shaving”. This certainly does not seem to indicate that the shaving analyses were not
2 done. It also raises for me concern that more effort is needed to understand which is a
3 better technique or more representative of “truth”. Thus I am worried that “time
4 constraints may be once again getting in the way of what the proper analysis should be.

5
6 Section 3.3., Charge Question 2 asks about the discussion of rationale for modeling
7 choices and selection of CR function. The section follows a logical rationale and
8 provides appropriate documentation of both model and site selection.

9
10 Page 3.49, Section 3.4.1.3. It might be worth adding a sentence that approximately 52 x
11 10⁶ people are represented in the 15 cities or about 20% of the total population of US
12 (even if not a representative sample).

13
14 Page 3.53-54, Table 3.11. there is far too much missingness in this table to make it at all
15 useful. It raises more questions about the quality of the health input than would be
16 justified. The lack of COPD for all sites but LA just doesn't compute for the 4th major
17 cause of death in the US. Why are there not values for All Ages for All Causes? The data
18 are presented for some but not all categories of disease. Someone needs to take the time
19 to get all these cells filled in. These data all exist at the Federal level in one book o(or on
20 one web site)!

21
22 Charge Question 3, Uncertainty and Variability. The discussion is quite complete dealing
23 with a number of issues. What is not fully considered is the role of the potential for non-
24 random missingness in both site selection within the 15 cities and thus the selection of the
25 cities themselves.

26
27 Chapter 4

28 Charge Question 4-Sensitivity Analysis

29 Section 4.3, Table 4.1, page 4.6. This table needs to define denominators for Incidence
30 (in title). Assuming both are the same then in general hospitalization considerably more
31 frequent than mortality. (My concern is mortality may be x/100,000 and morbidity might
32 be y/10,000 and if this is the case there are strikingly different numbers that need to be
33 discussed more fully).

34
35 Page 4.17, line 6. Need to indicate denominators for these incidence rates.

36
37 Page 4.46 Figures 4.7 & 4.8. I have difficulty in interpreting these figures. For HD total
38 incidence for LA go from 5-10% to as high as 19%. For Philadelphia from 10-15%
39 to 14-15% with outlier at 23%. For total mortality the variation are separately 2-6% and
40 3-4 to 8%. For both these cases the core analysis seems too high and the delta seems too
41 big, as tests of sensitivity. Is this the wrong interpretation?

42
43 Page 4.47, line 29. Should this be figures 4.7 and 4.8 rather than 4.6 and 4.7?

44 Page 4.54, line 5. This should be changed to Table 4.7 (page 4.56).

45 Line 21. Figure numbers seem not to match up with text being discussed. Please
46 check.

1
2 Page 4.67-4.69, Figures 4.17-4.19. These are very effective presentations of design
3 values. I suggest either change titles or text numbers so that 24 hour/annual are presented
4 consistently in both.

5
6 Charge Question 5a. Design values analysis and assessments. These analyses are quite
7 useful, as they are described. They provide visual interpretations directly of the range of
8 effects related to the alternative standards. Notably in Chapter 3 the suggestion is made
9 tht the use of peak rollback shaving is not to be considered; however, it looks like it is
10 being used here.

11
12 Charge Question 5b. Additional suggestions. I think the ranges of outcomes are well
13 demonstrated in figures 4.20 and 4.21. Rather than simply ending with the description of
14 how they were constructed it might be worth adding a paragraph or two that summarize
15 the degree to which the various alternative would provide changes in some fraction or all
16 of the cities, rather than just ending with selected examples (maybe this will happen in
17 Chapter 6).

18 19 Chapter 5

20 This chapter leaves me uneasy and I would like to suggest we spend some time
21 discussing it. It look to me that it is a straight forward extrapolation of results from the
22 two large cohort studies (ACS and 6 Cities). The level of exposure are extracted from the
23 15 urban areas and then scaled up to the US. This leads to ~88,000 (4-8% of total
24 mortality—a figure repeated in Chapter 6). However, this seems too high. At one point
25 there is a suggestion that the figures come from upper end of urban mortality risk, but
26 isn't the calculation for the whole nation where exposure levels must be a lot lower? Lets
27 discuss!

28 29 Chapter 6

30 Page 6;.7, line 25. Typo change “if “to “of”

31
32 General Comment: The integrated discussion pulls together and highlights some of the
33 specific details presented in Chapter 4 and the Appendices. It reflects the obvious and
34 not so obvious conclusions that results from manipulating the various alternative
35 scenarios between current existing exposure, estimates to current NAAQS levels and the
36 various alternatives proposed. In spite of the evidence that there does not appear to be a
37 threshold the lowest level assessed 12/25 seems appropriate in that the evidence of an
38 effect below those levels is simply too uncertain to evaluate. On the other hand the
39 question of margin of safety remains and it will need to be argued that any level chosen
40 (above that level) will need to be defended as to whether there is adequate margin of
41 safety.

42
43 With regard to the Charge Questions:

- 44 a. Staff has done an excellent job in presenting and capturing the key-policy relevant
45 questions. However, as indicated early on in this document there was to be a
46 qualitative discussion on PM10-2.5 and on those effects that were deemed only

3-30-10 Deliberative Draft Letter on the Quantitative Health Risk Assessment for Particulate Matter (Second External Review Draft). DO NOT CITE OR QUOTE.

1 “suggestive” but might have important public health implications (e.g. lung
2 cancer, reproductive effects), but for which quantitative risk assessment was not
3 thought warranted but that would appear in the PA (page 2.6). I would have
4 thought that some remarks in this chapter would be necessary to assure that the
5 PA would discuss the issues.
6

- 7 b. Key observations are presented and discussed with adequate discussion of the
8 relevant contribution of the role of annual and 24 hour design values and the role
9 of “peakiness” of distributions. One observation that appears to be focused upon
10 and may be a driving force is the uneven distribution among the 15 urban sites on
11 the impact of the various scenarios and whether this fact is sufficiently taken into
12 account in scaling up for the national estimates. More discussion and or analyses
13 on this point may be warranted. For example what role does the actual estimates
14 from these 15 sites play in coming to the estimates of 3-9% excess mortality? It
15 may be too much to expect (in spite of the statistics) that 63,000-88,000
16 premature deaths would be prevented. Part of the country is already well below
17 the proposed alternative levels and thus would not contribute to lives saved. Are
18 there additional alternative sensitivity analyses that would provide either
19 alternative estimates or put more confidence in these estimates by taking into
20 account better population weighted C-R analyses?
21
- 22 c. See above. In spite of the last comment, the uncertainties and variability of the
23 core assessments seems to be as good as it can be.
24
- 25 d. Evaluation of the several national scale analyses, as indicated above is of some
26 concern. If I read the Tables in Appendix E correctly, the effect of moving to the
27 lowest alternative (25/12) in some cases within the 15 urban sites produces a
28 range of 32-67% (with one outlier at 11% and one at 100%) reduction in the IHD
29 compared to the current standard. The question is, is this the best baseline for the
30 comparison or should it be the current recent measurements, which would drop
31 the percent changes considerably (and perhaps provide a more realistic estimated
32 of the potential benefits from implementing changes). Obviously, the
33 proportional ranking and changes would be the same, but the impact on “lives
34 saved” on a national scale might be considerably less and more realistic.
35
- 36 e. Key observations seem to be presented in a balanced and fair way. Although the
37 national assessment suggests a range of 63,000-88,000 premature deaths per year
38 attributable to PM2.5 does not jive with a fairly often quoted figure from 2006
39 that moving the annual standard from 15 to 14 ppm would result in “more lives
40 than perished in 9/11”. (That figure translated into about 3000 lives.) Staff
41 acknowledges that the range of effects are in two categories: 3-9% and 0-3% in
42 two halves of the country. This may be less precise than what the data indicate in
43 that it would appear from their own estimates that the bulk of the effect comes from
44 the upper end of the exposure in the counties (pg 5.8, line 12-15).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43

Comments from Dr. Helen Suh:

The Second Draft of the Risk Assessment is a clear and comprehensive presentation of the rationale, methods and results for the assessment the acute and chronic PM-mediated health risks. In this draft, the authors have thoughtfully considered and incorporated the comments and suggestions from the CASAC panel. As an overall comment, the rationale that was used in the decision to forego a risk assessment for PM_{10-2.5} made logical sense. Further, inclusion of a discussion of PM_{10-2.5} exposure and risk in the upcoming draft Policy Assessment document is welcomed. However, it still seems that a qualitative and indirect assessment of PM_{10-2.5} risks could be made based on PM₁₀ monitoring and health data for metropolitan areas where PM_{10-2.5} concentrations comprise a large fraction of PM₁₀. While certainly not ideal, this qualitative assessment may provide valuable information about PM_{10-2.5} risks.

Charge Question 2: Selection of model inputs (section 3.3): We have expanded and clarified the discussion of our rationale for identifying modeling choices comprising the core risk model, focusing in particular on selection of C-R functions (section 3.3.3). To what extent does the Panel consider this discussion to be clear and the model selections appropriate?

The revisions to section 3.3 represent a significant improvement over the previous draft, with the selection rationale for the core risk model clearly and cogently presented and previous concerns addressed. The rationale provided is a thoughtful and sensible approach to assess particle-mediated health risks. Further, the summary tables (Table 3.5 - 3.8) provide a useful and nice synopsis of the model inputs for the core risk models and sensitivity analyses. As a very minor comment, it might be possible to condense Table 3.7 somewhat by replacing certain columns with check boxes instead of text (for example to indicate short-term or long-term).

Charge Question 3: Addressing uncertainty and variability (section 3.5): We have clarified the process used to evaluate sources of variability and added coverage for specific sources of variability (section 3.5.2); expanded our discussion of the qualitative analysis of uncertainty (section 3.5.3); and included analyses of pair-wise interactions of sources of uncertainty (section 3.5.4). To what extent does the Panel consider these discussions to be clear and appropriate?

The discussions of uncertainty and variability were clear and thoughtful, representing a substantial improvement over the previous draft. The sources of variability and uncertainty were well represented, although some explicit discussion of the impact of seasonality on risks should be included given its effects on each of the other discussed sources.

1 **Comments from Dr. Sverre Vedal:**

2
3 **PM Risk Assessment, Charge Question 6.**

4
5 **General:**

6 1. There is a too much formulaic and detailed presentation of results, for example when
7 presenting findings of just meeting the current standards and just meeting alternative
8 standards. This seems out of place in an integrative discussion.

9 2. While there is some opinion that effects of long-term exposure on IHD mortality (as
10 opposed to all-cause cardiovascular mortality) are especially strong, this is based on
11 relatively weak evidence from the ACS cohort (Pope 2004), in my opinion. It is
12 therefore not clear why it was elected to present risk assessment results largely for IHD in
13 this discussion (p. 6-6) and in chapter 4 of the RA. On second thought, it is clear because
14 all-cause CVD mortality effects were not presented in Krewski 2009, unfortunately, and
15 Krewski 2009 was chosen to provide the most defensible effect estimates.

16 3. The remaining percent PM-attributable effect of long-term exposure on total mortality
17 is presented (p. 6-7, line 18; p. 6-8, line 8 and lines 14 & 15). These are ridiculously
18 high. Presumably this should be on IHD mortality. The first bullet under the first Key
19 Observation (p. 6-18) gets it right, I believe.

20 4. I question whether short-term exposure-related risk (p. 6-5, line 3) is also driven by
21 changes in long-term average PM concentrations. Short-term effects are observed
22 independent of long-term PM concentration, ie, down to the lowest baseline
23 concentrations. Therefore the argument in this paragraph for motivating attention to the
24 annual average is not sound.

25
26 **6.a. Key questions captured?**

27 Yes

28
29 **6.b. Role of additional factors?**

30 Observations in 6.2 are well-supported. No other observations are apparent.

31
32 **6.c. Characterization of confidence.**

33 Again (see above), the emphasis on IHD mortality as opposed to all-cause cardiovascular
34 mortality reduces our confidence in effect estimates; unfortunately, all-cause
35 cardiovascular effect estimates are not provided in Krewski 2009. IHD effect estimates
36 are highest in the ACS cohort. Otherwise, estimates are conservative.

37
38 **6.d. National context.**

39 I'm not sure I agree that the 15 cities capture "the overall distribution of risk for the
40 nation," (p.6-16, line 36) given the fact that these are the more polluted cities in the
41 nation. More correct is the contention that these cities reflect the experience of cities
42 with relatively elevated levels of PM-related risk and attributable mortality. Both can't
43 be true.

44
45 **6.e. Key observations.**

3-30-10 Deliberative Draft Letter on the Quantitative Health Risk Assessment for
Particulate Matter (Second External Review Draft). DO NOT CITE OR QUOTE.

1 These observations are fine. I see little utility in estimating risk below an annual standard
2 of 12 mcg/m³.
3
4
5
6