

Nitrogen Cycles: Past, Present and Future

JAMES N. GALLOWAY^{1*}, FRANK J. DENTENER², DOUGLAS G. CAPONE³,
ELISABETH W. BOYER⁴, ROBERT W. HOWARTH⁵, SYBIL P. SEITZINGER⁶,
GREGORY P. ASNER⁷, CORY CLEVELAND⁸, PAMELA GREEN⁹, ELISABETH
HOLLAND¹⁰, DAVID M. KARL¹¹, ANTHONY F. MICHAELS³, JOHN H. PORTER¹,
ALAN TOWNSEND⁸ and CHARLES VÖRÖSMARTY⁹

¹Environmental Sciences Department, University of Virginia, Charlottesville, Virginia, USA; ²Joint Research Centre, Institute for Environment and Sustainability Climate Change Unit, Ispra, Italy; ³Wrigley Institute for Environmental Studies, University of Southern California, Los Angeles, California, USA; ⁴College of Environmental Science and Forestry, State University of New York, Syracuse, New York, USA; ⁵Department of Ecology & Evolutionary Biology, Cornell University, Ithaca, New York, USA; ⁶Institute of Marine and Coastal Sciences, Rutgers, The State University of New Jersey, New Brunswick, New Jersey, USA; ⁷Department of Global Ecology, Carnegie Institution, Stanford University, Stanford, California, USA; ⁸Institute of Arctic and Alpine Research, University of Colorado, Boulder, Colorado, USA; ⁹Complex Systems Research Center, Institute for the Study of Earth, Oceans, and Space, University of New Hampshire, Durham, New Hampshire, USA; ¹⁰Atmospheric Chemistry Division, National Center for Atmospheric Research, Boulder, Colorado, USA; ¹¹School of Ocean and Earth Science and Technology, University of Hawaii, Honolulu, Hawaii, USA; *Author for correspondence (email: jng@virginia.edu; phone: 434-924-1303; fax: 434-982-2300)

Key words: nitrogen, Haber-Bosch, global cycles, fertilizer fossil fuel combustion, denitrification, nitrogen fixation, Asia

Abstract. This paper contrasts the natural and anthropogenic controls on the conversion of unreactive N₂ to more reactive forms of nitrogen. A variety of data sets are used to construct global N budgets for 1860 and the early-1990s, and make projections for the global N budget in 2050. Regional N budgets for Asia, North America and other major regions for the early-1990s, as well as the marine N budget, are presented to highlight the dominant fluxes of nitrogen in each region. Important findings are that human activities increasingly dominate the N budget at the global and most regional scales, which the terrestrial and open ocean N budgets are essentially disconnected, and the fixed forms of N are accumulating in most environmental reservoirs. The largest uncertainties in our understanding of the N budget at most scales are the rates of natural BNF, the amount of N_r storage in most environmental reservoirs and the production rates of N₂ by denitrification.

Introduction

Water, water everywhere, nor any drop to drink;
Water, water everywhere and how the boards did shrink.

This couplet from the Rime of the Ancient Mariner (Samuel Taylor Coleridge, 1772 - 1834) is an observation that although sailors were surrounded by water, they were dying of thirst because of its form. Just as water is a critical substance for life, so is nitrogen. And just as most of the water on the planet is not useable by most organisms, most of the nitrogen is also unavailable. Approximately 78% of the atmosphere is diatomic nitrogen (N_2) which is unavailable to most organisms because of the strength of the triple bond that holds the two nitrogen atoms together. Over evolutionary history, only a limited number of species of Bacteria and Archaea have evolved the ability to convert N_2 to reactive nitrogen (Nr)¹ and by to use nitrogen efficiently. However, even with adaptations to use nitrogen efficiently, many ecosystems of the world are limited by nitrogen.

To place the current alteration of N cycle into historical context, we begin this paper with a brief overview of the development of human understanding of the nitrogen cycle. We use as a primary reference Smil (2001) which thoroughly documents the history of N as part of a discussion of the Haber-Bosch process.

Jean Antoine Claude Chaptal (1756-1832) formally named the 7th element of the periodic table in 1790. By the beginning of the second half of the 19th century it was known that N was a common element in plant and animal tissues, that it was indispensable for plant growth, that there was constant cycling between organic and inorganic compounds, and that it was an effective fertilizer. However, the source of nitrogen was uncertain. Lightning and atmospheric deposition were thought to be the most important sources. Although the existence of biological nitrogen fixation (BNF) was unknown, in 1838 Boussingault demonstrated that legumes could restore Nr to the soil and that somehow they must create Nr directly. It was 50 more years before the puzzle was solved. In 1888 Herman Hellriegel (1831-1895) and Hermann Wilfarth (1853-1904) published their work on microbial communities: “The Leguminosae do not themselves possess the ability to assimilate free nitrogen in the air, but the active participation of living micro-organisms in the soil is absolutely necessary” (Smil, 2001). They went on to say that it was necessary that there was a symbiotic relationship between legumes and micro-organisms. Also around this time, the processes of nitrification and denitrification were identified, so by the end of the 19th century, the essential components of the nitrogen cycle were in place.

Over the past 100 years, our knowledge of Nr creation and its movement through ecosystems and environmental reservoirs has increased dramatically. We know that Nr

¹ The term reactive nitrogen (Nr) as used in this paper includes all biologically active, photochemically reactive, and radiatively active N compounds in the atmosphere and biosphere of the Earth. Thus Nr includes inorganic reduced forms of N (e.g., NH_3 , NH_4^+), inorganic oxidized forms (e.g., NO_x , HNO_3 , N_2O , NO_3^-), and organic compounds (e.g., urea, amines, proteins). Note that this definition is much broader than the term ‘reactive N’ as defined by the atmospheric chemistry community—they define reactive N as NO_x , which is any N-O combination except N_2O (e.g., NO_x , N_2O_5 , HNO_2 , HNO_3 , nitrates, organic nitrates, halogen nitrates, etc).

creation occurs in a number of ecosystems (via BNF) as well as by lightning. We also know that the productivity of many ecosystems is controlled by N availability (Vitousek et al., 2002). While this limitation is part of the natural process, it was not tenable for a growing human population that needed increasing amounts of Nr to grow food. This demand has resulted in a very significant alteration of the N cycle in air, land and water and at local, regional and global scales.

Two anthropogenic activities have greatly increased Nr availability. The first is food production. Early hunter-gatherer peoples were able to meet their nitrogen requirements by consuming protein from wild plants and animals. However, the establishment of settled communities ~10,000 years ago required the ability to 'grow your own'. Archeological evidence points to legume cultivation over 6,500 years ago (Smith 1995). Rice cultivation began in Asia perhaps as early as 7,000 years ago (Wittwer et al. 1987), and soybeans have been cultivated in China for at least 3,100 years (Wang 1987). These crops resulted in anthropogenic-induced creation of Nr since legumes can self-fertilize via symbioses with N-fixing organisms and rice cultivation creates anaerobic environments that encouraged high rates of BNF by cyanobacteria.

The annual per-area rates of transfer of atmospheric N₂ to Nr by cultivation can be large compared to natural rates of transfer. As thoroughly reviewed by Smil (1999), *Rhizobium* associated with seed legumes (e.g., peas, and beans) can fix N at rates ranging from 3×10^2 to 3×10^4 mg N m⁻². Most fixation rates are on the order of 4×10^3 to 3×10^4 mg N m⁻². *Rhizobium* associated with leguminous forages (e.g., alfalfa, clover) have higher average rates, 1×10^4 to 2×10^4 mg N m⁻². Non-*Rhizobium* N-fixing organisms associated with some crops (e.g., cereals) and trees have ranges from 5×10^2 to 2×10^3 mg N m⁻², while cyanobacteria associated with rice paddies, and endophytic diazotrophs associated with sugar cane can fix 2×10^3 to 3×10^3 mg N m⁻² and 5×10^3 , respectively (Smil 1999).

The second anthropogenic activity that increased Nr was energy production. While food production creates additional Nr on purpose, energy production creates it by accident (H. Levy, pers. comm., 1995). During combustion of fossil fuels nitrogen is emitted to the atmosphere as a waste product (NO) from either the oxidation of atmospheric N₂ or organic N in the fuel (primarily coal) (Socolow, 1999; Galloway et al., 2002). The former creates new Nr; the latter mobilizes sequestered Nr. The magnitude of Nr mobilization due to energy production is not as extensive as that from food production. Although there are records of coal use dating from 500 BC in China, up until the late 19th century most energy was produced from biofuels (e.g., wood). It wasn't until the beginning of the 20th century that fossil fuels overtook biomass fuels in supplying primary energy (Smil 1994).

By the beginning of the 20th century the importance of nitrogen in food production had been established and the major components of the N cycle had been identified. In addition, both legume/rice cultivation and fossil fuel combustion were creating Nr: the former as a means to provide N to produce food, and the latter as a consequence of energy production. Many realized that there was not enough nitrogen available from naturally occurring sources to provide food for a growing global population (Smil 2001). The only sources for 'new' N at that time (in addition to cultivation) were guano deposits on arid islands and evaporite nitrate deposits in South America (Chile), that supplied about 0.2 Tg N yr⁻¹ (Smil 2000). This was not enough.

The pressure to obtain additional Nr for food production (and the need for nitrate to produce munitions) led to the 1913 development of the Haber-Bosch process in Germany to produce NH₃ from N₂ and H₂ (Smil 2001).

We are now at the beginning of the 21st century. Food and energy production have grown with population and, in some regions, with significant increases in the per-capita resource use. Currently, food and energy production has increased the anthropogenic Nr creation rate by over a factor of ten compared to the late-19th century. The magnitude of this production raises critical questions as to the consequences and fate of new Nr in the environment. Answers to these questions are problematic. With seven oxidation states, numerous mechanisms for interspecies conversion, and a variety of environmental transport/storage processes, nitrogen has arguably the most complex cycle of all the major elements. This complexity makes tracking anthropogenic nitrogen through environmental reservoirs a challenge. However, such work is necessary because of nitrogen's role in all living systems and in several environmental issues (e.g., greenhouse effect, smog, stratospheric ozone depletion, acid deposition, coastal eutrophication and productivity of freshwaters, marine waters and terrestrial ecosystems).

The analysis of the magnitude and consequences of human intervention in the N cycle is not new. More than thirty years ago Delwiche (1970) stated that humans were mobilizing about the same amount of N as natural processes, and that the fate of the new Nr was uncertain. Since Delwiche's seminal work, anthropogenic Nr creation has doubled, while natural terrestrial BNF has decreased due to land use change. Many uncertainties remain and have become all the more important to resolve.

In the last several years, a number of recent papers have addressed the N cycle on the global scale (e.g., Ayres et al. 1994; Mackenzie 1994; Galloway et al. 1995; Vitousek et al. 1997; Galloway 1998; Seitzinger and Kroeze 1998; Galloway and Cowling 2002), the regional scale (Asia—Galloway 2000; Zheng et al. 2002; Bashkin et al. 2002; North Atlantic Ocean and watershed—Galloway et al. 1996; Howarth et al. 1996; oceans—Karl 1999; Capone 2001; Europe—von Egmond et al. 2002; United States—Howarth et al. 2002), and on major components of the N cycle (food production—Smil 1999, 2002; Oenema and Pietrzak 2002; Cassman et al. 2002; Roy et al. 2002; fertilizer production—Fixen and West 2002; fossil fuel combustion—Bradley and Jones 2002; Moomaw 2002; industrial uses of Nr—Febre Domene and Ayres 2001), the atmosphere—Holland et al. 1999; BNF—Cleveland et al. 1999; Karl 1999, 2002; Vitousek et al. 2002), and on the relationship to public policy (Socolow 1999; Mosier et al. 2001; Melillo and Cowling 2002). Using these previous papers as a foundation, and utilizing the results of the multi-year SCOPE project (Boyer and Howarth 2002) and the findings of the Second International Nitrogen Conference (Galloway et al. 2002) this paper addresses the following questions:

- How has the global nitrogen budget changed from the late 19th century to the late 20th century?
- What is the global N budget projected to be in the mid-21st century?
- How have atmosphere-biosphere N exchanges been altered by human activity?
- What are the connections between the terrestrial and marine N cycles?
- How much of the Nr created by human activity is denitrified back to N₂?

These are important questions to address. Nr influences biogeochemical processes in the atmosphere, in terrestrial ecosystems and in freshwater and marine

aquatic ecosystems. Increases in the concentration of Nr species can enhance ecosystem productivity through fertilization or decrease it through nutrient imbalances, and decrease ecosystem biodiversity through acidification and eutrophication (Vitousek et al. 1997; Aber et al. 1998; NRC 2000; Matson et al. 2002; Rabalais 2002; Tartowski and Howarth 2000). Higher Nr concentrations in the atmosphere can increase the incidence of air pollution-related illness due to O₃ and particulate matter inhalation (Follett and Follett 2001; Wolfe and Patz 2002; Townsend et al. 2003). A unique aspect of the impact of Nr on the environment and on people is that the effects can occur in series. Referred to as the nitrogen cascade (Galloway et al. 2003), one atom of nitrogen can, in sequence, increase atmospheric O₃ (human health impact), increase fine particulate matter (visibility impact), alter forest productivity, acidify surface waters (biodiversity loss), increase coastal ecosystem productivity, promote coastal eutrophication and increase greenhouse potential of the atmosphere (via N₂O production). The magnitude of the consequences, coupled with the magnitude of current rates of Nr creation makes the issue of Nr accumulation an important one to address.

The paper begins with the global N budget and the primary natural process that creates Nr—BNF. After assessing natural terrestrial rates of Nr creation, the paper addresses anthropogenic Nr creation rates in 1860 and the early-1990s (defined as 1990 to 1995, depending on the data set). The next two sections track Nr through global terrestrial systems for both time periods. The next section addresses the N budget on regional scales as geopolitical units. Such an analysis illustrates the spatial heterogeneity in both Nr creation and distribution. The paper then presents an assessment of the marine component of the N budget and its linkages with the terrestrial component. The final section discusses projections for the N budget in 2050.

The budgets presented in this paper are constructed from data previously published (or data previously published but adjusted for the time periods covered in this paper), and data that are presented for the first time. In general, previously published data and time-adjusted data are: Nr creation by Haber-Bosch process; terrestrial BNF; cultivation-induced BNF; atmospheric emission for NO_x, NH₃ and N₂O; and NO_y and NH_x deposition. The marine N budget terms are based upon an assessment presented in this paper using data from a number of sources. The riverine fluxes represent a new analysis and are based primarily on other data bases used in this paper. Estimates for N₂ production via denitrification from soils, rivers and estuaries are based upon previously published data as well as assessments presented in this paper. Additional information about data sources is provided in the appropriate sections.

The data and model results used in this paper have differing levels of uncertainty, which we have presented to the degree possible. Because of these uncertainties, we generally present fluxes to three significant figures or 0.1 Tg N yr⁻¹. This practice at times produces sums that are slightly different, due to rounding errors, from what would be obtained by numerical addition. The uncertainty associated with most fluxes is discussed within the paper. The uncertainties associated with the atmospheric NO_y and NH_x budgets (emissions, transport, transformation, deposition) are discussed in Appendix I).

Terrestrial Nr Creation

Natural

Lightning

High temperatures occurring in lightning strikes produce NO in the atmosphere from molecular oxygen and nitrogen. Subsequently this NO is oxidized to NO₂ and then to HNO₃, and quickly (i.e., days) removed by wet and dry deposition, thus introducing Nr into ecosystems primarily over tropical continents. Most current estimates of Nr creation by lightning range between 3-10 Tg N yr⁻¹ (Prather et al. 2001). In this analysis we use a global estimate of 5.4 Tg N yr⁻¹ (Lelieveld and Dentener 2000) (Table 1). While this number is small relative to terrestrial BNF, it can be important for regions that do not have other significant Nr sources. It is also important because it creates NO_x high in the free troposphere, compared to NO_x emitted at the earth's surface. As a result it has a longer atmospheric residence time and is more likely to contribute to tropospheric O₃ formation, which significantly impacts the oxidizing capacity of the atmosphere.

BNF

Problems and Uncertainties

Quantifying the magnitude of natural, terrestrial Nr creation by BNF is tenuous, owing most notably to uncertainty and variability in the estimates of rates of BNF at the plot scale. Specifically, methodological differences, uncertainties in spatial coverage of important N-fixing species, and locational biases in the study of BNF all suggest critical gaps in our understanding of natural BNF at large scales (Cleveland et al. 1999). In addition, for many large areas where BNF is likely to be important, particularly in the tropical regions of Asia, Africa, and South America, there are virtually no data on natural terrestrial rates of BNF. In a recent compilation of rates of natural BNF by Cleveland et al. (1999), symbiotic BNF rates for several biome types are based on one-to-few published rates of symbiotic BNF at the plot scale within each particular biome. For example, based on the few estimates of symbiotic BNF available for tropical rain forests, estimated BNF in these systems represents ~24% of total natural terrestrial BNF globally on an annual basis (Cleveland et al. 1999). While the relative richness of potential N₂-fixing legumes in tropical forests suggests that symbiotic BNF in these systems is relatively high (Crews 1999), the paucity of actual BNF rate estimates in these systems suggests caution when attempting to extrapolate plot scale estimates of BNF, and highlight the difficulties of attempting to estimate natural BNF at the global scale.

Previous Estimates

Difficulties notwithstanding, prior estimates of BNF in terrestrial ecosystems range from ~40-200 Tg N yr⁻¹ (e.g. Schlesinger 1991; Söderland & Rosswall 1982; Stedman & Shetter 1983; Paul & Clark 1997). Most studies merely present BNF estimates as "global values" that, at best, are broken into a few very broad components (e.g. "forest", "grassland", and "other"; e.g. Paul & Clark 1997). Such coarse divisions average enormous land areas that contain significant variation in both BNF data sets and biome types, thus diminishing their utility. Many studies do not list data sources from which their estimates were derived (e.g. Schlesinger 1991; Söderland & Rosswall 1982; Stedman & Shetter 1983; Paul & Clark 1997). In contrast, Cleveland et al. (1999) provided a range of estimates of BNF in natural ecosystems from 100-290 Tg N yr⁻¹

(with a "best estimate" of 195 Tg N yr⁻¹). These estimates were based on published, data-based rates of BNF in natural ecosystems, and differ only in the percent cover estimates of symbiotic N fixers used to scale plot-level estimates to the biome scale.

Current Estimates

While the data-based estimates of Cleveland et al. (1999) provide more documented, constrained range of terrestrial BNF, there are several compelling reasons to believe that an estimate in the lower portion of the range is more realistic than higher estimates. First, rate estimates of BNF presented in the literature are inherently biased, as investigations of BNF are frequently carried out in areas where BNF is most likely to be important (i.e. where there are large assemblages of N-fixing species that don't reflect average community composition for the entire biome). For example, many rates of BNF in temperate forests were derived from studies that include N inputs from alder and black locust (Cleveland et al., 1999). While rates of BNF may be very high within stands dominated by these species (Binkley et al. 1994; Boring and Swank 1984), these species are certainly not dominant in temperate forests as a whole (Johnson & Mayeux 1990). Similarly, while species with high rates of BNF are often common in early successional forests (Vitousek 1994), they are often rare in mature or late successional forests, especially in the temperate zone (Gorham et al. 1979; Boring & Swank, 1984; Blundon & Dale 1990). Literature derived estimates based on *reported* coverage of N-fixing species are thus inflated due to these inherent biases.

We suggest that annual global BNF contributed between 100 and 290 Tg N yr⁻¹ to natural terrestrial ecosystems prior to large-scale human disturbance. However, we contend that due to the inherent biases noted in plot-scale studies of N fixation rates, the true rate of BNF lies at the lower end of this range. Thus, we used actual evapotranspiration (ET) values generated in Terraflux (Asner et al., 2001; Bonan, 1996) and the strong, positive relationship between ET and BNF (Cleveland et al. 1999) to generate a new, single estimate of BNF prior to large-scale, human disturbance. Our new analysis is based on the relationship between ET and BNF, but uses rates of BNF calculated using the low percent cover values of symbiotic N fixers over the landscape (i.e. 5%; Cleveland et al. 1999). This analysis suggests that within the range of 100 Tg N yr⁻¹ to 290 Tg N yr⁻¹, natural BNF in terrestrial ecosystems contributes 128 Tg N yr⁻¹. This value is supported by an analysis comparing BNF to N requirement (by biome-type). Using the Cleveland et al. (1999) relationship between ET and BNF, a global N fixation value of 128 Tg N yr⁻¹ would suggest an average of ~15% of the N requirement across all biome types is met via BNF; higher estimates of BNF would imply that at least 30% of the N requirement across all biomes is met via natural BNF (Asner et al. 2001). However, BNF in even the most active leguminous crop species frequently accounts for < 30% of total plant N (Peoples et al. 1995).

Our estimate (128 Tg N yr⁻¹) represents potential BNF prior to large-scale human disturbance, and does not account for decreases in BNF due to land use change or decreases in BNF due to other physical, chemical, or biological factors. To estimate natural terrestrial BNF for 1860 and the early-1990s, we scale BNF to the extent of altered land at those two times. Of the 11,500 million ha of natural vegetated land (Mackenzie 1998), Houghton and Hackler (2002) estimate that in 1860 and 1995, 760 million ha and 2,400 million ha, respectively, had been altered by human action (e.g.,

cultivation, conversion of forests to pastures). Therefore, in this analysis of BNF in the natural terrestrial landscape, we use 128 Tg N yr⁻¹ for the natural world, 120 Tg N yr⁻¹ for 1860, and 107 Tg N yr⁻¹ for the present world.

Summary

The conversion of N₂ to Nr requires energy to break the N:N triple bond. In the natural world, physical (lightning) and biological (BNF) processes provide this energy. Nr creation by lightning is highest in tropical terrestrial regions, where convective activity is the largest. BNF rates in terrestrial systems are also generally highest in tropical regions where late-successional tropical forests often contain many potentially N-fixing canopy legumes (Cleveland et al. 1999; Vitousek et al. 2002). Thus, relative to temperate regions tropical regions are important source areas of Nr due in part to ecosystem structure and energy availability. As will be seen in the next section, human Nr creation in many temperate regions is primarily controlled by a different factor—the use of fossil energy to produce energy and fertilizer. Thus the latitude dependency changes from one driven by solar intensity and ecosystem type, to one driven by population density and industrial productivity.

Anthropogenic

1860

Van Aardenne et al. (2001) estimated 0.6 Tg N yr⁻¹ of Nr was created in the form of NO_x during fossil fuel combustion in 1890, primarily from coal combustion. Scaling this estimate by population and other factors, we estimate that in 1860 the equivalent value was ~0.3 Tg N yr⁻¹. Since the Haber-Bosch process was not yet invented, the only new N created by food production was by cultivation of legumes. Galloway and Cowling (2002) estimate, using assessments by V. Smil (pers. comm.), that ~15 Tg N yr⁻¹ was produced in 1900 by cultivation-induced BNF. Given the uncertainty about this estimate, we believe it to be reasonable to use this value to represent conditions in 1860.

Thus prior to the 20th century, humans created new Nr almost entirely to support food production. The total anthropogenic Nr produced (~15 Tg N yr⁻¹) was small relative to BNF occurring in unmanaged terrestrial ecosystems (120 Tg N yr⁻¹) (Table 1).

Early-1990s

Between 1860 and 1995 the world's population increased ~4.5-fold, from 1.3 to 5.8 billion. Cultivation-induced Nr creation increased by only ~2-fold from ~15 Tg N yr⁻¹ in 1860 to ~33 Tg N yr⁻¹ in the mid-1990s (Smil 1999). Symbiotic biofixation by *Rhizobium* associated with seed legumes resulted in 10 Tg N yr⁻¹ (8-12 Tg N yr⁻¹) of new nitrogen. Biofixation by leguminous cover crops (forages and green manures such as clover, alfalfa, vetches) accounted for an additional 12 Tg N yr⁻¹ (10-14 Tg N yr⁻¹) of new nitrogen. As Smil notes, biofixation by non-*Rhizobium* N-fixing species was of less importance, fixing on the order of 4 Tg N yr⁻¹ (2-6 Tg N yr⁻¹). Cyanobacteria fixed on the order of 4-6 Tg N yr⁻¹ in wet-rice fields, while endophytic N-fixing organisms in sugar cane fixed an additional 1-3 Tg N yr⁻¹. The global total from cultivation is thus ~33 Tg N yr⁻¹ within a range of 25-41 Tg N yr⁻¹ (Smil 1999). When we applied Smil's crop-specific mean fixation rates to the crop area data on a regional basis from FAO (2002), we

estimate that for 1995 that total global C-BNF was 31.5 Tg N yr⁻¹, very similar to Smil's value of 33 Tg N yr⁻¹ (Table 1).

Relative to cultivation-induced BNF, about three times as much Nr was created with the Haber-Bosch process. In 1995, 100 Tg N of NH₃ was created for food production and other industrial activities (Kramer 1999). Of this amount, about 86% (~86 Tg N yr⁻¹) was used to make fertilizers. The remaining 14 Tg N yr⁻¹ was dispersed to the environment during processing or used in the manufacture of synthetic fibers,

refrigerants, explosives, plastics, rocket fuels, nitroparaffins, etc. (Smil 1999; Febre Domene and Ayres 2001). As with the production of fertilizer, this also represents creation of new Nr that is introduced into environmental systems.

The increase in energy production by fossil fuels resulted in increased NO_x emissions from 0.3 Tg N yr⁻¹ in 1860 to ~24.5 Tg N yr⁻¹ in the early-1990s—by the early-1990s over 90% of energy production resulted in the creation of new reactive nitrogen, contrasting to 1860 where very little of energy production caused creation of Nr.

Summary

In the early-1990s, Nr creation by anthropogenic activities was ~156 Tg N yr⁻¹, a factor of ~10 increase over 1860, contrasted to only a factor of ~4.5 increase in global population (Table 1). Food production accounted for ~77%, energy production accounted for 16%, and production for industrial uses accounted for ~9%.

Global Terrestrial N Budget Introduction

This section examines the extent of Nr distribution via atmospheric and hydrologic pathways. It provides a context to evaluate the extent to which human

Table 1: Global Nr creation and distribution, Tg N yr⁻¹

	1860	early-1990s	2050	Notes
Nr Creation				1
Natural				
Lightning	5.4	5.4	5.4	
BNF-terrestrial	120	107	98	
BNF-marine	121	121	121	
subtotal	246	233	224	
Anthropogenic				
Haber-Bosch	0	100	165	
BNF-cultivation	15	31.5	50	
Fossil fuel combustion	0.3	24.5	52.2	
subtotal	15	156	267	
Total	262	389	492	
Atmospheric emission				2
NO_x				
fossil fuel combustion	0.3	24.5	52.2	
lightning	5.4	5.4	5.4	
Other emissions	7.4	16.1	23.9	
NH₃				
terrestrial	14.9	52.6	113	
marine	5.6	5.6	5.6	
N₂O				
terrestrial	8.1	10.9	13.1+?	
marine	3.9	4.3	5.1	
Total (NO_x and NH₃)	13.1	46	82	
Atmospheric deposition				3
NO_y				
Terrestrial	6.6	24.8	42.2	
Marine	6.2	21	36.3	
subtotal	12.8	45.8	78.5	
NH_x				
Terrestrial	10.8	38.7	83	
Marine	8	18	33.1	
subtotal	18.8	56.7	116.1	
Total	31.6	103	195	
Riverine fluxes				4
Nr input into rivers	69.8	118.1	149.8	
Nr export to inland systems	7.9	11.3	11.7	
Nr export to coastal areas	27	47.8	63.2	
Denitrification				5
Continental				
Terrestrial		67	95	
Riverine		47.8	63.2	
subtotal	98	115	158	
Estuary and Shelf				
riverine nitrate	27	47.8	63.2	
open ocean nitrate	145	145	145	
subtotal	172	193	208	
Open Ocean	129	129	129	
Total	125	163	221	

Notes

- Nr Creation**
 'BNF-terrestrial'--based on Cleveland et al (1999) as discussed in the text.
 'BNF-marine'--Table 8 (average of the minimum and maximum values).
 'Lightning'--Lalleveld and Dentener (2000)
 'Haber-Bosch'--early-1990s (Kramer 1999); 2050 (see text)
 'BNF cultivation'--based on Smil (1999; personal communication)
 'Combustion'--Klein Goldewijk and Batjes (1997); van Aardenne et al. (2001)
- Atmospheric Emission**
 'NO_x (other emissions)'--Klein Goldewijk and Batjes (1997); van Aardenne et al. (2001)
 'NH₃'--see text
 'N₂O'--see text
- Atmospheric Deposition**
 'NO_x and NH_x'--see text
- Riverine fluxes**
 'Nr inputs in to rivers'--assumed to be twice riverine discharge. Range is 30-70% (Seitzinger 2000.)
 'River Fluxes'--see text and Appendix II.
- Denitrification**
 'Terrestrial'--1860, no storage; 1990s, nat denit reduced, 0.25 of anthro Nr denit, & excess river N lost
 'Riverine'--assumed to be equal to difference between 'river input' and 'river discharge'
 'Estuary and Shelf'--Shelf estimate (Table 8) minus 1860 'riverine discharge'
 'Open Ocean'--Table 8 (average of the minimum and maximum values).

intervention in the N cycle in the early-1990s has substantially changed N distribution on a global and regional basis. The global fluxes presented in this section represent an update and an expansion over those presented in Galloway and Cowling (2002).

1860

Fixation

As discussed above, natural rates of Nr creation in 1860 were 120 Tg N yr⁻¹ by BNF and 5.4 Tg N yr⁻¹ by lightning. Anthropogenic Nr creation rates were 0.3 Tg N yr⁻¹ and ~15 Tg N yr⁻¹ by fossil fuel combustion and cultivation-induced BNF, respectively. Total terrestrial Nr creation was ~141 Tg N yr⁻¹ (Figure 1a, Table 1).




Figure 1a: Components of the global nitrogen cycle for 1860. All shaded boxes represent reservoirs of nitrogen species in the atmosphere.. Creation of Nr is depicted with bold arrows from the N₂ reservoir to the Nr reservoir (depicted by the dotted box). 'N-BNF' is biological nitrogen fixation within natural ecosystems; 'C-BNF' is biological nitrogen fixation within agroecosystems. Denitrification creation of N₂ from Nr within the dotted box is also shown with bold arrows. All arrows that do not leave the dotted box represent inter-reservoir exchanges of Nr. The dashed lines within the dotted box associated with NH_x represent natural emissions of NH₃ that are re-deposited on fast time-scales to the oceans and continents.

Emission and Deposition

Atmospheric emissions for 1860 and 1993 (except for marine NH₃ emissions) were derived from a 1° x 1° data-base (van Aardenne et al. 2001) that covered 1890 to 1990. The adjustment to 1860 used estimated activity data (e.g. population) together with emission factors for 1890. For 1993 we extrapolated 1990 anthropogenic emissions using the reported increase of CO₂ emissions for the period 1990-1993 (Brenkert 1997). Emission factors were assumed to be unchanged. Marine NH₃ emissions are based on the monthly oceanic NH₄⁺ and pH values derived from the HAMOCC3 biological ocean model (for further information see Bouwman et al. (1997) and Quinn et al. (1996)) and

model-calculated exchange velocities. The atmospheric deposition data on a 5° by 3.75° grid were generated from a global transport-chemistry model (Lelieveld and Dentener 2000). Each grid was subdivided into a 50 km x 50 km sub-grid to create a spatial defined deposition map (Figure 2). The gridded data were assigned to continental and marine regions needed for this study using boundaries delineated on a world data coverage from ESRI (ESRI 1993). See Appendix I for a discussion of uncertainties in these estimates.

NO_x and NH₃ emissions can result from natural processes, food production and energy production. Although anthropogenic creation rate of Nr was only ~16% of that created naturally in terrestrial environments, in 1860 humans had an observable effect on atmospheric Nr emissions. For NO_x, natural emissions occur from soil processes, lightning, wildfires (biomass burning) and stratospheric injection. These amounted to a total of 10.5 Tg yr⁻¹ (Table 2a). Energy production resulted in total NO_x emissions of 0.6 Tg N yr⁻¹, (0.3 Tg N yr⁻¹ from fossil fuel combustion, and 0.4 Tg N yr⁻¹ from biofuel combustion). Food production-related combustion of agricultural waste yielded 0.9 Tg

Table 2a: Global atmospheric emissions of NH₃ and NO_x, Tg N yr⁻¹

	1860		1993		2050		Note
	NO _x	NH ₃	NO _x	NH ₃	NO _x	NH ₃	
Food							A1
sav	0.9	0.2	2.9	1.8	3.8	2.2	
agl	0.0	-	2.6	-	5.1	-	
agr	0.9	0.6	2.4	1.4	5.0	3.1	
def	0.2	0.2	1.1	1.4	0.9	1.1	
fer	-	0.0	-	9.7	-	15.7	
anm	-	5.3	-	22.9	-	69.4	
lan	-	0.1	-	3.1	-	7.2	
cro	-	0.2	-	4.0	-	7.7	
subtotal	2.0	6.6	9.0	44.3	14.8	106	
Energy							A2
e1*	0.3	0.0	20.4	0.1	39.0	0.2	
nrt&ots*	0.0	-	3.6	-	11.2	-	
aircraft*	0.0	-	0.5	-	2.0	-	
his	0.0	0.0	1.5	0.2	3.6	0.2	
bf3	0.4	0.7	1.3	2.6	1.3	1.7	
subtotal	0.6	0.7	27.2	2.9	57.0	2.1	
Natural							A3
aglnat	2.9	-	2.9	-	2.9	-	
lightning*	5.4	-	5.4	-	5.4	-	
firenat	1.6	1.6	0.8	0.8	0.8	0.8	
strat	0.6	-	0.6	-	0.6	-	
soil&veg	-	6.0	-	4.6	-	3.6	
ocean	-	5.7	-	5.6	-	5.6	
subtotal	10.5	13.3	9.7	11.0	9.7	10.0	
Emission, total	13.1	20.6	45.9	58.2	81.5	118	
Deposition, total	12.8	18.8	45.8	56.7	78.5	116	B1

NOTES

A. Emissions (van Aardenne et al., 2001)

1. Food --'sav' represents NO_x and NH₃ emissions from savannah burning, some fraction of which could be considered natural; 'agl' is NO_x emissions from agricultural soils; 'agr' is NO_x and NH₃ emissions from agricultural waste burning; 'def' is NO_x and NH₃ emissions from combustion, as part of deforestation; 'anm' is NH₃ emissions from agricultural animal waste; 'lan' is NH₃ emissions from humans, pets and waste water; 'cro' is NH₃ emissions from agricultural crops.

2. Energy --'e1' is NO_x and NH₃ emissions from fossil fuel burning; 'his' is NO_x and NH₃ emissions from industrial processes; 'nrt&ots' is NO_x emissions from non road transport and one specific industrial sector; 'aircraft' is NO_x emissions from stratospheric aircraft; 'bf3' is non road transport and one specific industrial sector from biofuel combustion. An asterisks means that the combustion process lead to the creation of new Nr.

3. Natural --'aglnat' is NO_x emissions from natural soils; 'lightning' is NO_x formation due to lightning; 'firenat' is NO_x and NH₃ emissions from natural burning at high latitudes; 'strat' is NO_x injection from stratosphere; 'soil&veg' is NH₃ emissions from natural soils, vegetation and wild animals, calculated using a compensation point (see text); 'oceans' is NH₃ emission from oceans calculated using a compensation point (see text).

B. Deposition (Lelieveld and Dentener, 2000)

1. Data are for wet and dry deposition of NO_y and NH₃, (Table 1).

N-NO_x yr⁻¹. Slash-and-burn of forests yielded 0.2 Tg N yr⁻¹, and the annual burning of savanna grass and shrubs emitted 0.9 Tg N yr⁻¹. In 1860, total NO_x emissions were 13.1 Tg N yr⁻¹ of which ~10.5 Tg N yr⁻¹ were from natural sources. Total NO_y deposition is ~12.8 Tg N yr⁻¹ (Table 2a), with 6.6 Tg N yr⁻¹ to continents and 6.2 Tg N yr⁻¹ to oceans (Table 1).

The equivalent analysis for NH₃ shows that energy production resulted in 0.7 Tg N yr⁻¹ from the combustion of biofuels (Table 2a). Food production resulted in NH₃ emissions to the atmosphere of 6.6 Tg N yr⁻¹, ~10-fold greater than energy production. The component parts were combustion of agricultural waste (0.6 Tg N yr⁻¹), forests (0.2 Tg N yr⁻¹), and savannas (0.2 Tg N

yr⁻¹). Low temperature NH₃ emissions were 5.3 Tg N yr⁻¹ from domestic animal waste, 0.1 Tg N yr⁻¹ from human sewage, and 0.2 Tg N yr⁻¹ from crops (Table 2a). Natural emissions of NH₃ from oceans and natural vegetation were calculated using a compensation point approach. This results in a net exchange of NH₃ between the atmosphere and the underlying surface (Conrad and Dentener 1999; Dentener and Crutzen 1994). These emissions were part of a rapid local cycling, meaning that most of the emissions are deposited within the same region (Table 2a), as illustrated with the dotted lines in Figure 1a. Deposition of NH_x to the continents from non-local sources was 4.8 Tg N yr⁻¹ and to the ocean, 2.3 Tg N yr⁻¹. Including the rapid deposition of the natural NH₃ emissions, these numbers would have been 10.8 Tg N yr⁻¹ and 7.9 Tg N yr⁻¹, to continents and oceans, respectively in 1860 (Table 1).

Given the contributions to atmospheric N emissions from energy and food production, in contrast to natural emissions, it is not surprising that a map of N deposition for 1860 to the earth surface shows the imprint of human activity (Figure 2a). N deposition was more pronounced near populated areas with rates up to few hundred mg N m⁻² yr⁻¹, while in more remote regions it is on the order of 50 to 100 mg N m⁻² yr⁻¹ for terrestrial region and 5 to 25 mg N m⁻² yr⁻¹ for marine regions. The largest deposition rates were in Asia (maximum on the order of 1,000 mg N m⁻² yr⁻¹), which is not surprising as it had 65% of the world's population. The evidence of atmospheric transport downwind from these areas is also evident (Figure 2a).

Another Nr species emitted to the atmosphere, N₂O, bears mention. In 1860




Figure 2a: Spatial patterns of total inorganic nitrogen deposition in 1860, mg N m⁻² yr⁻¹.

terrestrial systems contributed 6.6 Tg N yr⁻¹ (assuming the Bouwman et al. (1995) natural soil emissions for 1990 are applicable to 1860 conditions) (Table 2b). As noted in Kroeze et al. (1999), anthropogenic N₂O emissions from animal waste, cultivation-induced BNF, crop residue and from biomass burning were approximately 1.4 Tg N yr⁻¹. Rivers are estimated to have contributed another 0.05 Tg N yr⁻¹ (Seitzinger et al. 2000). Estuaries and marine shelves accounted for 0.02 Tg N yr⁻¹ and 0.4 Tg N yr⁻¹, respectively (Seitzinger et al. 2000),

and the open ocean contributed ~3.5 Tg N yr⁻¹ (Nevison et al. 1995) (Table 2b, Figure 1a). Thus total N₂O emissions were ~12 Tg N yr⁻¹ in 1860. N₂O is globally distributed and either accumulates in the troposphere or is lost to the stratosphere (Prather et al. 2001). While there are considerable uncertainties in the magnitude of N₂O emissions from any one source, a closed N₂O budget for the period 1500-1994 is obtained when emissions from all sources are used as input to an atmospheric model indicating that increases in atmospheric N₂O can be primarily attributed to direct and indirect emissions associated with changes in food production systems (Kroeze et al. 1999).

The atmospheric portion of this analysis of the global N cycle only addresses inorganic N (NO_y, NH_x and N₂O) and does not include oxidized atmospheric organic N

(i.e., organic nitrates (e.g., PAN), reduced atmospheric organic N (e.g., aerosol amines and urea), and particulate atmospheric organic N (e.g., bacteria, dust). The emission and deposition of organic N is probably a significant component of the atmospheric N cycle. Neff et al. (2002) estimate for the present time a range of 10 to 50 Tg N yr⁻¹ emitted and deposited on a global basis, with the range reflecting the substantial unresolved uncertainties. Human activities certainly contribute to organic N in the atmosphere both from direct emissions and indirect (e.g., organic nitrates formation due to increased NO_x emissions). However, due to the limited number of measurements the quantification of the influence of humans has not been possible (Neff et al. 2002).

Riverine Export

Our estimates of riverine Nr fluxes to inland receiving waters and to the coastal ocean are based on a modification of an empirical, mass-balance model relating net anthropogenic Nr inputs per landscape area (NANI) to the total flux of Nr discharged in rivers, originally established using data from land regions draining to the coastal zone of the North Atlantic Ocean (Howarth et al. 1996). The NANI model considered new inputs of Nr that are human controlled, including inputs from fossil-fuel derived atmospheric deposition, fixation in cultivated croplands, fertilizer use, and the net import (or export) in food and feed to a region. Subsequent studies have found that the form of the relationship holds when considering other world regions in the temperate zone (e.g., Boyer et al. 2002; Boyer et al. in prep). Further, a model inter-comparison by Alexander et al. (2002) found the NANI model to be the most robust and least biased of several models used to estimate N fluxes from a variety of large watersheds.

For this paper we use a modification (as described Appendix II and detailed in Boyer et al. (in prep.) of the NANI model to quantify riverine Nr export from world regions for 1860, 1990, and 2050. The modification considers new inputs of Nr to a region from natural BNF in forests and other non-cultivated vegetated lands in addition to anthropogenic Nr inputs: the net total nitrogen inputs per unit area of landscape (or NTNI, which includes anthropogenic plus natural N inputs). Using data for a variety of coastal watersheds throughout the world, Boyer et al. (in prep.) found that riverine export was approximately 25% of NTNI. Aggregation of Nr input data for each region by Boyer et al. was based on the same data sources discussed in this paper (Appendix II).

Using this approach, we estimate that in 1860 riverine export of total Nr to inland receiving waters was 7.9 Tg N yr⁻¹, and that riverine export of total Nr in waters draining to the coastal ocean was ~27.0 Tg N yr⁻¹. The remainder of the Nr inputs not accounted for in riverine export were stored, emitted to the atmosphere as N₂ or transported to the oceans via the atmosphere.

Denitrification and Storage

About 30% of the 141 Tg N of terrestrial Nr created in 1860 was lost from continents: ~8 Tg N of N₂O are emitted to the atmosphere where it is stored or transported to stratosphere; 28.4 Tg N are transferred to coastal systems by rivers (27 Tg N) and atmospheric deposition (~1.4 Tg N), where most is denitrified (see below); and 7.1 Tg N are deposited to the open ocean surface (Figure 1a). The remaining ~98 Tg N is either stored within terrestrial systems as a reactive species or is denitrified to N₂, the primary process that converts Nr back to its unreactive form. Note that a small portion of

the denitrified N_r will not become N_2 , but rather chemically (NO) or radiatively (N_2O) important gases. The relative importance of storage vs. denitrification production of N_2 is arguably the largest uncertainty that exists for nitrogen budgets at most any scale. For pristine ecosystems over long time scales it is reasonable to propose that N_r creation should be balanced by denitrification production of N_2 . This hypothesis is supported by the constancy of the atmospheric N_2O recorded in ice bubbles back to 1000 AD, which implies that BNF and denitrification in pre-anthropogenic times were approximately equal (Ayres et al. 1994), and thus N_r was not accumulating in environmental reservoirs prior to human intervention. While the assumption of equality between BNF and denitrification may be reasonable over the very long time scale, there is certainly significant variability year to year. Thus we choose not to make a specific estimate for 1860, except to say that the upper limit of denitrification is 98 Tg N yr^{-1} , or 70% of total terrestrial N_r inputs. This implies that no net N_r storage occurred in terrestrial systems. A more detailed estimate of denitrification from soils, and the stream-river-estuary-shelf continuum is presented below for the present world.

Summary

In 1860 natural terrestrial BNF created $\sim 120 \text{ Tg N}$ of N_r and lightning created another 5.4 Tg N . Taken together, these natural processes were ~ 8 -fold greater than the N_r created by anthropogenic activities (fossil fuel combustion, 0.3 Tg N , and cultivation-induced BNF, 15 Tg N). After accounting for the transfer to the ocean via atmosphere and rivers, and the emission of N_2O to the atmosphere, $\sim 98 \text{ Tg N yr}^{-1}$ of the 141 Tg N yr^{-1} of new N_r created is ‘missing’ and was either stored as N_r in terrestrial systems or denitrified to N_2 .

Early-1990s

Fixation

BNF by natural terrestrial ecosystems in the early-1990s was $\sim 107 \text{ Tg N yr}^{-1}$. Lightning provided a minor additional source of 5.4 Tg N yr^{-1} . Anthropogenic activities created an additional $\sim 156 \text{ Tg N yr}^{-1}$ from food and energy production (Table 1).

Emission and Deposition

NO_x emissions totaled $\sim 45.9 \text{ Tg N yr}^{-1}$ in the early-1990s. Energy production accounted for $\sim 27.2 \text{ Tg N yr}^{-1}$ (from fossil fuel combustion, $20.4 \text{ Tg N yr}^{-1}$; biofuel combustion, 1.3 Tg N yr^{-1} ; non-road transport (e.g., ships), 3.6 Tg N yr^{-1} ; industrial emissions, 1.5 Tg N yr^{-1} ; 0.5 Tg N yr^{-1} by aircraft; and miscellaneous processes, 0.7 Tg N yr^{-1}) (Table 2a). Food production resulted in a total of 9.0 Tg N yr^{-1} of NO_x emissions, composed of combustion of agricultural waste (2.4 Tg N yr^{-1}), forests (1.1 Tg N yr^{-1}), and savanna grass (2.9 Tg N yr^{-1}) and emissions from fertilized soils (2.6 Tg N yr^{-1}). This final value was estimated by subtracting the 1860 soil NO_x soil emission value from the 1993 soil emission value under the assumption that fertilizer induced NO_x emissions were trivial in 1860. Other emissions were 0.6 Tg N yr^{-1} coming from the stratosphere, and 0.8 Tg N yr^{-1} from temperate wildfires (including human set fires). Total NO_x emissions were $45.9 \text{ Tg N yr}^{-1}$ of which $\sim 9.7 \text{ Tg N yr}^{-1}$ were from natural sources (Table 2a). Total NO_y deposition is $\sim 45.8 \text{ Tg N yr}^{-1}$, with $24.8 \text{ Tg N yr}^{-1}$ to continents and $21.0 \text{ Tg N yr}^{-1}$ to oceans (Table 1). It should be noted that there are other assessments of NO_x emissions

from soils that are larger than that used in this paper for the early 1990s (5.5 Tg N yr⁻¹). Davidson and Kingler (1997) estimate that global NO_x emissions range from 13 Tg N yr⁻¹ to 21 Tg N yr⁻¹. The differences in these estimates provide an indication of the uncertainty about our knowledge.

Anthropogenically induced NH₃ emissions in the early-1990s totaled 47.2 Tg N yr⁻¹. This value is higher from that of Bouwman et al. (2002) who estimated anthropogenic NH₃ emissions in 1990 to be 43 Tg N yr⁻¹, and lower than that of Schlesinger and Harley (1992), who estimated ~52 Tg N yr⁻¹. In our study, energy production resulted in 2.6 Tg N yr⁻¹ from the combustion of biofuels, 0.1 Tg N yr⁻¹ from fossil fuels, and 0.2 Tg N yr⁻¹ from industrial sources for a total of 2.9 Tg N yr⁻¹ (Table 2a). Food production resulted in NH₃ emissions to the atmosphere of 44.3 Tg N yr⁻¹. The component parts were combustion of agricultural waste (1.4 Tg N yr⁻¹), forests (1.4 Tg N yr⁻¹), and savannas (1.8 Tg N yr⁻¹). Low temperature NH₃ emissions were domestic animal waste (22.9 Tg N yr⁻¹), sewage and landfills (3.1 Tg N yr⁻¹), fertilizer (9.7 Tg N yr⁻¹) and crops (4.0 Tg N yr⁻¹). Natural emissions included high latitude burning (0.8 Tg N yr⁻¹).

Table 2b: Global atmospheric emissions of N₂O, Tg N yr⁻¹

	1860	early 1990s	2050	Notes
Soils				
natural	6.6	6.6	6.6	1
anthropogenic	1.4	3.2	3.2 +/-	2
Rivers				
natural	0.05	0.05	0.05	3
anthropogenic	-	1.05	3.22	4
Estuaries				
natural	0.02	0.02	0.02	3
anthropogenic	-	0.2	0.9	4
Shelves				
natural	0.4	0.4	0.4	5
anthropogenic	-	0.2	0.32	6
Ocean (natural)	3.5	3.5	3.5	7
TOTAL	12	15.2	18.2 +/-	8

Notes

1. We have assumed that the Bouwman et al. (1995) estimate of N₂O emissions from natural soils for for 1990s was applicable to 1860 and 2050 conditions.
2. 1860, Kroeze et al (1999); early 1990s, Bouwman et al. (1995).
3. Seitzinger and Kroeze 1998; Kroeze and Seitzinger 1998; Seitzinger et al. 2000
4. Seitzinger and Kroeze 1998; Kroeze and Seitzinger 1998; Seitzinger et al. 2001
5. Seitzinger et al. (2000)
6. early-1990s, Seitzinger et al. (2000); 2050, Kroeze and Seitzinger (1998)
7. Our value of 3.5 was obtained by subtracting shelf N₂O emissions from estimate of Nevison et al. (1995) for 1990. Further we assumed that Nevison et al estimate for 1990 was applicable to 1860 and 2050 conditions.
8. The '+/-' reflects the uncertainty in anthropogenic emissions in 2050.

In addition, emissions from oceans and soils/vegetation amounted to 5.6 Tg N yr⁻¹ and 4.6 Tg N yr⁻¹. As described before, rapid re-deposition in near-by regions effectively removes these emissions. Total NH₃ emissions were 58.3 Tg N yr⁻¹ of which ~11 Tg N yr⁻¹ were from natural sources (Table 2a). NH_x deposition was 34.1 yr⁻¹ and 12.4 Tg N yr⁻¹, to the continents and oceans respectively. Including the rapid deposition of the natural NH₃ emissions, these numbers would have been 38.7 Tg N yr⁻¹ and 18 Tg N yr⁻¹ respectively (Table 1).

In the early-1990s, global atmospheric anthropogenic NH₃ and NO_x emissions were ~47.2 Tg N yr⁻¹ and ~36.2 Tg N yr⁻¹, respectively, which is over half of the Nr created by human action (Figure 1b). About two-thirds of the NO_x emissions were as a consequence of energy

production (primarily fossil fuel combustion); the remainder was due to food production. About 95% of NH_3 emissions were from food production, with about half being from animal waste. For total N, about 70% of N emissions to the atmosphere are a consequence of food production. Nr deposition rates increased over a large area. In 1860, Nr deposition $>750 \text{ mg N m}^{-2} \text{ yr}^{-1}$ occurred over a very small area of southern Asia (Figure 1a). In the early-1990s, large portions of North America, Europe and Asia had rates $>750 \text{ mg N m}^{-2} \text{ yr}^{-1}$, and significant regions received $>1,000 \text{ mg N m}^{-2} \text{ yr}^{-1}$ (Figure 2b).

Global N_2O emissions increased from $\sim 12 \text{ Tg N yr}^{-1}$ in 1860 to $\sim 15 \text{ Tg N yr}^{-1}$ in the early-1990's (Figure 1b; Table 2b). Emissions from terrestrial soils and rivers were $\sim 8 \text{ Tg N yr}^{-1}$ in 1860 and increased to $\sim 11 \text{ Tg N yr}^{-1}$ in 1990 due primarily to increases in N use in food production systems. Specific sources were agricultural soils, animal waste management systems, biomass burning, biofuel combustion, energy/transport sources, industrial processes and from rivers associated with increased N inputs from leaching (Bouwman et al., 1995). Over the same time period N_2O emissions from estuaries and shelves also increased from $\sim 0.4 \text{ Tg N yr}^{-1}$ to $\sim 0.8 \text{ Tg N yr}^{-1}$ mainly due to indirect agricultural effects (Seitzinger et al. 2000) (Table 2b, Figure 1b), but see cautions above.




Figure 1b: Components of the global nitrogen cycle for early 1990s. See legend of Figure 1a for details

Riverine Export

As described above and in Appendix II, riverine Nr fluxes to inland receiving waters and to coastal systems was determined using the NTNI model relating net total Nr inputs to net total Nr fluxes discharged in rivers. Accounting for about 25% of the 230 Tg N yr⁻¹ net total Nr inputs to the terrestrial area of the continents in 1990, $\sim 11 \text{ Tg N yr}^{-1}$ were transported in rivers to inland receiving waters and drylands not draining to coastal

areas, and 48 Tg N yr⁻¹ are transported in rivers to coastal systems. Note that the *net total Nr inputs* to the global watersheds (230 Tg N yr⁻¹) used in the NTNI model calculation is less than the 268 Tg N yr⁻¹ of Nr *created* in the early-1990s as presented in Table 1 (Figure 1b). There are two reasons for this difference. First, the NTNI model considers net anthropogenic Nr input in atmospheric deposition from fossil fuel combustion, not total atmospheric NO_x emission from fossil fuel combustion. Second, the NTNI model considers net inputs of Nr in fertilizers used on the landscape. Not all the Nr created by the Haber-Bosch process is consumed as fertilizer—about 14% is used for industrial purposes. Further, the amount of N fertilizer consumed on an annual basis globally is about 6% less than the amount produced (FAOSTAT 2000).

Several other recent efforts have quantified global riverine Nr fluxes to the world's oceans for the 1990 timeframe. A study by Seitzinger and Kroeze (1998) predicted global riverine Nr exports of 21 Tg Nr yr⁻¹, though this estimate included only the dissolved inorganic fraction of Nr export to the coastal zone. Our approach included all fractions of Nr export (total Nr: dissolved, particulate, and organic forms) and suggests a global riverine export of 48 Tg Nr yr⁻¹. A lower global riverine loading estimate of 35 Tg Nr yr⁻¹ was predicted by Green et al. (2004), using similar data-sets to those described herein and an empirical model relating watershed characteristics to Nr export. In contrast, a higher global riverine loading estimate of 54 Tg Nr yr⁻¹ (total Nr) was predicted by Van Drecht et al (2001), based on point- and non-point sources of Nr and a model of their hydro-ecological transport and transformations. Such differences in these riverine Nr estimates highlight uncertainties stemming from data quality and resolution, scaling issues, and model approaches.

Nr Storage vs. Denitrification

In the early-1990s, human activities created ~156 Tg N yr⁻¹ of Nr. The Nr was extensively distributed by both anthropogenic (e.g., commodity transport, see below) and natural (e.g., atmospheric and hydrologic transport) processes. While we have a good understanding of the amount of Nr created, and a reasonable understanding of its dispersion, we have a poor understanding of its ultimate fate, especially about how much is denitrified to N₂. This question is critical to answer for it limits our ability to determine the Nr accumulation rate in environmental systems.

Denitrification generally requires an anoxic environment and a source of both organic matter and nitrate (Mosier et al. 2002). Thus, larger-scale hotspots for denitrification in terrestrial ecosystems tend to be characterized by high water contents (e.g., riparian zones, wetland rice, heavily irrigated regions, animal manure holding facilities, and other wet systems such as rain saturated soil). However, while average rates in well-drained upland systems are typically fairly low, a combination of large areas, periods of intense precipitation and the existence of anaerobic microsites even in well-oxygenated soils can in theory add up to a significant potential for N₂ loss at the landscape scale. When these large, well-aerated areas are combined with smaller hotspots, significant amounts of anthropogenic Nr may be denitrified (e.g., Yavitt and Fahey (1993)). Furthermore, the location and timing of denitrification in the environment may be more extensive than previously thought based on recent information on the potential for aerobic denitrification and other alternative denitrification pathways

(Robertson et al. 1995; see also review by Zehr and Ward 2002). The significance of these alternative denitrification pathways in the environment is uncertain.

For 1860, since we assumed that no Nr was stored in terrestrial systems, any Nr that was not lost from continents via net atmospheric losses or rivers was converted to N₂. For the early-1990s we present an assessment of the potential for denitrification by ecosystem type, and then at the end of the section we rely on a limited number of landscape-scale studies to estimate N₂ production relative to Nr inputs.

Atmosphere

In the early-1990s, the troposphere received ~46 Tg N yr⁻¹ of NO_x, ~58 Tg N yr⁻¹ of NH₃, and ~15 Tg N yr⁻¹ of N₂O. All of the emitted NO_x and NH₃ were deposited to the earth's surface, thus there was no accumulation (or conversion back to N₂ in the atmosphere). However, ~25% of the N₂O remained in the troposphere, the remainder was destroyed in the stratosphere (Dentener and Raes, 2002). Thus of the ~156 Tg N yr⁻¹ created by human action in the early-1990s, ~2.5% can be accounted for by tropospheric accumulation of N₂O.

Forests and grasslands:

Temperate ecosystems historically poor in N should in theory have a substantial capacity to store excess N, as suggested by several multiple-site studies. For example, NITREX (Dise and Wright 1992) and EXMAN (Rasmussen et al. 1990), established in Europe, manipulated the amount of Nr deposition to entire watersheds or large forest stands (Wright and Rasmussen 1998). In North America an informal collection of studies done at scales of plot to small catchments has used ¹⁵N fertilization additions to address questions concerning the fate of Nr deposited into forests (Nadelhoffer 2001; Seely and Lajtha 1997; Nadelhoffer et al. 1992, 1995, 1999; Preston and Mead 1994). Findings from these studies support the fact that most deposited N is retained within the watershed when the inputs are fairly small, primarily in the soil. Schlesinger and Andrews (2000) reviewed several studies that assessed the fate of N added to natural ecosystems. They also found that more N accumulates in the soil than in the woody biomass. It should be noted however, that with time some of the soil nitrogen can be later mineralized and made available to trees (Goodale et al. 2002). However, upland storage in biomass and soils can only be a transient phenomenon; at some point such sinks will begin to saturate, and losses to the atmosphere and aquatic systems will rise. The timing of a transition from substantial N storage to measurable saturation of biomass and soil sinks in a system experiencing chronic elevated N loading will vary with climate, vegetation type and pre-disturbance nutrient levels, but is unlikely to exceed several decades in most ecosystems (Dise and Wright 1995; Aber et al. 1998; Fenn et al. 1998).

Due to the interest in temperate forests as a potential sink for atmospheric CO₂, much of our recent information about the potential fate of excess Nr in natural ecosystems comes from northern mid-latitude forests (Nadelhoffer 2001; Seely and Lajtha 1997; Dise and Wright 1992). Many studies suggest that total rates of denitrification tend to be relatively low in temperate forest soils because most are well-drained with limited anoxic regions (Gundersen 1991; Nadelhoffer 2001). However, in forest systems with high rates of N deposition, not only are leaching rates increased but also gaseous losses of N are enhanced (Aber et al. 1995). For example, research at

Hogwald Forest, Germany on spruce and beech plots suggests that over a 4-year period that N_2 fluxes can be a significant fraction of the N deposited (Butterbach-Bahl et al. 2002).

Denitrification rates are typically larger in moist tropical forests, where both moisture and nitrate levels can be substantially higher, thus increasing N additions to tropical regions may produce more rapid and sizable denitrification losses (Matson et al. 1999). The potential for denitrification is quite large in the riparian zone of forested ecosystems where there is a supply of organic matter and a source of nitrate (Steinhart et al. 2000). Overall, as stated above, our knowledge of total potential denitrification at the scale of large forested watersheds is far too incomplete, and further research in this area is sorely needed.

Finally, many of the most dramatic recent increases in atmospheric deposition to natural ecosystems have not been in temperate forests, but rather in grassland, high elevation, and semi-arid regions (Williams et al. 1996; Asner et al. 2001). Total denitrification rates in such systems are likely to be quite low due to a combination of largely drier conditions and well-drained soils. However, our knowledge about how excess N may be partitioned in such systems, most notably semi-arid zones, is far worse than it is for temperate forests, and the primary controls over the fate of N in such regions are likely to be quite different than seen in forested systems (Asner et al. 2001). There can be bursts of denitrification under certain conditions. For example, Peterjohn and Schlesinger (1991) found that there are high potential N_2O losses from Chihuahuan desert soils (USA), after periods of rainfall.

Agroecosystems

Agroecosystems receive ~75% of the N_r created by human action. As the amount stored within the systems each year is small relative to the amount that enters, most N_r is dispersed to other systems. Smil (1999) estimates that ~170 Tg $N \text{ yr}^{-1}$ were added to the world's agroecosystems in the early-1990s, ~120 Tg $N \text{ yr}^{-1}$ from the addition of new N_r (fertilizer and cultivation-induced BNF), and 50 Tg $N \text{ yr}^{-1}$ from the addition of existing N_r (e.g., atmospheric deposition, crop residues, animal manure). He concludes that very little of the N accumulates in the agroecosystem soil (on the order of 2% - 5%), and that most is either removed in the crop (~50%), emitted to the atmosphere (~25%) or discharged to aquatic systems (~20%). This low rate of accumulation is supported by van Breemen et al. (2002), who estimate for 16 large watersheds in eastern USA that only ~10% of the inputs are stored within the agroecosystem. In fact, agroecosystems in some regions of the world are losing N_r . Li et al. (2003), using the DNDC model, report that in 1990 Chinese agroecosystems were losing organic C at 1.6% yr^{-1} . Losses of soil organic carbon are commonly observed in agroecosystems (Davidson and Ackerman 1993) and usually are accompanied by soil organic N losses because of the relative narrow C:N ratio associated with soil organic matter (8-15).

Denitrification production of N_2 can be an important loss of N_r from agroecosystems (Mosier et al. 2002), however the rates are quite variable. In a collection of papers, Freney and his colleagues find a wide range of denitrification rates. For 15 studies of flooded rice, denitrification ranged from 3% to 56% of the N_r applied, with a median value of 34% (Simpson et al. 1984; Galbally et al. 1987; Keerthisinghe et al. 1993; Simpson and Freney 1988; Freney et al. 1995; Cai et al. 1986; Zhu et al. 1989; De

Datta et al. 1989; Freney et al. 1990). An investigation of irrigated wheat found 50% of the applied Nr denitrified (Freney et al. 1992). Two investigations of irrigated cotton reported on five studies that ranged from 43% to 73% of the Nr applied was denitrified (median was 50%) (Freney et al. 1993; Humphreys et al. 1990). For 3 studies of dryland wheat (Bacon and Freney 1989) the values ranged from 2 to 14% of the Nr applied, and a median value of 11%. As expected, irrigated crops had higher rates. However, the denitrification rates from some irrigated crops may be lower. Mosier et al. (1986) in a study of irrigated corn and barley reported that the total volatile loss of Nr from N₂O and N₂ was 1-2.5% of the applied fertilizer. These findings correspond with the findings of Rolston et al. (1978, 1982) and Craswell and Martin (1975a,b) who found a loss of 3-4% of the applied fertilizer N.

In the comparative assessment of N dynamics, Li et al. (2003) found that denitrification fluxes of NO, N₂O and N₂ ranged from 1.4 - 5.0 Tg N yr⁻¹ in China and 2.2 - 5.9 Tg N yr⁻¹ in the USA. Production of N₂ was 0.6 - 2.6 Tg N yr⁻¹ for China and 1.0 - 3.2 Tg N yr⁻¹ for the USA, with the range set by whether the soil had high or low levels of soil organic carbon. Relative to the external inputs (fertilizer, atmospheric deposition, cultivation induced BNF), on average <5% of the Nr was converted to N₂ in agroecosystems.

In the Netherlands, with very high levels of Nr production, 30-40% of the Nr applied to agroecosystems is either stored or denitrified (Erisman et al. 2001). Van Breemen et al. (2002) analyzed the fate of nitrogen introduced into 16 large watersheds in the eastern USA. After accounting for all other sinks, they estimate (by difference) that relative to Nr inputs to agricultural lands, denitrification within soils of the agroecosystems ranges from 34% to 63% with a weighted mean of 49%. The large differences in these estimates of denitrification relative to inputs reflect both regional variability in the conditions that promote denitrification (e.g., soil moisture) and the general uncertainty in estimating storage and loss of N in the terrestrial landscape.

On the largest scale, in an analysis of global agroecosystems, Smil (1999) reviews estimates of denitrification production of N₂ based on N₂O/N₂ ratios, and based on estimates of total denitrification. He concludes that annual N₂ fluxes range from 11-18 Tg N yr⁻¹ (mean, 14 Tg N yr⁻¹). Relative to the ~170 Tg N yr⁻¹ added to the agroecosystem (including atmospheric deposition, fertilizer and manure), ~8% of the applied Nr is denitrified to N₂.

In summary we suggest that >75% of the Nr applied to agroecosystems is removed as Nr (either via crop removal, or biogeochemically into the atmosphere or water). The amount stored in the agroecosystem is small (<10%) and, while there is substantial variability, the amount denitrified to N₂ appears to be on the order of 10-40% of inputs.

Groundwater

Human activities, particularly food production and use of septic systems, have resulted in increases in nitrate levels in groundwater, which are of growing concern in many regions of the world. In the United States, nitrate levels are higher than 10 mg N L⁻¹ (the standard for drinking water recommended by the World Health Organization) in approximately 20% of wells in farmland areas, between 2 to 10 mg L⁻¹ in 35% of wells, and below 2 mg L⁻¹ in only 40% of wells. Nitrate levels in China and India are also often

high, and are generally correlated with the rate of use of nitrogen fertilizer (Agrawal et al. 1999; Zhang et al. 1996). In Europe high levels of nitrate are also associated with agriculture (Howarth et al. 1996).

High levels of nitrate in groundwater are of concern because they can pose a significant health risk (Follet and Follet 2001; Townsend et al 2003). However, it does not appear that accumulation of nitrogen in groundwater is a major sink for the nitrogen mobilized in the landscape by human activity. For Europe and North America, the average rate of increase of nitrogen in the groundwater in areas of intense agricultural activity is in the range of 25 to 40 mg N m⁻² yr⁻¹, which accounts for at most a few percent of the nitrogen inputs to the landscape from human activity (Howarth et al. 1996). Where intensive agriculture occurs on areas of highly permeable sandy soils, rates of nitrogen accumulation in the groundwater can be as high as 200 mg N m⁻² yr⁻¹ (Howarth et al. 1996). For comparison, the average rate of fertilizer application on all agricultural lands in the United States as of 1997 was 6 x 10³ mg N m⁻² yr⁻¹ (Howarth et al. 2002) and for intensively farmed corn culture in the American mid-west was approximately 1 x 10⁴ mg N m⁻² yr⁻¹ (Boesch et al. 2002).

Surface waters

The wetland, stream, river, estuary, shelf region provides a continuum with substantial capacity for denitrification. Nitrate is commonly found, there is abundant organic matter, and sediments and suspended particulate microsites offer anoxic environments. In this section we discuss denitrification in the stream/river/estuary/shelf continuum. While there are several specific studies of denitrification in wetlands, there is a need for a better understanding of the role of wetlands in Nr removal at the watershed scale.

Seitzinger et al. (2002) estimate that of the Nr that enters the stream/river systems that drain 16 large watersheds in eastern USA, 30 to 70% can be removed within the stream/river network, primarily by denitrification. In an independent analysis on the same watersheds, Van Breemen et al. (2002) estimated by difference that the lower end of the Seitzinger et al. (2002) range is most likely. In a global analysis, Green et al. (2004) showed that on a basin-wide scale that there was an average loss/sequestration of 18% (range 0-100%) of the Nr through the combined effects of soil, lake/reservoir, wetland and riverine systems, based on residency time and temperature differences across watersheds. For Nr that enters estuaries, 10% to 80% can be denitrified, depending primarily on the residence time and depth of water in the estuary (Seitzinger 1988; Nixon et al. 1996).

Of the Nr that enters the continental shelf environment of the North Atlantic Ocean from continents, Seitzinger and Giblin (1996) estimated that >80% was denitrified. The extent of the denitrification is dependent in part on the size of the continental shelf. Regions where the shelf denitrification potential is large are the south and east coast of Asia, the east coast of South America and North America. It is interesting that these are also the regions where riverine Nr inputs are the largest. While most of the riverine/estuary Nr that enters shelf region is denitrified, total denitrification in the shelf region is larger than that supplied from the continent and thus Nr advection from the open ocean is required.

In summary, based on the ranges above, we estimate that ~50% (range 30% to 70%) of the N that enters the stream-river continuum is denitrified, and that of the N_r remaining another 50% (range 10% to 70%) is denitrified in the estuary, leaving 25% of the original N_r that entered the stream; most of which is denitrified in the shelf region. It should be stressed that there is substantial uncertainty about these 'average' values.

Landscape-scale estimates

The previous material focused on denitrification in specific systems. What about denitrification on the landscape scale? Several recent studies have estimated denitrification in land and associated freshwaters relative to inputs for large regions. At the scale of continents, denitrification has been estimated as 40% of N_r inputs in Europe (van Egmond et al. 2002) and ~30% in Asia (Zheng et al. 2002). At the scale of large regions, estimates of denitrification as a percentage of N_r inputs include 33% for land areas draining to the North Atlantic Ocean (Howarth et al. 1996) and 37% for land areas draining to the Yellow-Bohai Seas (Bashkin et al. 2002). Country-scale estimates of the percentage of N_r inputs that are denitrified to N_2 in soils and waters include ~40% for the Netherlands (Kroeze et al. in press), 23% for the USA (Howarth et al. 2002), 15% for China (Xing and Zhu 2002), and 16% for the Republic of Korea (Bashkin et al. 2002).

On a watershed scale, Van Breemen et al. (2002) estimate that 47% of total N_r inputs to the collective area of 16 large watersheds in the northeastern U.S. are converted to N_2 : 35% in soils and 12% in rivers. Within the Mississippi River watershed, Burkart and James (2002) divided the basin into 6 large sub-basins, concluding that soil denitrification losses of N_2 ranged from a maximum of about 10% of total inputs in the Upper Mississippi Region to less than 2% in the Tennessee and Arkansas/Red Regions. Goolsby et al. (1999) estimate denitrification within soils of the entire Mississippi-Atchafalaya watershed to be ~8% but did not quantify additional denitrification losses of N_r inputs in river systems. Relative to inputs, landscape-scale estimates of N_r denitrified to N_2 in terrestrial systems and associated freshwaters are quite variable, reflecting major differences in the amount of N inputs available to be denitrified and in environmental conditions that promote this process. All of the cited studies reporting denitrification at regional scales claim a high degree of uncertainty in the estimates, highlighting the fact that our knowledge of such landscape-level rates of denitrification is quite poor. The few estimates that do exist are subject to enormous uncertainties, and must often be derived as the residual after all other terms in a regional N budget are estimated, terms which themselves are often difficult to constrain (e.g. Erisman et al. 2001; Van Breemen et al. 2002). Notwithstanding these uncertainties, for the purposes of this paper, for continental systems we estimate that 25% (within a range of 10% to 40%) of the N_r applied is denitrified to N_2 .

Global-scale denitrification estimates

The previous sections have presented a brief review of denitrification by ecosystem type and at the landscape scale. This section presents our estimates of global denitrification for continental regions of the world. In the early-1990s, 268 Tg N yr^{-1} of new N_r was added to the earth's surface. Of this amount, ~93 Tg N yr^{-1} (~34%) was lost from the continents, via river discharge to the coasts (48 Tg N yr^{-1}), N_2O emission to the atmosphere (11.3 Tg N yr^{-1}) and NO_x and NH_3 emitted to the continental atmosphere and

deposited to the ocean surface ($33.4 \text{ Tg N yr}^{-1}$). Thus 175 Tg N yr^{-1} is unaccounted for. Some is stored in soils and biota, the rest is denitrified in either terrestrial systems or in the stream-river continuum prior to discharge to the coast. Our estimate of the former relies on the landscape scale studies where we concluded that 25% of inputs are denitrified in the soil, or 67 Tg N yr^{-1} . Our estimate of the latter is that in the stream-river continuum, denitrification is equal to discharge to the coastal zone, or 48 Tg N yr^{-1} . Thus of the 175 Tg N yr^{-1} remaining, 115 Tg N yr^{-1} is denitrified and 60 Tg N yr^{-1} is stored in terrestrial systems. Concerning denitrification in the estuary and shelf regions, we assume that of the Nr that enters the estuary via rivers that 50% is denitrified or 24 Tg N yr^{-1} . We further assume that all of the continental Nr that enters flows from the estuaries to the shelf is denitrified, together with additional Nr advected to the shelf from the open ocean (see below).

The uncertainties about these estimates are large enough that the relative importance of denitrification vs. storage is unknown. But the calculations do suggest that denitrification is important in all portions of the stream/river/estuary/shelf continuum and that on a global basis the continuum is a permanent sink for Nr created by human action.

Summary

Between 1860 and the early-1990s, the amount of Nr created by natural terrestrial processes decreased by ~15% (120 Tg N yr^{-1} to $\sim 107 \text{ Tg N yr}^{-1}$) while Nr creation by anthropogenic processes increased by ~10-fold ($\sim 15 \text{ Tg N yr}^{-1}$ to $\sim 156 \text{ Tg N yr}^{-1}$). Human creation of Nr went from being of minor importance, to becoming the dominant force in the transformation of N_2 to Nr on continents.

Much of the Nr created in the early-1990s was dispersed to the environment. Of the $\sim 268 \text{ Tg N yr}^{-1}$ created by natural terrestrial and anthropogenic processes, $\sim 98 \text{ Tg N yr}^{-1}$ of NO_x and NH_3 was emitted to the atmosphere. Of that amount, $\sim 65 \text{ Tg N yr}^{-1}$ was deposited back to continents and $\sim 33 \text{ Tg N yr}^{-1}$ was deposited either to the estuary and shelf region ($\sim 8 \text{ Tg N yr}^{-1}$) or to the open ocean ($\sim 25 \text{ Tg N yr}^{-1}$). An additional $\sim 59 \text{ Tg N yr}^{-1}$ was injected into inland (11 Tg N yr^{-1}) and coastal (48 Tg N yr^{-1}) systems via rivers. Thus losses of Nr from continents to the marine environment total $\sim 81 \text{ Tg N yr}^{-1}$ from atmospheric and riverine transport. Rivers are more important than is atmospheric deposition in delivering Nr to coastal/shelf regions ($\sim 48 \text{ Tg N yr}^{-1}$ vs. $\sim 8 \text{ Tg N yr}^{-1}$, respectively). Conversely, since most of the Nr introduced to coastal systems is converted to N_2 along the continental margins, the atmosphere is more important than rivers in delivering Nr to the open ocean.

A small ($\sim 11 \text{ Tg N yr}^{-1}$), yet environmentally important, amount of the Nr created is emitted to the atmosphere as N_2O from continents, estuaries and the shelf region, where a portion ($\sim 25\%$) accumulates in the troposphere until eventual destruction in the stratosphere.

Thus of the $\sim 268 \text{ Tg N yr}^{-1}$ of new Nr that entered continents, $\sim 81 \text{ Tg N yr}^{-1}$ was transferred to the marine environment via atmospheric and riverine transport, and $\sim 12 \text{ Tg N yr}^{-1}$ was emitted to the atmosphere as N_2O . Of the remaining 175 Tg N yr^{-1} of Nr we estimate that $\sim 115 \text{ Tg N yr}^{-1}$ is converted to N_2 and that $\sim 60 \text{ Tg N yr}^{-1}$ is accumulating in terrestrial systems. As discussed earlier, there is large uncertainty about the storage and N_2 production values. Since it is unlikely that either one is near zero and more likely that the ranges overlap, it seems clear that significant fractions of the missing N are routed to

both denitrification and storage in biomass and soils. We view improved resolution of the partitioning of anthropogenic Nr as a critical research priority. The purpose of the above analysis is to try to track the fate of the Nr introduced to environmental reservoirs. It is important to note that the analysis does not include an assessment of landuse change on the mobilization of N in soil and vegetation pools.

Regional Nitrogen Budgets

Introduction

To gain insight into the spatial heterogeneity of Nr creation and distribution, we examine N budgets by geopolitical region for the early-1990s. The geographical units in the regional analysis are Asia, Africa, Europe (including the former Soviet Union, FSU), Latin America, North America and Oceania. These units are collections of countries as defined by the FAO (2000). This analysis on the regional scale is important as it illustrates the differences in Nr creation and distribution as a function of level of development and geographic location. In addition, the short atmospheric lifetime of NO_y, and NH_x (1 day to 1 week depending on total burden and altitude) means that the concentration of these chemical species varies substantially in both space and time. Atmospheric transport and dynamics, N emissions, chemical processing, removal mechanism (dry vs wet deposition) all interact to alter the spatial distribution of reactive N. Dividing the Earth's land surface by geopolitical region offers a convenient mechanism for examining this spatial variation.

Table 3: Nr creation by region, Tg N yr⁻¹

	Nr Inputs (Note 1)						Sum Inputs
	Lighthing	BNF	Fossil fuel	Fertilizer Prod	C-BNF	Imports	
Africa	1.4	25.9	0.8	2.5	1.8	1.2	33.6
Asia	1.2	21.4	5.7	40.1	13.7	13.8	95.9
Europe/FSU	0.1	14.8	6.1	21.6	3.9	9.6	56.1
L. America	1.4	26.5	1.3	3.2	5.0	2.5	39.9
N. America	0.2	11.9	7.3	18.3	6.0	5.0	48.7
Oceania	0.2	6.5	0.4	0.4	1.1	0.6	9.2
TOTAL	4.4	107	21.5	86.1	31.5	32.7	283

	Nr Outputs (Note 2)						Sum Outputs (not riv-inland)	% Input/Output
	Exports	Atmos-NO _y	Atmos-NH _x	Riverine-Inland	Riverine-Coastal	Total Riverine		
Africa	1.1	1.9	1.4	2.0	6.6	8.5	11.0	33%
Asia	5.1	1.9	1.4	5.1	16.7	21.8	25.1	26%
Europe/FSU	15.2	3.0	2.4	0.7	8.4	9.1	29.0	52%
L. America	2.7	2.3	2.0	1.4	8.2	9.7	15.2	38%
N. America	8.3	3.8	0.6	0.6	7.2	7.8	19.9	41%
Oceania	0.3	0.6	0.5	1.5	0.7	2.1	2.1	23%
TOTAL	32.7	13.5	8.3	11.3	47.8	59.0	102	36%

Notes

1. Nr Inputs *

- 'Lightning'--Lelieveld and Dentener (2000)
- 'BNF'--based on Cleveland et al (1999) as discussed in the text.
- 'Fossil Fuel'--Klein Goldewijk and Batjes (1997); van Aardenne et al. (2001)
- 'Fertilizer'--Bouwman 1995 (via FJ Dentener)
- 'C-BNF cultivation'--Table 4
- 'Imports'--Table 5

2. Nr Outputs *

- 'Exports'--Table 5
- 'Atmos-NO_y'--by difference between NO_x emission and NO_y deposition (Table 6)
- 'Atmos-NH_x'--by difference between NH₃ emission and NH_x deposition (Table 6)
- 'Riverine'--from Boyer et al. (in prep), see text and Appendix A. Note that 'Riverine-Inland' output is not included in "Sum Outputs".

* Values in bold are the largest flux for each country within the 'input' or 'output' subclass (e.g., the largest Nr input for Africa is BNF).

Regional N fluxes

Natural Nr creation

Global production of NO_x by lightning is 5.4 Tg N yr^{-1} with most of it occurring over continents with significant tropical regions; Africa, Asia and Latin America have 1.4, 1.2, and 1.4 Tg N yr^{-1} , respectively, of lightning fixation. Europe and North America account for only 0.1 and 0.2 Tg N yr^{-1} , respectively (Table 3, Figure 3). Marine lightning fixed 1.1 Tg N yr^{-1} . On a global basis, natural BNF created 107 Tg N yr^{-1} in the early-1990s (Table 1). The regional breakdown has most ($\sim 70\%$) of the BNF occurring in regions with warmer climates—Africa ($25.9 \text{ Tg N yr}^{-1}$), Latin America ($26.5 \text{ Tg N yr}^{-1}$) and Asia ($21.4 \text{ Tg N yr}^{-1}$). Other regions create lesser amounts of Nr by BNF—North America ($11.9 \text{ Tg N yr}^{-1}$), Europe/FSU ($14.8 \text{ Tg N yr}^{-1}$) and Oceania (6.5 Tg N yr^{-1}) (Table 3). In support of these rough estimates, Zheng et al. (2002) estimate that natural BNF in Asia created $15.8 \text{ Tg N yr}^{-1}$ in 2000, similar to our estimate of $21.4 \text{ Tg N yr}^{-1}$ (Table 3).

Anthropogenic Nr creation

Asia dominates Nr creation as a consequence of food production. Fertilizer N production is almost double that of the next closest region (Europe/FSU) (Table 3). Cultivation-induced BNF is almost twice the next closest region (N. America) with




Figure 3: Nr creation and distribution by geopolitical region for the mid-1990s, Tg N yr⁻¹

legumes, forages and rice production all playing major roles. Asia is followed by Europe/FSU and North America (Tables 3 and 4; Figure 3). For Nr creation by energy production, North America leads the way followed closely by Europe/FSU and Asia. Overall, the creation of Nr in Asia approximates that in Europe/FSU and North America combined (Table 3). Globally, fossil fuel creation of total Nr (in form of NO_x) is approximately 16% (note that only fertilizer production is considered in the regional analysis and not the other industrial uses of NH_3 created by the Haber-Bosch process (Febra Domene and Ayres, 2001)). The larger terms of fertilizer production and cultivation are also more uncertain because of difficulties in establishing reliable inventories in regions like Asia (Smil 1995).

Table 4: Cultivation-induced Nr creation by region, TgN/yr¹ (1995)

	Legumes	Forage+Misc	Rice	Sugar Cane	Total
Africa	1.5	0	0.2	0.1	1.8
Asia	4.1	4.5	4.3	0.8	13.7
Europe+FSU	0.4	3.5	0	0	3.9
L. Amer.	1.9	2	0.2	0.9	5
N. Amer.	2.5	3.5	0	0	6
Oceania	0.2	0.8	0	0.1	1.1
Total	10.6	14.3	4.7	1.9	31.5

Nr commodity exchange

Once produced, Nr containing commodities (primarily fertilizer and grain) are exchanged among regions. Asia was the largest net importer ($\sim 8.7 \text{ Tg N yr}^{-1}$). Europe/FSU and North America were the largest net exporters ($\sim 5.6 \text{ Tg N yr}^{-1}$ and $\sim 3.3 \text{ Tg N yr}^{-1}$, respectively). All other regions had net exchanges that were $< \sim 0.5 \text{ Tg N yr}^{-1}$ (Table 5, Figure 3) (Galloway and Cowling 2002).

Table 5: Exchange of N (import and export) in commodities among regions, Tg N/yr¹ (1995)

	Import Fertilizer	Import Plant	Import Meat	Export Fertilizer	Export Plant	Export Meat	Net Fertilizer	Net Plant	Net Meat	Import Total	Export Total	Net Total
Africa	0.7	0.5	0.0	1.0	0.0	0.0	-0.3	0.5	0.0	1.2	1.1	0.2
Asia	10.7	2.9	0.1	4.3	0.8	0.1	6.4	2.2	0.1	13.8	5.1	8.7
Europe+FSU	6.6	2.7	0.2	13.2	1.7	0.2	-6.6	1.0	0.0	9.6	15.2	-5.6
L. Amer.	1.9	0.7	0.0	1.2	1.5	0.0	0.7	-0.9	0.0	2.5	2.7	-0.2
N. Amer.	4.8	0.2	0.0	5.2	3.1	0.1	-0.4	-2.9	-0.1	5.0	8.3	-3.3
Oceania	0.6	0.0	0.0	0.0	0.3	0.1	0.6	-0.2	-0.1	0.6	0.3	0.3
Total	25.2	7.1	0.5	24.9	7.3	0.5	0.4	-0.3	0.0	32.8	32.7	0.1

Riverine discharge

In the early-1990s $\sim 59 \text{ Tg N yr}^{-1}$ was discharged via riverine export, with $\sim 11 \text{ Tg N yr}^{-1}$ transported to inland receiving waters and drylands and $\sim 48 \text{ Tg N yr}^{-1}$ transported to coastal waters. Asia had the most Nr transported to inland receiving waters/drylands (5.1 Tg N yr^{-1}) and N. America had the least. Interestingly, about twice as much Nr was transported to inland receiving waters/drylands of Oceania than was transported to the

coast. The comparison of Nr riverine transport to coast shows Asia with the largest transport at $\sim 16.7 \text{ Tg N yr}^{-1}$, and all other regions except for Oceania transporting in the range of $6 - 9 \text{ Tg N yr}^{-1}$ (Table 3; Figure 3).

Atmospheric emission, deposition and export to marine atmosphere

The rates of N emission, deposition and export to the marine atmosphere (emission minus deposition) are consistent with the patterns of Nr creation (Tables 3 and 6). Asia has the greatest rates of emission, deposition and export of both oxidized N (NO , NO_2 and NO_y) and reduced N (NH_3 and NH_x). Asian export of NO_y is 4.2 Tg N yr^{-1} , less than the 6.0 Tg N yr^{-1} of reduced N exported (Table 6). For comparison, total export of both oxidized and reduced N from North America is 4.4 Tg N yr^{-1} , and 5.4 Tg N yr^{-1} from Europe/FSU. NH_3 emissions are 4.4 Tg N yr^{-1} higher in Europe/FSU than in North America and NO_x emissions are similar for the two regions. These modeling results are consistent with estimated N deposition budgets based on measurements and models done for the US and Western Europe (Holland et al., pers. comm.; Holland et al. 1999; Whelpdale et al. 1996, 1997).

In these model simulations, roughly twice as much NO_y falls on the continents relative to the oceans (Table 6, Figure 1b). A greater proportion of emitted NH_3 falls on land than oceans with a total of $36.6 \text{ Tg NH}_x\text{-N yr}^{-1}$ out of a total of $\sim 58.2 \text{ Tg N yr}^{-1}$.

Table 6: Regional atmospheric emissions and deposition for the early-1990s, Tg N yr⁻¹

	Emission NOx	Deposition NOy	Export/Import NOy	Emission NH3	Deposition NHx	Export/Import NHx
Continents						
Africa	6.8	5.0	-1.9	7.0	5.6	-1.4
Asia	10.7	6.5	-4.2	22.1	16.1	-6.0
L. America	5.3	3.1	-2.2	7.8	5.8	-2.1
N. America	8.5	4.7	-3.8	3.6	3.0	-0.6
Oceania	1.1	0.5	-0.6	1.0	0.5	-0.5
Europe/FSU	7.9	5.0	-2.9	8.0	5.6	-2.4
subtotal	40.4	24.8	-15.6	49.5	36.5	-13.0
Oceans						
Arctic		0.4	0.4	0.3	0.4	0.1
S. Pacific		1.8	1.8	1.8	2.6	0.8
N. Pacific		6.3	6.3	1.6	5.8	4.2
N. Atlantic		8.1	8.1	0.6	3.8	3.2
S. Atlantic		1.2	1.2	0.7	1.7	1.0
Indian		3.2	3.2	1.1	4.1	3.0
subtotal	4.2	21.0	16.8	5.6	18.3	12.7
TOTAL	44.6	45.8		55.1	54.8	

* Emissions based upon van Aardenne et al., (2001). NOx emissions for Oceans are from lightning and ships, and are not broken down by ocean basin. See text for discussion of deposition. Exchanges are either imports (+) via the atmosphere to the region or exports (-) from the region and are calculated by the difference between emissions and deposition.

Global NH_3 emissions are $\sim 25\%$ greater than global NO_x emissions. The imbalance between global emissions and deposition suggests that N may be transported to the middle and upper atmosphere, out of the realm of this model focused on tropospheric chemistry, and/or the model has a problem with mass conservation. A comparison of five earlier generation 3D chemical transport models simulated 20% more NO_y deposition on

land for fossil fuel derived NO_y only (Holland et al. 1997). Four of the five models simulated as much as 2 times more NO_y deposition on land when the full NO_y budget was considered. One model GRANTOUR, lacked mechanisms for formation of long range transport species, i.e. PAN, and simulated greater NO_y deposition on oceans than on land. The contrasting model results underscore the gaps in our understanding and the advances that are needed in understanding and measuring global deposition patterns. Thorough evaluation of the models with available measurements is needed to resolve some of the differences and to point to important areas for more measurements.

Points of Discussion

Natural vs. anthropogenic N_r creation

Although anthropogenic inputs of N at the global scale may have roughly doubled the amount of N_r entering the terrestrial environment each year; a more detailed analysis of the spatial distribution of natural BNF suggests a much different story. In heavily industrialized or agricultural regions of the world, anthropogenic sources of nitrogen are much higher than natural background levels, while in more pristine systems, BNF may still contribute the bulk of fixed N inputs (Chameides et al. 1994; Howarth et al. 1996) and for Africa and Latin America, it is the largest N_r source (Table 3). For example, the highest reported rates of natural BNF are in tropical savanna, tropical evergreen rain forest, tropical floodplain, and wet savanna ecosystems (Cleveland et al. 1999). Together, these ecosystems supply more than 60% of the N fixed in natural terrestrial ecosystems each year. However, these non-agricultural areas are receiving relatively low inputs of anthropogenic N via atmospheric deposition (Figure 2b). In contrast, N inputs via BNF in temperate ecosystems are relatively low. Altogether, BNF in temperate grasslands, temperate forests, and boreal ecosystems contributes <15% of the total N




Figure 2b: Spatial patterns of total inorganic nitrogen deposition in early 1990s, $\text{mg N m}^{-2} \text{yr}^{-1}$.

fixed naturally per year (from Cleveland et al. (1999) analysis) while inputs of N_r via N deposition in the temperate zone may be relatively large (Figure 2b). These patterns suggest that while human activities have undoubtedly increased global N fixation, a "global average" represents a deceptive portrayal of the effect to which humans have altered global N fixation. In still relatively undisturbed areas of the world, N inputs via natural N fixation dominate, and external inputs of

newly fixed N may have changed little from the pristine ecosystem (Hedin et al. 1995). Alternatively, in the highly developed temperate zone, inputs of fixed N may actually be several times higher than in pre-industrial times (Cleveland et al. 1999).

Nr transfer to other regions

Nr is transferred out of regions by atmospheric transport, primarily to marine regions, by riverine transport to the coastal zone, and by commodity export (primarily grain and fertilizer) to other regions. The relative importance of the three processes differs by region. North America and Europe/FSU transfer most of their Nr by commodity export, with riverine transfers being the next most important. For the other regions, riverine transfers are the most important. In fact for Asia, Latin America and Africa, riverine transfers are greater than all the other transfers combined (Table 3).

Unknown losses of Nr

For the regions we consider, the three Nr transfers discussed above account for ~26% to ~52%, relative to inputs (Table 3). There is important regional variability. For Africa known outputs account for 33% of the Nr inputs. At the other extreme, for Europe/FSU they account for 52%. While the magnitude of the fate of the remaining Nr is unknown, we do know the processes. About 12 Tg N yr⁻¹ of the Nr is emitted to the atmosphere as N₂O. Our knowledge of spatial emissions patterns is not sufficient to estimate this on a regional basis. Most of the balance is either stored in the region in soil, biomass, etc., or it is denitrified to N₂ in the terrestrial system or the wetland-stream-river continuum. Earlier we assumed that on a global basis, as much Nr was denitrified in surface waters as was discharged to coastal systems. Given the probable strong regional variability of N₂ production vs. Nr storage, we do not extend that assumption here, but note that N₂ production is likely to occur in warmer regions with a large P/ET ratio—such as Africa, Latin America and Asia. In other regions, a greater portion of the Nr has the potential to accumulate.

Summary

Overall, Asia, Europe and North America account for nearly 90% of the current human increase in BNF and use, thus the majority of our current focus on problems associated with an accelerated N cycle should be on these regions. However, as will be seen below, substantial increases are projected over the next 50 years for other regions of the world. As for the already heavily changed continents, it is essential that we begin to improve our understanding of the potential fates for additional N in these future hotspots. This statement is especially true for tropical regions, where N often is not the limiting nutrient even in little disturbed ecosystems (Martinelli et al. 1999). This fact, combined with warm, often wet climates, can lead to high rates of N loss to atmospheric and aquatic realms, making it likely that even a modest rise in anthropogenic N inputs could lead to rapid increases in Nr losses to air and water (Matson et al. 1999).

Marine N Budgets

Introduction

In this section, we have focused on the Nr creation in the continental regions and its dispersion both within continents, and via atmospheric and riverine transport to the oceans. To put these losses from continents into context with the rest of the global system, in this section of the paper we analyze the N budget of the ocean by basin. The base time period is the early-1990s. With the exception of riverine N fluxes,

denitrification on the shelf and atmospheric deposition of N and Fe, the analysis is time-independent.

Nr Creation

N₂ Fixation in the Oceans by extrapolation of direct measurements

Most research efforts on marine BNF to date have focused on the conspicuous planktonic non-heterocystous cyanobacterium, *Trichodesmium* spp. (Capone et al. 1997, Capone and Carpenter 1999; Karl 2002). *Trichodesmium* has a cosmopolitan distribution in the world's oligotrophic oceans and is most common in waters of 20° C and warmer (Carpenter 1983 a,b) with substantial populations and blooms restricted to waters of 25° C or greater (Carpenter and Capone 1992).

We have therefore chosen to provide a minimum estimate for oceanic BNF by extrapolating the accumulated data on directly determined rates of N₂ fixation by *Trichodesmium* to warm oligotrophic surface waters. Studies considered generally coupled estimates of the population density of *Trichodesmium* through the upper water column with estimates of nitrogenase activity based on either the C₂H₂ reduction or ¹⁵N₂ uptake methods (Table 7).

An analysis of nine studies in tropical oceans (largely of isolated macroscopic colonies of the planktonic cyanobacterium, *Trichodesmium* spp. [Capone et al. 1997]), accounting for a total of 138 discrete observations of depth-integrated BNF, yielded an average rate of BNF of 1.79 mg N m⁻²day⁻¹ (or 128 μmol m⁻²day⁻¹, based on the average derived from each study) and 1.89 mg N m⁻²day⁻¹ (or 133 μmol m⁻²day⁻¹, weighted for the number of discrete observations in each study) (Table 7).

We chose 1.82 mg N m⁻²day⁻¹ (130 μmol N m⁻² day⁻¹) and simply extrapolated this value to all waters within each basin with a sea surface temperature 25°C and warmer as a proxy for oligotrophic surface waters (as derived from AVHRR monthly averages from a 1° global grid) for an estimate of pelagic BNF. This approach yielded estimates of 12, 5, 29, 19 and 20 Tg N yr⁻¹ for nitrogen fixation in the North and South Atlantic,

Table 7: Average areal rate of marine BNF. Values are means from indicated studies. Standard errors of the means are given for the individual studies, where available. For recent cruise based studies of Capone et al., stations are chosen arbitrarily along a cruise track at 0600 each morning.

Location	# stations or observations	N ₂ Fixation		N ₂ Fixation		Weighted for stations		Reference		
		μmol N/m ² * d	S.E.	mg N/m ² * d	S.E.	S.E.				
SW N. Atlantic 0°-24°N, 45°-66°W	19	41	+/-	7.6	0.574	+/-	0.106	779	pc	Goering et al. 1966
	17	108	+/-	24	1.512	+/-	0.336	1836	pc	Goering et al. 1966
N. Pacific, 21°N, 159°W	2	134			1.876	+/-	0.000	268	pc	Gunderson et al. 1976
Caribbean, 12° to 22°N	12	161	+/-	20	2.254	+/-	0.280	1932	it	Carpenter & Price 1977
SE E. China Sea, 10°-25° N	32	126			1.764	+/-	0.000	4032	tc	Saino 1977
Arabian Sea, 7-10°N	9	35	+/-	7.4	0.490	+/-	0.104	315	it	Capone et al. 1998
HOT/ ALOHA	3	84	+/-	49	1.176	+/-	0.686	252	it	Karl et al. 1997
Tropical N. Atlantic	15	258	+/-	98	3.612	+/-	1.372	3870	it	Capone et al. submitted
SW N. Atlantic	20	206	+/-	63	2.884	+/-	0.882	4120	it	Capone et al. submitted
Average	129	128	+/-	27	1.79	+/-	0.379			
Weighted Average		135			1.89					

pc = plankton concentrates
tc = *Trichodesmium* concentrates
it = isolated *Trichodesmium* colonies

Table 7. Average areal rate of marine N₂ fixation

North and South Pacific and Indian Oceans, respectively (Table 8). Total annual pelagic BNF was estimated to be 85 Tg N. This estimate provides a lower bound in that it is primarily the background contribution of *Trichodesmium* and does not capture either bloom conditions or the N₂ fixing activities of other diazotrophs. Further, this extrapolation does not imply a direct or scaled temperature control for *Trichodesmium* but rather a threshold, or step function, above which the nominal rate derived above is scaled. It should also be recognized that other diazotrophs might not have the same temperature constraints as *Trichodesmium*.

Other important factors unaccounted for in this calculation are the input of N₂ due to dense surface aggregations or surface “blooms” and the contribution of free trichomes (Capone 2001, Carpenter et al. in press). Surface accumulations of *Trichodesmium* are episodic. Several field efforts found amplified input of nitrogen during such occurrences (e.g. see Capone et al. 1998). In some areas, the bulk of *Trichodesmium* has been reported to occur as free trichomes (rather than as aggregates or colonies) (Letelier and Karl 1996, Orcutt et al. 2001, Carpenter et al. in press). Other marine diazotrophs, such as *Richelia* the endosymbiont of some diatoms, can form large blooms and be responsible for intense N inputs through BNF (Carpenter et al. 1999). *Trichodesmium* also occurs in cooler waters (i.e. < 25°C) and this input would not be captured in the current extrapolation.

Nitrogen fixation has also been documented in a range of shallow marine habitats. Capone (1983) summarized data for a range of benthic habitats and provided the only available global estimate of nitrogen fixation by the benthic components of marine ecosystems, about 15.4 Tg N yr⁻¹. In a subsequent analysis, Howarth et al. 1988, suggested that areal rates used by Capone (1983) may be high, perhaps in some cases by an order of magnitude. Hence, for our conservative estimate of benthic nitrogen fixation, we have taken the areal estimates of Capone (1983) for particular ecosystems, and reduced them by a factor derived from the ratio of the average areal rates derived in these two studies for particular types of ecosystems. This input was apportioned among the ocean basins based on the approximate relative extent of < 200m shelf among the 3 basins (i.e. 4:3:1 Atlantic to Pacific to Indian). The resulting values, while finite, are insubstantial relative to the water column values. Similarly, BNF in deep sea sediments was assumed to be negligible (Capone 1983). We therefore estimate that, conservatively,

Table 8: N fixation and denitrification by ocean basin

Basin	N ₂ Fixation Tg Ny		HIGH N ₂ Fixation Tg Ny		Denitrification Tg Ny		HIGH Denitrification Tg Ny		PON Flux Tg Ny	N ₂ O Flux [*] Tg Ny	
N Atlantic	12 [*]	This exercise	42 [*]	Gruber & Sarmiento 97	0		3	Codispoti et al. 01		0.364 C. Nevison, pers. comm.	
	0.38	Howarth et al. 88	3.7	Capone 83	15	Christensen et al. 87	75 [*]	Devol 91	3.9 [*]		Hedges & Keil 95
	0		0		1.2	This exercise	2.4 [*]	Bender et al. 77	0.11 [*]		
S. Atlantic	5 [*]	This exercise	21 [*]		0		2	Codispoti et al. 01		0.476 C. Nevison, pers. comm.	
	0.38	Howarth et al. 88	3.7	Capone 83	15	Christensen et al. 87	75 [*]	Devol 91	3.9		Hedges & Keil 95
	0		0		1.2	This exercise	2.4 [*]	Bender et al. 77	0.11 [*]		
N. Pacific	29 [*]	This exercise	35 [*]	Deutsch et al. 01	22	Deutsch et al. 01	40	Codispoti et al. 01		1.20 C. Nevison, pers. comm.	
	0.28	Howarth et al. 88	2.8	Capone 83	14	Christensen et al. 87	70 [*]	Devol 91	3.7		Hedges & Keil 95
	0		0		2.1	This exercise	4.2 [*]	Bender et al. 77	0.20 [*]		
S. Pacific	19 [*]	This exercise	24 [*]	Deutsch et al. 01	26	Deutsch et al. 01	40	Codispoti et al. 01		0.756 C. Nevison, pers. comm.	
	0.28	Howarth et al. 88	2.8	Capone 83	7	Christensen et al. 87	35 [*]	Devol 91	1.9		Hedges & Keil 95
	0		0		2.1	This exercise	4.2 [*]	Bender et al. 77	0.20 [*]		
Indian Ocean	20 [*]	This exercise	19 [*]	Bange et al. 00	33	Bange et al. 00	65	Codispoti et al. 01		0.90 C. Nevison, pers. comm.	
	0.19	Howarth et al. 88	1.9	Capone 83	6.4	Christensen et al. 87	32 [*]	Devol 91	1.6		Hedges & Keil 95
	0		0		1.8	This exercise	3.6 [*]	Bender et al. 77	0.17 [*]		
All Pelagic	85		141		81		150		x		
All Shelves	1.5		14.9		57		287		15.0		
All Deep	0		0		8.4		17		0.79		
TOTAL	86.5		156		147		454		15.8	3.696 C. Nevison, pers. comm.	

biological BNF in the oceans should account for at least 86.5 Tg N yr⁻¹ (Table 8).

For pelagic BNF, the available studies are limited in both their numbers and in the global extent of their coverage (Tables 7 and 8) and have both spatial and temporal biases. Further, direct determinations based on sample containment and incubation may suffer from potential artifacts (e.g., bottle effects) (Zobell and Anderson 1943).

Moreover, recent evidence suggests 1) a much greater diversity of diazotrophs may be contributing to overall BNF for which we do not currently have robust direct estimates because of their small size and relatively dilute concentrations *in situ* (Zehr et al. 2000; Capone 2001; Zehr et al. 2001; Dore et al. 2002) and 2) much higher *in situ* rates based on geochemical analyses using mass balance (e.g. Karl et al. 1992), stable isotope mass balance (e.g. Karl et al. 1997; Brandes et al. 1998) or N*, an index of the relative regeneration of N and P below the euphotic zone (e.g. Michaels et al. 1996; Gruber and Sarmiento 1997, see below).

Geochemical estimates of Oceanic N₂ Fixation

Recent evidence indicates that there are likely other sources of BNF in the open ocean in addition to *Trichodesmium* (Zehr et al. 2000; Capone 2001) and thus, it is not surprising that several geochemical analyses generally suggest a greater overall input than estimates based on direct scaling of rates by *Trichodesmium* alone. Geochemical indices (e.g. $\delta^{15}\text{N}$ natural abundance (Brandes et al. 1998), N* (Gruber and Sarmiento 1997) and dissolved inorganic carbon removal in the absence of fixed nitrogen (Michaels et al. 1996)) provide integrative measures of the net effect of BNF, nitrogen deposition from the atmosphere, and denitrification. For our second analysis we used estimates of pelagic BNF derived either directly or indirectly from a parameter termed N* and from mass-balance calculations on basin scales.

The N* approach calculates a metric that compares the amount of nitrate in a water mass with that predicted from the remineralization of planktonic material with a characteristic C:N:P ratio of 106:16:1, generally referred to as the Redfield ratio (Redfield 1958; Falkowski et al. 2000; Michaels et al. 2001).

This metric is usually some variant of the form:

$$\text{N}^* = [\text{NO}_3] - 16 \times [\text{PO}_4]$$

The variants on this metric take into account the varying composition of remineralization materials.

N* decreases if organic matter is exported and remineralized at a N:P ratio higher than 16, such as has been found in diazotroph-dominated plankton communities, or if phosphorus is preferentially removed from the water column after organic mineralization has taken place. N* decreases when nitrate is removed by denitrification, if exported organic matter is phosphate-rich or organic matter rich in phosphorus is remineralized. The changes in the N* value of a water mass are a balance of these processes. Thus an N* near zero can indicate the absence of nitrogen fixation and denitrification or, equally, very high rates of each within the same water mass since the time of its formation. Persistent oceanic gradients in N* can indicate mixing between areas of BNF and denitrification and a knowledge of residence times in these water masses can allow for calculation of the rate of net nitrogen transformation in each area. Michaels et al. (1996) derived a N* based estimate of BNF in the N. Atlantic of between 52 and 90 Tg N yr⁻¹ using the observed profiles of N* in the middle of the Sargasso Sea and the residence

time of each layer of water. Gruber and Sarmiento (1997), using a more elegant N* approach, estimated an annual input of 28 Tg N for the North Atlantic Ocean based on N*. In their analysis, they assumed an N:P ratio for diazotrophs of 125:1 (see their Figure 18) based on one observation of the N:P ratio in surface particulate matter after a *Trichodesmium* bloom in the Pacific (Karl et al. 1992). However, subsequent field observations indicate that the N:P ratio in *Trichodesmium* is generally much lower and in the range of 30-50 (Letelier and Karl 1996, 1998). Using a value of 40 for the N:P would result in an estimate of BNF of about 42 Tg N yr⁻¹ for the North Atlantic (see Gruber and Sarmiento 1997). Both of these estimates are for net nitrogen fixation rates, and the presence of denitrification in these waters would require a comparably higher gross nitrogen fixation rate. The estimate of 15 Tg N yr⁻¹ for North Atlantic shelf denitrification (Table 8) may mean that gross pelagic BNF estimates are much higher.

Data on the distribution and activity of diazotrophs in the South Atlantic are relatively sparse. While the positive N* anomalies of the S. Atlantic are not nearly as strong as in the North Atlantic (possibly because of lower aeolian dust fluxes and iron inputs which may limit rates of nitrogen fixation, see below), the area of oligotrophic surface waters suitable for *Trichodesmium* and other diazotrophs is only about 42% that in the N. Atlantic. Hence, we have assumed that average areal rates of BNF in the S. Atlantic are half those in the North Atlantic.

In a N* analysis of the Pacific Ocean, Deutsch et al. (2001) were unable to use N* to derive BNF because of the basin-scale impact of denitrification (i.e. complex admixtures in the surface gyre of waters with and without the N* imprint of denitrification) and instead took a mass balance approach to estimate BNF. They calculated an input of 59 Tg N yr⁻¹ to balance denitrification and N transport. The areal rates derived from the Deutsch et al. (2001) calculation, about 546 mg N m⁻² yr⁻¹, agree very favorably with areal estimates from the HOT station (Karl et al. 1997; Capone 2001). We have apportioned this input based on the distribution of warm surface waters of the Pacific (S. Pacific has about 40% of the area of warm surface waters). However, as mentioned above, the input of Fe from terrigenous sources is much greater in the North Pacific, coincident with positive N* anomalies in the northwest Pacific, while aeolian sources of iron to the South Pacific are minimal, and this may have some bearing the relative proportions of BNF in the 2 basins (Behrenfeld and Kolber 1999; Wu et al. 2000; Gao et al. 2001; Karl 2002).

For the Indian Ocean, we scaled our nominal rate of BNF in warm surface waters to the > 25° C area based on average SST (Table 8). For our geochemical-based estimate, we used the estimate of Bange et al. (2000) of 3.3 Tg N yr⁻¹ by BNF for the Arabian Sea, an area of about 4.9 x 10⁶ km². Their estimate was an average based on the ¹⁵N isotope mass balance of Brandes et al. (1998), which derived an areally integrated BNF in the central portion of the Arabian Sea (1.2 x 10⁶ km²) to be about 6 Tg N yr⁻¹ and the data based upon the study of Capone et al. (1998) for the spring inter-monsoon, which, when scaled to the same area, yielded an input of 0.6 Tg yr⁻¹. The annual average area of warm surface waters of the Indian Ocean proper is about 30 x 10⁶ km². Thus, we scaled the Bange et al. (2000) estimate for the central Arabian Sea over this broader area which yielded about 19 Tg N yr⁻¹ for the entire Indian Ocean basin, very close to the estimate based on our nominal BNF rate for *Trichodesmium* scaled to the basin (Table 8).

Our geochemical estimate of annual pelagic BNF is thus 141 Tg N. For the complementing benthic component, we use the more liberal estimates of Capone (1983), for salt marshes (6.3 Tg N yr⁻¹), coral reefs (2.8 Tg N yr⁻¹), mangroves (1.5 Tg N yr⁻¹), seagrasses (1.5 Tg N yr⁻¹), and estuarine and shelf sediments above 200 m of depth (3.1 Tg N yr⁻¹), yielding a total benthic input (including minor of about 15 Tg N yr⁻¹). This was apportioned among basins as for the more conservative estimate. Thus, our total oceanic BNF is estimated to range from about 87 Tg N yr⁻¹ to 156 Tg N yr⁻¹. For the purposes of illustration, we use the mean value of 121 Tg N yr⁻¹ in Figure 1 and assume that it does not vary with time.

Lee et al. (2002) have recently conducted a global analysis of inorganic C removal from nitrate depleted tropical surface waters. They have derived a new production rate of 0.8 ± 0.3 Pg C yr⁻¹ which they conclude implies a BNF input of 114 ± 43 Tg N yr⁻¹ (based on a C:N ratio of 7). This is largely consistent with our analysis. Based on global isotope budgets, Brandes et al. (2002) have also recently concluded that nitrogen fixation must be much greater than 100 Tg N yr⁻¹.

Iron issues/ distributions of BNF

In apportioning the amounts of pelagic BNF among and within basins, we have simply assumed a direct relationship between the extent of warm waters and BNF. However, there appears to be a direct and strong correlation between the input of iron by aeolian flux. Areas with large dust inputs (tropical N Atlantic, N Pacific) have very high positive N* anomalies (Michaels et al. 1996; Gruber and Sarmiento 1997; Gao et al. 2001).

Recent results indicate strongly that BNF in the N. Atlantic basin is less likely to be constrained by Fe compared to the N. Pacific (Wu et al. 2000), but may be more directly limited by P availability in the upper water column (Sañudo-Wilhelmy et al. 2001). Hence, the extent of iron and phosphorus limitation of pelagic BNF should be more comprehensively assessed to better understand the distribution and extent of oceanic BNF.

Denitrification

Denitrification for entire ocean and within basins

Pelagic denitrification is not thought to be a major process in the basins of the North or South Atlantic Ocean. However, substantial water column denitrification does occur in anoxic plumes in the Eastern Tropical North (ETNP) and South (ETSP) Pacific and in the Arabian Sea because of the large-scale circulation of the deep ocean. The proximal cause of these anoxic plumes is upwelling, increased surface productivity and vertical flux of organic matter over areas of slow mid-water circulation and poor ventilation. However, the larger scale oceanic circulation known as the “conveyor belt” also contributes to the development of these plumes. Deep water is formed in the N. Atlantic where cold surface water sinks, losing contact with the atmosphere. Those waters transit south through the deep Atlantic, picking up more sinking water around Antarctic. Deep waters then traverse into the Indian Ocean, and through the South Pacific with the oldest waters finally reaching the deep North Pacific. Thus, the oldest waters have been out of contact with the atmosphere for the longest period, have the lowest O₂ content (highest apparent oxygen utilization) and highest concentrations of nitrate and phosphate, as a result of the cumulative effect of microbial respiration. Large

areas of hypoxic waters occur in the Indian Ocean and in the ETSP and ETNP and are globally significant sites of denitrification.

We have used Deutsch et al.'s (2001) N^* based estimates for denitrification in the ETNP and ETSP of 22 and 26 Tg N yr⁻¹, respectively. These values are consistent with earlier estimates (Codispoti and Richards 1976). While previous studies had suggested relatively low values for pelagic denitrification in the Arabian Sea (e.g. 2 Tg N yr⁻¹, Somasundar et al. 1990) before full recognition of the size of the anoxic plume, more recent estimates are much higher, around 30 Tg N yr⁻¹ (Naqvi et al. 1992). We used the most recent estimate of Bange et al. (2000), 33 Tg N yr⁻¹ for our calculations although this does not include any denitrification occurring in sub-oxic pockets of the Indian Ocean. Thus, total oceanic pelagic denitrification is conservatively estimated at 81 Tg N yr⁻¹.

Shelf sediments are globally important sites of denitrification (Christensen et al. 1987). Denitrification rates have been measured at specific locations within the North Atlantic, North Pacific, and Arctic shelf regions (Tables 8 and 9). To provide a conservative estimate of the contribution by shelf sediments to total N flux within each basin, we chose a value of 5.6 mg N m⁻² day⁻¹ (400 μ mol N m⁻² day⁻¹) rounded up from the mean (5.3 mg N m⁻² day⁻¹ or 381 μ mol N m⁻² day⁻¹) derived from several earlier studies (Table 9), scaled to the shelf area for each basin. This yielded estimates of 30 Tg N yr⁻¹ for the Atlantic (apportioned 50% for each of the North and South Atlantic), 21 Tg N yr⁻¹ for the Pacific (apportioned by 66% to the North Pacific and 33% to the South Pacific) and 6.4 Tg N yr⁻¹ for the Indian Ocean (Table 8) for a total of 57 Tg N yr⁻¹. Christensen et al. (1987) originally estimated between 50 and 75 Tg N yr⁻¹ for a global annual rate of shelf denitrification.

Results by Devol (1991) suggest that denitrification on shelf sediments may be >5-fold higher than earlier geochemically-based estimates (e.g. compare estimates for older studies based on NO₃⁻ flux to several more recent estimates based on N₂ flux in Table 9). Also, recent analyses by Codispoti et al. (2001) and Devol et al. (1997) suggest that denitrification on the Arctic shelf, which represents 25% of all shelf area, may account for about 40-45 Tg N yr⁻¹, with the Antarctic shelf adding another 4 Tg N yr⁻¹. The estimates of shelf area we used were from Sverdrup et al. (1942, derived from Kossinna 1921 original analysis), which included shelf areas of the Arctic and Antarctic in the basins of the Atlantic, Pacific and Antarctic (e.g. compare to Menard and Smith 1966). Another recent development is the detection of the anammox reaction in shelf sediments (Thamdrup and Dalsgaard 2002) that can also contribute to the N₂ flux through the biological oxidation of NH₄⁺ by NO₂⁻.

A second, less conservative, estimate of global marine denitrification is therefore made by applying a caveat to the existing database. For shelf sediments, we increase our conservative estimate by a factor of 5 (Devol 1991) that increases the shelf contribution to 287 Tg N yr⁻¹ (Table 8). This is consistent with Codispoti et al. (2001) recent estimate of about 300 Tg N yr⁻¹.

Many of the measurements used for the extrapolations are based on sediment incubations or geochemical modeling of sediment nitrate distributions and are at relatively small scales (<1 m²). Seitzinger and Giblin (1996) took a modeling approach to estimate the larger scale spatial distribution of denitrification rates in continental shelf sediments of the North Atlantic basin. The model was based on an empirical regression

that related denitrification rates (coupled nitrification/denitrification) to sediment oxygen consumption rates which in turn were related to depth-integrated water column primary production. Data from a range of geographic locations were used to develop this model. Model predicted denitrification rates in various shelf regions within the North Atlantic basin compared favorably with measured rates (e.g., Gulf of Mexico, North Sea, South Atlantic Bight, Massachusetts Bay). Denitrification in the shelf sediments of the North Atlantic basin as a whole was estimated to be 102 Tg N yr⁻¹. Bange et al. (2000) used the model of Seitzinger and Giblin (1996) to derive an Arabian Sea shelf sediment denitrification rates of 6.8 Tg N yr⁻¹. This would scale to a total of 42 Tg N yr⁻¹ for the shelves of the entire Indian Ocean. These estimates compare favorably to the extrapolated rates under the High Denitrification scenario (Table 8). The model was also more generally applied to global shelf regions by multiplying the average model estimated denitrification rate calculated for each of three latitudinal areas in the North Atlantic Basin (Table 9) by the area of continental shelves in each of those latitudinal belts (Seitzinger and Kroeze 1998). This resulted in an estimated denitrification rate of 214 Tg N yr⁻¹ for global shelf sediments, again in reasonable agreement with the globally extrapolated value derived here. These model rates may underestimate total sediment

Table 9: Some estimates of marine sedimentary denitrification

Location	Comments	Units reported			Reference	
		pmol NO ₃ cm ² sec	μmol N cm ² y	x 10 ⁻¹⁰ g mol m ² sec		mg N m ² day
Shelves	North Sea	NO ₃ consump/ model	0.79		9.56	Billen 1978
	Bering Sea shelf	¹⁵ N-tracer	0.25		3.08	Koike & Hattori 1979
	Bering Sea shelf	C ₂ H ₂ Blockage	0.45		6.02	Haines et al. 1981
	East China Sea	geochem model	0.38		4.60	Aller et al. 1985
	Mexican shelf	geochem model	0.15		1.83	Christiansen et al. 1987
	Washington shelf	geochem model	0.57		6.89	Christiansen et al. 1987
	Average of shelves- older values					5.33
	Baltic Sea	geochem model	2.6		31.4	Shaffer & Ronner 1984
	Mid-Atlantic Bight	N ₂ flux			23.1	Laursen and Seitzinger 2002
	Washington state shelf	N ₂ flux	3.7		44.8	Devol 1991
	Chukchi	N ₂ flux			15.4	Devol 1997
	Bering	N ₂ flux			7.42	Devol 1997
	Greenland	N ₂ flux			8.82	Devol 1997
	Svalbard	N ₂ flux			11.34	Devol 1997
	Average for all					13.40
	Latitudes 45-90°	modeled	19.7		7.56	Seitzinger & Kroeze 1998
	Latitudes 20-45°	modeled	57.6		22.1	Seitzinger & Kroeze 1998
	Latitudes 0-20°	modeled	31.4		12.0	Seitzinger & Kroeze 1998
Deep Sea	Eastern eq. Atlantic	standard denitrif stoichiometry	0.13		0.05	Bender et al. 1977
		Anammox stoichiometry	0.29		0.11	Bender et al. 1977
	Northeast Atlantic	N ₂ gas profile		0.6	0.15	Wilson 1978

denitrification because they only include coupled nitrification/denitrification and not denitrification of nitrate diffusing into the sediments from the overlying water.

Codispoti et al. (2001) have recently proposed that marine pelagic denitrification has been substantially underestimated, and suggest an annual rate of 150 Tg N which we have used and apportioned largely to the Indian and Pacific Oceans on the basis of their analysis, with minor amounts of denitrification assigned to the North and South Atlantic associated with anoxic basins (e.g. Cariaco Trench and Baltic Sea) and aerobic waters (Codispoti et al. (2001).

For deep sediments, Bender et al. (1977) suggested that, depending on the stoichiometry of denitrification in the deep sea, rates of denitrification based on NO_3^- diffusion could be twice as great. Applying this factor would yield a total abyssal rate of 17 Tg N yr^{-1} (Table 8). In this regard, a recent geochemical modeling study (Middleburg et al. 1996) concluded that deep sea denitrification might make a considerably greater contribution to oceanic denitrification, estimated to be 100-130 Tg yr^{-1} .

Thus, our conservative estimate based on extrapolation of total (pelagic plus benthic) oceanic denitrification is 147 Tg N yr^{-1} . Using the more liberal assumptions for shelf sediment denitrification, an upper limit for total oceanic denitrification would be about 454 Tg N yr^{-1} .

For the purposes of illustration, in Figure 1 we use the mean value of 116 Tg N yr^{-1} for pelagic denitrification and the mean value of 13 Tg N yr^{-1} for deep sediment denitrification to yield a total of 129 Tg N yr^{-1} for denitrification in the open ocean and assume that it does not vary with time. Our range of shelf denitrification is 57 Tg N yr^{-1} to 287 Tg N yr^{-1} , with a mean of 172 Tg N yr^{-1} , which we apply for 1860 as a natural background. For the early-1990s and 2050, we add to this value the additional Nr injected into coastal regions based upon the assumption that all riverine Nr is denitrified in estuaries or shelf regions. Thus for the early-1990s and 2050 denitrification from estuaries and shelves produces 193 and 210 Tg N yr^{-1} of N_2 , respectively.

Particulate Organic Nitrogen (PON) Flux and Storage in Sediment

Finally, we (guess)timated the flux of PON into oceanic sediments. Hedges and Keil (1995) suggested an annual burial rate of C for the oceans of 160 Tg C yr^{-1} . They further suggested that this burial occurred in the proportion of 45% in deltaic sediments, 45% in coastal margin regions, 5% under coastal upwelling areas and 5% on the continental rise and in abyssal sediments. We assumed a C:N ratio of 10:1 (Jahnke, pers. comm.), and a similar proportional distribution of the various shelf types among the ocean basins. Hence, we suggest a shelf burial rate in the Atlantic of 7.8 Tg N yr^{-1} (apportioned equally between the north and south Atlantic), 5.7 Tg N yr^{-1} in the Pacific (66% in the North Pacific and 33% in the South Pacific) and 1.7 Tg N yr^{-1} in the Indian Ocean. The global total for the early-1990s is ~15 Tg N yr^{-1} . For 1860 and 2050, we simply scale PON sediment burial on the shelf to the riverine flux (Figure 1).

PON burial in the deep sea is estimated to be 0.11 Tg N yr^{-1} in each of the North and South Atlantic, 0.2 Tg N yr^{-1} in each of the North and South Pacific and 0.17 Tg N yr^{-1} in the Indian Ocean (Figure 2). Bange et al. (2000) have proposed a sedimentation rate of 0.26 Tg N yr^{-1} for the central and northern portions of the Arabian Sea. The global total is 0.8 Tg N yr^{-1} ; given the the lack of connection between the continental and marine N budgets we assume that this rate also applies to 1860 and 2050 (Figure 1).

N₂O

While not a major constituent in the oceanic nitrogen budget, N₂O contributes to greenhouse warming and ozone depletion, so it has been the subject of numerous studies. Concentrations of atmospheric N₂O have increased over the past several decades (Khalil and Rasmussen 1982; Nevison et al. 1995). Marine N₂O emissions contribute to the overall sources of this important trace gas. The marine source strength, including continental shelf regions, is thought to be ~ 4 Tg N yr⁻¹ (Nevison et al. 1995) (Figure 1b). Estuaries may contribute another 0.2 Tg N yr⁻¹ of N₂O (Seitzinger and Kroeze 1998).

Using the N₂O gas flux model of Nevison et al. (1995 and pers. comm.), the greatest fluxes of N₂O occur in the North Pacific, Indian Ocean and South Pacific, all with major zones of anoxic waters. Lower effluxes occur from the North and South Atlantic. The N₂O flux from the Southern Ocean (waters below 60° S) was estimated to be about 0.2 Tg N yr⁻¹. However, this flux was apportioned into the three major southern hemisphere basins, the South Pacific, Indian Ocean and South Atlantic in a ratio of 3:2:2, respectively.

Ocean Basin Budgets

Global

The global ocean contains ~8 x10⁵ Tg N of nitrate (Mackenzie 1998). It currently (1995) receives most of its Nr additions from marine BNF (87 - 156 Tg N yr⁻¹), riverine input of total N (48 Tg N yr⁻¹) and atmospheric deposition of NO_y and NH_x (33 Tg N yr⁻¹) (Figure 1b)). Nr losses are primarily from denitrification formation of N₂ (150 – 450 Tg N yr⁻¹), and minor losses due to PON storage in shelf and deep sediments (14 Tg N yr⁻¹ and 0.8 Tg N yr⁻¹, respectively), and N₂O emission (~4 Tg N yr⁻¹) from the shelves and open ocean. Although the range of Nr sources overlaps that of Nr sinks, it is clear that there is substantial uncertainty in both BNF and denitrification.

The globally integrated estimates for both marine BNF and denitrification have increased substantially over the last several decades with the recognition of previously unrecognized sources (e.g. Zehr et al. 2001) and sinks (Christensen et al. 1987, Codispoti et al. 2001). Along with the steadily progressive increase in the respective rates, the estimated residence time of combined N in the ocean has gotten progressively shorter (Codispoti et al. 2001). Given the fluxes presented here, the average residence time of Nr in the oceans would be about 1500-5000 years (see also Codispoti et al. 2001, Brandes & Devol 2002).

Atmospheric deposition and riverine inputs of Nr are the two major connection points between continents and oceans, and their importance as Nr sources to the oceans is increasing with time. Atmospheric NO_y and NH_x deposition to oceans increased about 4-fold from 8.5 Tg N yr⁻¹ in 1860 to 33.4 Tg N yr⁻¹ in the early-1990s, with NO_y deposition being about 2x as important as NH_x deposition. The transfer of Nr from continents to ocean margins via rivers increased slightly under 2-fold between 1860 and 1990, from ~27 Tg N yr⁻¹ to ~48 Tg N yr⁻¹, respectively (Figures 1a and 1b).

The importance of the atmosphere, relative to rivers, in transferring Nr to marine ecosystems is also changing with time. In 1860, ~24% of the continental Nr supplied to oceans came from atmospheric deposition; rivers supplied ~76%. By the early-1990s, the relative importance of atmospheric deposition had increased to ~41%, and as will be seen

in a later section, by 2050, atmospheric deposition will supply almost 50%. This means that as additions of new Nr to continents increase with time, that the atmosphere becomes increasingly important relative to rivers in transferring Nr from continents to marine systems.

There is another factor that strongly influences the degree of transport of Nr to marine systems. While the increase in riverine exports of Nr has substantial consequences for the health of coastal ecosystems, most of the Nr that enters coastal regions via river flow is converted to N₂ in the coastal and shelf region and does not reach the open ocean. There is however widespread distribution of anthropogenic Nr to the oceans via the atmosphere (Paerl 1993, Cornell et al. 1995). In 1860, most coastal regions received on the order of 25-50 mg N m⁻² yr⁻¹. In the early-1990s some coastal regions of most continents were receiving up to 2,000 mg N m⁻² yr⁻¹, a greater than an order of magnitude increase in N deposition (Figures 2a and 2b). Deposition to the open ocean has also increased. In 1860 most marine regions received <50 mg N m⁻² yr⁻¹. In the early-1990s most of the North Atlantic received >100 mg N m⁻² yr⁻¹, and large portions of the Pacific and Indian Oceans received >50 mg N m⁻². In addition, the deposition patterns show clear transport patterns from North America to Europe and Africa to South America (Figure 2a and 2b).

Basin

Given the general paucity of data for most marine regions, it is not surprising that the ranges of the basin estimates for BNF and denitrification are large and in some cases do not overlap (Figure 4). In fact, there may be imbalances in these processes within basins.. As put forth above, it is evident that there are substantial differences in the intensity of pelagic denitrification among the major basins. The high rates of denitrification in the Indian Ocean, ETNP and ETSP probably account for these imbalances. Similarly, the geochemical N* evidence suggests that BNF may be a much more important process in the North Atlantic (and perhaps the South Atlantic) than in the other major basins, possibly because of the substantial iron fluxes into this system.




Figure 4: N₂O creation and distribution by ocean basin, Tg N yr⁻¹, in the early-1990s

However, a recent re-analysis of the distribution of the excess nitrate anomaly in the N. Atlantic by Hansell et al. (in press) concluded that N_2 fixation was far less than the N^* estimate by Gruber & Sarmiento (1997).

The ocean receives new N_r by riverine injection, atmospheric deposition and BNF. The relative importance of river injection and atmospheric deposition compared to BNF is largest for the North Atlantic and North Pacific, and lowest for the South Pacific, in keeping with the patterns of N_r on upstream and upwind continental regions (Figure 4). However, as discussed previously, most N_r injected into coastal systems is denitrified, thus inputs of new N_r to the open ocean are either by BNF or atmospheric deposition. In the early-1990s, the only ocean basin where atmospheric deposition is in the same range as BNF is the North Atlantic Ocean (Figure 4). However, given the increase in N_r emissions projected to occur over the next few decades, it is likely that other basins will receive substantial increases in atmospheric deposition (Figure 1c). As N_r creation rates increase in the future, that external supply of N_r to ocean basins will become increasingly important relative to marine BNF.




Figure 1c: Components of the global nitrogen cycle for 2050. See legend of Figure 1a for details.

Outlook

The apparent imbalance in sources and sinks of the marine N cycle has important implications with respect to future trends in the marine N cycle, with direct ramifications

to marine productivity. At the estimated excess of losses over sources (assuming a mean rate for marine BNF and denitrification of 122 and 319 Tg yr⁻¹, respectively) of about 160 Tg N/y, about 2.0% of the current stocks of nitrate would be removed each century (see also Codispoti et al. 2001, Brandes & Devol 2002). A pressing question, therefore, is whether this apparent imbalance is real or if, as has been suggested, marine BNF is still substantially underestimated (Zehr et al. 2001, Capone 2001, Brandes, 2002).

The oceanic N cycle is clearly dynamic at various time scales as can be gleaned from the geological record. Large fluctuations in major processes appear to occur over long time scales (e.g. glacial-interglacial) periods and there is not *a priori* reason to expect that the oceans are in balance with respect to its nitrogen cycle (and many reasons to think otherwise). A current hypothesis is that during warm interglacial periods, such as the present, denitrification exceeds BNF resulting in a general decline in oceanic N inventories, while during glacial periods, an increase in iron flux to the oligotrophic ocean (coupled with a decreased denitrifying shelf area) allows an excess of BNF relative to denitrification (Falkowski 1997; Broecker and Henderson 1998). Support for this model comes from the systematic variability in the isotopic concentration of N in organic matter in deep sea cores, with isotopically lighter nitrogen deposited during glacial periods and heavier N during interglacial periods (Altabet et al. 1995; Farrell et al. 1995; Ganeshram et al. 1995; c.f. Haug et al. 1998). This has been interpreted as evidence for diminished denitrification, relative to BNF, in the glacial ocean. However, Ganeshram et al. (2002) have recently speculated that while denitrification appears to have declined and marine N inventories may have increased during glacial periods, the concomitant increases in phosphorus were less leading to higher N:P ratios and a negative feedback to marine BNF. Most recently, Gruber (in press) and others (D. Sigman, pers. comm.) have alternatively proposed a much closer coupling of nitrogen fixation and denitrification over glacial-interglacial cycles.

Substantial fluctuations in marine N dynamics have occurred within the Holocene. For instance, Emmer & Thunell (2000) and Suthof et al. (2001) have inferred higher frequency fluctuations in denitrification over the last 50,000 years, with a general increase in the extent of this process in the ETNP and Arabian Sea, respectively, since the last glacial interrupted with a decrease, for instance, during the Younger-Dryas event 10,000 years ago.

On even shorter time scales, climate oscillations such as the El Nino/ Southern Oscillation (ENSO), The North Atlantic Oscillation (NAO) and the Pacific Decadal Oscillation (PDO) have direct impacts on large-scale climatology, rainfall patterns, nutrient delivery, as well as on upper water column thermal structure, circulation and primary productivity. the intensity of continental shelf denitrification is tied to patterns of N delivery and primary productivity on the shelves forced by these climate oscillations (Seitzinger and Kroeze 1998).

Similarly, upper water column BNF should also be sensitive to nutrient delivery and upper water column stability. Decadal scale variability in BNF and a shift towards more chronic P limitation has been noted at Station ALOHA in the subtropical North Pacific Ocean coincident with a period of shallow stratification and greater oligotrophy (Karl et al. 1997; Karl 1999). However, whether this variability exists at the basin scale for BNF and whether denitrification exhibits similar decadal scale variability is presently unknown.

While it is probably somewhat premature to attempt to project such fluctuations in the marine N cycle in the future, we can speculate about some of the possible forcings. The broad trend in warming in the upper ocean (Levitus et al. 2000) could lead to both greater upper water column stability through much of the ocean as well as decreased circulation in the deep ocean (Sarmiento et al. 1998; Joos et al. 1999), both with significant implication to the marine N cycle. Reduced circulation and increased stratification may result in the expansion of oxygen minimum zones and therefore expanded zones of pelagic denitrification (Codispoti et al. 2001). Reduced upwelling and greater upper water column stability in the tropics may result in oligotrophy fostering increased diazotrophy (Sachs & Repeta 1999).




Figure 2c: Spatial patterns of total inorganic nitrogen deposition in 2050, mg N m⁻² yr⁻¹.

The delivery of combined nitrogen to the ocean through both the atmosphere and by riverine input is projected to increase steadily over the upcoming centuries (Figure 2c). In coastal regions subjected to major nutrient perturbations and eutrophication, such as the Gulf of Mexico, short term human effects are already clearly manifest (Howarth 1998). Furthermore, the increasing areas of hypoxia in some of these areas could have direct implications with regard to the flux of trace gases such as N₂O from

marine sources, and thereby a short term impact on atmospheric N₂O. (e.g. Naqvi et al. 2000, Codispoti et al. 2001). The extent of human intervention and impact in the major components of the open ocean's N cycle will be less easy to directly discern because of the natural fluctuations mentioned, the relatively slow circulation of the deep sea, the long residence time (>1000 y) of major nitrogen pools, such as NO₃⁻ relative to the longevity of individual humans (and researchers!).

It has been argued that iron is an important determinant of oceanic BNF (Michaels et al. 1996, Falkowski 1997, et al. 1998, Gruber & Sarmiento 1997) and the primary source of iron to the upper ocean is through the deposition of aeolian dust. Human activities are actively modifying the flux of this iron source to the oceans, for instance through reforestation and improved agricultural practices (Metz et al. 2001). Recent modeling efforts have predicted reduced primary production in the Oceans due to a reduction in dust flux from (Ridgwell et al. 2002). Oceanic BNF may also be expected to be adversely affected (Michaels et al. 2001). Ironically, purposeful large-scale fertilization of the upper ocean with iron in order to promote sequestration of atmospheric CO₂, as has been proposed (e.g. www.greenseaventure.com/iron2.html), could also have unintended effects with respect to the N cycle (e.g. Fuhrman and Capone 1991, Codispoti et al. 2001).

Projections to 2050

Introduction

The global population is projected to be ~9 billion people in 2050, an increase of 50% over 2000. In addition to the increase in population, there will likely be increases in per-capita energy and food consumption in many parts of the world. These changes will result in additional Nr creation via the Haber-Bosch process, cultivation-induced BNF and fossil fuel combustion, and increased Nr distribution via atmospheric and hydrologic pathways. For this analysis we present a range of Nr creation rates on a global basis, and then within each of those ranges we select a value to use to illustrate the inputs of Nr to the global N cycle (Figure 1c).

Nr Creation

Natural BNF

In the absence of people, BNF in natural terrestrial systems created ~128 Tg N yr⁻¹. By 1995, we estimate that the addition of 5.8 billion people reduced BNF occurring in natural landscapes to ~107 Tg N yr⁻¹. The addition of 50% more people to the planet will certainly require the conversion of natural landscapes to altered ones. As a consequence, the processes that occurred in those natural landscapes, including BNF, will be reduced in scope. As a rough estimate we scale the reduction in natural BNF inversely to the increase in population between 1990 and 2050. The resulting estimate is 98 Tg N yr⁻¹. To approximate the regional distribution of the global BNF in 2050, we assumed that the ratio of regional BNF to global BNF (in full or active watershed boundaries, as appropriate) remained the same in 2050 as it was in 1990.

Haber-Bosch

For 2050 we estimate that N fertilizer consumption will be ~ 135 Tg N yr⁻¹ (Appendix II). We use this estimate in two ways. First it is a Nr input to the terrestrial landscape used to calculate riverine Nr fluxes in 2050 with the NTNI model. Second it serves as the basis for estimating total Haber-Bosch production of Nr. In that regard, we make two adjustments to the 135 Tg N yr⁻¹. The first is based on the fact that not all fertilizer-N that is produced is consumed, and the second is based on the fact that not all Haber-Bosch production of Nr is used for fertilizer. Over the period 1990 to 2000, the annual average fertilizer consumption was 7% less than the fertilizer production (FAOSTAT 2000). (We assume that this is due to post-production losses, but other possible explanations are incorrect data or differences in accounting of production vs. consumption.) Over the same period, the annual average amount of fertilizer produced via the Haber-Bosch process was 15% less than the total Nr created by Haber-Bosch (FAOSTAT 2000). Making these two adjustments gives an estimate of ~165 Tg N yr⁻¹ of Nr produced by the Haber-Bosch process (Figure 1c), ~145 Tg N yr⁻¹ produced by Haber-Bosch to be used as fertilizer, and our base figure, ~135 Tg N yr⁻¹ of Haber-Bosch fertilizer N that was consumed.

These estimates of fertilizer use can be compared to other estimates for fertilizer use in 2050 (future projections of the total amount of Nr created by the Haber-Bosch process are lacking). IMAGE (2001) estimates that N fertilizer use in 2050 for the four

B1, B2, A2 and A1 SAES scenarios range from 140 Tg N yr⁻¹ to ~200 Tg N yr⁻¹, respectively. Tilman et al. (2001) estimate a range of 165 Tg N yr⁻¹ to 339 Tg N yr⁻¹. The magnitude of the ranges for both studies reflects the number of factors (and associated uncertainties) that determine future rates of fertilizer use.

Cultivation-induced BNF

Nr creation by cultivation-induced BNF increased from ~21 Tg N yr⁻¹ in 1961 to ~31.5 Tg N yr⁻¹ in 2000 (FAO 2000; Smil 1999). On a per-capita basis, cultivation induced Nr creation rate was ~7 kg N yr⁻¹ in 1961, and steadily decreased to 5.5 kg N yr⁻¹ in 1991 where it remained relatively constant to date. If we assume the *per-capita* rate remains constant, then by 2050 (global population of 9 billion), Nr creation by cultivation-induced BNF would be ~50 Tg N. This number is very uncertain and the correct value could be higher (i.e., increased reliance on C-BNF to provide Nr) or lower (conversion of forage-producing land to animal-producing land). An additional factor is the fact that much of the additional land used for cultivation-induced BNF will be in the tropics, which increases the possibility for larger per area losses of natural BNF, and makes more uncertain the fate of the replaced land. An additional complicating factor is that many systems in the tropics are P-limited. Given the uncertainty about the number, we choose a range of 45 Tg N yr⁻¹ to 55 Tg N yr⁻¹, with the mean value of 50 Tg N yr⁻¹ (Figure 1c). It should be noted however, that for our 2050 projections for riverine Nr discharge that we used a value of 31.5 Tg N yr⁻¹.

Combustion of fossil fuels

There are large ranges in the estimates of Nr creation by fossil fuel combustion in 2050. IPCC (1996) estimate a range of ~40 Tg N yr⁻¹ to ~70 Tg N yr⁻¹. More recently, IPCC (2001) give a range of 38.8 Tg N yr⁻¹ to 94.9 Tg N yr⁻¹, depending on the scenario used. For our analysis, we use 52.2 Tg N of Nr created by fossil fuel combustion in 2050 (Table 1; Figure 1c), which is well within the range of the IPCC estimates.

Nr Distribution

Atmospheric emission and deposition

We have used specific scenarios to examine dispersion of Nr via atmospheric and hydrologic pathways for 2050. For the atmosphere, emissions of NO_x and NH₃ were based on the activity data underlying the IS92a scenario (IPCC 1996). When comparing this scenario for NO_x with the very recent IPCC-TAR scenario, the IS92a emissions are between the 'B2' and 'A2' scenario. A2 corresponds to a situation where the world develops according to a business-as-usual scenario, and large differences in technology are found in the various regions around the world. Also scenario B2 assumes a heterogeneous development of the world economy, but within all regions there is more social awareness (e.g., on pollution issues).

IPCC has never developed the corresponding scenarios for NH₃. Comparing 2050 to the early-1990s, our estimate based on IPCC92a calculates an increase of anthropogenic emissions by a factor 2.3. Although certainly not fully comparable, we may compare this increase with the corresponding increases of anthropogenic N₂O emissions (see below). Given the specific issues involving anthropogenic ammonia emissions, it seems that the wide range of possible developments that are possible, and

the emissions used in this study have substantial uncertainty. However, when coupled with modeling estimates of deposition they do show the regions that will experience the largest increases in deposition.

The large increases in NO_x and NH_3 emissions drastically change the pattern and magnitude of deposition of total inorganic nitrogen (Figure 2c). Asia has the largest changes. Relative to the early-1990s the area receiving $>1,000 \text{ mg N m}^{-2} \text{ yr}^{-1}$ and $>2,000 \text{ mg N m}^{-2} \text{ yr}^{-1}$ has grown significantly, and for the first time, large regions of South and East Asia are projected to receive $>5,000 \text{ mg N m}^{-2} \text{ yr}^{-1}$. Other regions also exhibit increases. In the early-1990s, a small region of SE South America received inorganic N_r deposition $>1,000 \text{ mg N m}^{-2} \text{ yr}^{-1}$. By 2050, this area has grown significantly and there is a large region receiving $>2,000 \text{ mg N m}^{-2} \text{ yr}^{-1}$. There is a similar change in Central America with deposition increasing from $<750 \text{ mg N m}^{-2} \text{ yr}^{-1}$ over most of the region to $>2,000 \text{ mg N m}^{-2} \text{ yr}^{-1}$ for almost the entire region. Inorganic nitrogen deposition increases in the central portion of Africa from $<750 \text{ mg N m}^{-2} \text{ yr}^{-1}$ to $>1,000 \text{ mg N m}^{-2} \text{ yr}^{-1}$. As might be expected to its existing level of emissions, North America does not experience significant increases. Neither does western Europe but southern and Eastern Europe have significant increases in deposition.

With the increased emissions on continents, deposition of inorganic nitrogen to downwind oceans increases. While there are slight increases in the North Atlantic Ocean (deposition over most of the region is $>100 \text{ mg N m}^{-2} \text{ yr}^{-1}$), the North Pacific has the largest increases, with large regions receiving $>100 \text{ mg N m}^{-2} \text{ yr}^{-1}$.

IPCC (2001) makes several estimates of anthropogenic N_2O atmospheric emissions from continents in 2050 that range over a factor of ~ 2 with the minimum being less than their estimates of current emissions, and the maximum being significantly greater than current emissions. Given this uncertainty we do not choose to make an estimate for 2050. For aquatic ecosystems increased anthropogenic N inputs to rivers and estuaries are expected to increase the emissions of N_2O . By 2050 N_2O emissions from rivers are predicted to increase from the 1990 level of 1 Tg N yr^{-1} to 3.3 Tg N yr^{-1} , while estuarine emissions are predicted to increase from 0.2 Tg N yr^{-1} to 0.9 Tg N yr^{-1} (Kroeze and Seitzinger 1998) (Table 2b, Figure 1c). However, as has been noted above, there are considerable uncertainties about N_2O emissions at the global scale for any one source and with an increasing uncertainty as one attempts to project into the future.




Figure 5: Riverine N_r export to the coastal zone (Tg N yr⁻¹) in the past (1860), present (1990) and future (2050). Dry and inland watershed regions that do not transmit to coastal areas are shown in gray.

Riverine transport

As described in Appendix II, using moderate increases for fertilizer consumption and nitrate deposition, and assuming a conservative scenario where there are no increases of cultivation-induced BNF or decreases in natural BNF, Boyer et al. (in preparation) estimate that in 2050 Nr transport to inland receiving waters will be $\sim 11.7 \text{ Tg N yr}^{-1}$ and to the coastal ocean will be $\sim 63.2 \text{ Tg N yr}^{-1}$ (Figure 5).

Riverine transport vs. atmospheric emissions

Riverine and atmospheric transport are both vectors for the distribution of Nr. Over the period 1860 to the early-1990s the amount of Nr transported has increased and is projected to keep on increasing to 2050 (and probably beyond). However the rate of increase is quite different for the two vectors. Relative to the amount of Nr creation, the emission of NO_x plus NH_3 to the atmosphere is increasing much faster than the discharge of Nr to the coastal zone (Figure 6). The reasons for the relatively limited response of riverine systems to increases in Nr creation rate are most probably the ability of terrestrial ecosystems to accumulate Nr, and the fact that significant amounts of Nr added to terrestrial systems are denitrified either within the system or in the stream-river continuum prior to transport to the coast. These reasons notwithstanding, it does appear clear that with time the atmosphere is growing increasingly important in distributing Nr created by human action.




Figure 6: Nr creation in 1860, early-1990s and 2050, as a function of atmospheric emissions of $\text{NO}_x + \text{NH}_3$ and riverine Nr discharge to the coastal zone, Tg N yr^{-1} .

Nr Storage vs. Denitrification

Denitrification from terrestrial system and the stream-river-estuary continuum is estimated in a similar fashion as for the early-1990s. We estimate that in 2050, terrestrial Nr creation will be 370 Tg N. Again assuming that 25% (range 10% to 40%) is denitrified in terrestrial systems, we calculate a N_2 flux of 92 Tg N. For streams/ivers,

since 63.2 Tg N are transported to the coast, we estimate an equivalent amount produced as N₂ during transport, for a total N₂ production of 155 Tg N for continents. We further assume that all of the 63.2 Tg N that are injected into the coastal and shelf environment from continents is denitrified, primarily to N₂ which together with the 145 Tg N supplied to the shelf from the open ocean, makes total estuary/shelf denitrification 208 Tg N (Table 1, Figure 1c). Again, the uncertainties about these estimates should be stressed.

Conclusions

From 1860 to the early-1990s, anthropogenic Nr creation globally increased from ~15 Tg N yr⁻¹ to ~156 Tg N yr⁻¹. We project that by 2050 anthropogenic Nr creation will be ~270 Tg N yr⁻¹. The dispersion of Nr has kept pace with the increased creation. Total atmospheric emissions of NO_x and NH₃ increased from 23 Tg N yr⁻¹ in 1860, to 93 Tg N yr⁻¹ and 189 Tg N yr⁻¹ in the early-1990s and in 2050, respectively. The associated data for riverine Nr fluxes were 35 Tg N yr⁻¹, 59 Tg N yr⁻¹ and 75, Tg N yr⁻¹ respectively, most of which is delivered to coastal waters (Figure 6). Atmospheric NO_x and NH₃ emissions have a much more direct response to changes in Nr creation than do riverine fluxes. As Nr creation rates increase, atmospheric emissions increase much faster than does riverine discharge (Figure 5). There are some obvious reasons for the greater responsiveness of atmospheric emissions (e.g., Nr creation by fossil fuel combustion results in direct atmospheric emission), which indicate that atmospheric dispersal will become increasingly important in the future, as Nr creation rates become larger. This assumes that the degree of retention by terrestrial systems remains unchanged. There are at least two reasons why this may not be the case. First, riverine export seems likely to increase more rapidly in the future as terrestrial sinks become increasingly saturated. Secondly, the continued removal of wetland and riparian landscapes will reduce denitrification, increasing losses to rivers.

Nr creation in Africa and Latin America are dominated by natural BNF, while in Asia, North America and Europe/FSU anthropogenic process dominate. We project that in the future Nr creation rates will increase in all areas, with the largest absolute increases occurring in Asia.

The open oceans are largely decoupled from the direct impact of human alteration of the nitrogen cycle. The primary link between people and the oceans, relative to N, is through runoff and atmospheric deposition. However, with the projected increases in atmospheric deposition and runoff as described here, human induced perturbation in coastal waters will necessarily increase and the terrestrial and marine sub-components (at least in surface waters) will become more closely coupled. The effects of climate warming may also alter components of the oceanic N cycle. Increased coastal eutrophication along with a generalized warming trend will, at a minimum, likely result in the expansion of anoxic plumes.

The biggest unknown in the N cycle in managed and unmanaged ecosystems is the rate of denitrification production of N₂, and its relationship to Nr creation rates and ecosystem characteristics that control Nr cycling and storage. Until a more complete understanding of the magnitude of N losses in managed and unmanaged ecosystems is gained, determining the true, ultimate fate and long-term impact of Nr created by human actions will remain an important but unanswered scientific question.

Acknowledgements

We thank David Kicklighter, Arvin Mosier, Changsheng Li, and Knute Nadelhoffer for their comments on portions of the paper. We also thank Carolien Kroeze for her help with the N₂O budgets, Rick Jahnke for insights into deep ocean particle fluxes, and Ajit Subramaniam (UMd/ESSIC) who helped get the SST data for the BNF extrapolation. We thank Mary Ann Seifert for putting the paper into the correct format and to Sue Donovan and Bob Smith for assistance in table and figure preparation. JNG is grateful to The Ecosystems Center of the Marine Biological Laboratory, and the Woods Hole Oceanographic Institution for providing a sabbatical home to write this paper, and to the University of Virginia for the Sesquicentennial Fellowship. G. P. Asner was supported by NASA New Investigator Program (NIP) grant NAG5-8709 and NASA Interdisciplinary Science (IDS) grant NAG5-9356 and NAG5-9462. UNH contributions to this work were funded through the NASA Biological Oceanography Program (Grant # NAG5-10260), NASA Earth Observing System (NAG5-10135), Office of Naval Research (N000140110357), and the GEMS-Water Programme (UNEP/WHO/UNESCO). This paper is a contribution of the International SCOPE Nitrogen Project that received support from both the Mellon Foundation and from the National Center for Ecological Analysis and Synthesis.

REFERENCES

- Aber JD, Magill A, McNulty SG, Boone RD, Nadelhoffer KJ, Downs M, Hallett R (1995) Forest biogeochemistry and primary production altered by nitrogen saturation. *Water Air Soil Pollut.* 85: 1665-1670
- Aber JD, McDowell, WH, Nadelhoffer, KJ, Magill, A, Berntson G, Kamakea M, McNulty SG, Currie W, Rustad L & Fernandez I (1998) Nitrogen saturation in temperate forest ecosystems: hypotheses revisited. *BioScience* 48: 921-934
- Agrawal GD, Lunkad SK & Malkhed T (1999) Diffuse agricultural nitrate pollution of groundwaters in India. *Water Sci. Technol.* 39: 67-75
- Alexander RB, PJ Johnes, EW Boyer & RA Smith (2002) A comparison of methods for estimating the riverine export of nitrogen from large watersheds. *Biogeochem.* 57/58: 295-339
- Aller RC, Mackin JE, Ullman WJ, Wang CH, Tsai SM, Jin JC, Sui YN, Hong JZ. 1985. Early chemical diagenesis, sediment-water solute exchange, and storage of reactive organic-matter near the mouth of the Changjiang, East China Sea. *Continental Shelf Research* 4 (1-2): 227-251.
- Altabet MA, Francois R, Murray DW & Prell WL (1995) Climate-related variations in denitrification in the Arabian Sea from sediment 15N/14N ratios. *Nature* 373: 506-510
- Asner GP, Townsend AR, Riley WJ, Matson PA, Neff JC & Cleveland CC (2001) Physical and biogeochemical controls over terrestrial ecosystem responses to nitrogen deposition. *Biogeochem.* 54: 1-39
- Ayres RU, Schlesinger WH & Socolow RH (1994) Human impacts on the carbon and nitrogen cycles. In: Socolow RH, Andrews C, Berkhout R, & Thomas V (Eds) *Industrial Ecology and Global Change* (pp 121-155). Cambridge University Press
- Bacon PE & Freney JR (1989) Nitrogen loss from different tillage systems and the effect on cereal grain yield. *Fert. Res.* 20: 59-66
- Bange H, Rixen T, Johansen A, Siefert R, Ramesh R, Ittekkot V, Hoffmann M & Andreae M (2000) A revised nitrogen budget for the Arabian Sea. *Global Biogeochem. Cycles* 14: 1283-1297
- Bashkin VN, Park SU, Choi MS, Lee CB (2002) Nitrogen budgets for the Republic of Korea and the Yellow Sea region. *Biogeochem.* 57-58 :387-403
- Behrenfeld M & Kolber Z (1999) Widespread iron limitation in the South Pacific Ocean. *Science* 283: 840-843
- Bender M, Ganning KA, Froelich PM, Heath GR & Maynard V (1977) Interstitial nitrate profiles and oxidation of sedimentary organic matter in the eastern equatorial Atlantic. *Science* 198: 605-609
- Berman-Frank I, Cullen J, Hareli Y, Sherrell R & Falkowski PG (2001) Iron availability, cellular iron quotas, and nitrogen fixation in *Trichodesmium*. *Limnol. Oceanogr.* 46: 1249-1277
- Billen G. 1978. Budget of nitrogen recycling in North Sea sediments off Belgian Coast. *Estuarine Coastal Marine.* 7 (2): 127-146.
- Binkley D, Cromack K Jr & Baker DD (1994) Nitrogen fixation by red alder: Biology, rates, and controls. In: Hibbs DE, et al. (Eds) *The Biology and Management of Red Alder* (pp 57-72). Oregon State Univ. Press

- Blundon DJ & Dale MRT (1990) Dinitrogen fixation (acetylene reduction) in primary succession near Mount Robson, British Columbia, Canada. *Arct. Alp Res.* 22: 255-263
- Boesch DF, Brinsfield RB, Howarth RW, Baker JL, David MB, Downsing J, Fretz AT, Jaynes DB, Keeney DR, Lowarnace R, Miller K, Mitsch WJ, Nemazie D, Paerl HW, Rabalais NN, Randall GW, Scavia D, Schepters JS, Shabman L, Sharpley AN, Simposon TW, Staver KW & Townsend A (2002) Improving water quality while maintaining agricultural production. Submitted manuscript.
- Bonan GB (1996) A land surface model (LSM Version 1.0) for ecological, hydrological and atmospheric studies: Technical description and user's guide. NCAR Technical Note TN-417+STR, National Center for Atmospheric Research, Boulder, CO.
- Boring LR & Swank WT (1984) The role of black locust (*Robinia pseudo-acacia*) in forest succession. *J. Ecol.* 72: 749-766
- Bouwman AF, Boumans LJM, Batjes NH (2002) Estimation of global NH₃ volatilization loss from synthetic fertilizers and animal manure applied to arable lands and grasslands. *Global Biogeochem. Cycles* 16: 10.1029/2000GB001389.
- Bouwman AF, van der Hoek KW, Olivier JGJ (1995) Uncertainty in the global source distribution of nitrous oxide. *J. Geophys. Res.* 100: 2785-2800
- Bouwman AF, Lee DS, Asman WAH, Dentener FJ, van der Hoek KW, Olivier JGJ (1997) A global high-resolution emission inventory for ammonia. *Global Biogeochem. Cycles* 11: 561-578
- Boyer EB & Howarth RH (Eds.) (2002) *The Nitrogen Cycles at Regional to Global Scales*. Kluwer Academic Publishers
- Boyer EW, Goodale CL, Jaworski NA & Howarth RW (2002). Anthropogenic nitrogen sources and relationships to riverine nitrogen export in the northeastern USA. *Biogeochem.* 57-58: 137-169
- Boyer EW, Howarth RW, Asner G, Cleveland C, Dentener F, P Green P & Vörösmarty C (and others as warranted) Nitrogen enrichment of the coastal zone from world rivers (title may change). Manuscript in preparation
- Bradley MJ & Jones BM (2002) Reducing global NO_x emissions: promoting the development of advanced energy and transportation technologies. *Ambio* 31: 141-149
- Brandes J. A. and Devol A. H. (2002) A global marine-fixed nitrogen isotopic budget: Implications for Holocene nitrogen cycling. *Global Biogeochemical Cycles* 16(4,1120), 10.1029/2001GB001856.
- Brandes J, Devol A, Yoshinari T, Jayakumar D & Naqvi S (1998) Isotopic composition of nitrate in the central Arabian Sea and eastern tropical North Pacific: A tracer for mixing and nitrogen cycles. *Limnol. Oceanogr.* 43: 1680-1689
- Brenkert AL (Ed) (1997) CDIAC (Carbon Dioxide Information Analysis Center), Northern Hemisphere Biome and process specific forest areas and gross merchantable volumes: 1890-1990. Database DB1017 (2/1997). Oak Ridge National Laboratory, Oak Ridge
- Broecker WS & Henderson GM (1998) The sequence of events surrounding Termination II and their implications for the cause of glacial-interglacial CO₂ changes. *Paleoceanogr.* 13: 352-364

- Burkart MR & James DE (2003) Agricultural nitrogen trends in the Mississippi Basin, 1949-1997. In: Hellums D (Ed) Agricultural Production Systems of the 21st Century: Challenges and Opportunities in Plant Nutrient Management. ASA Special Publication, Madison, Wisconsin, in press
- Butterback-Bahl K, Gasche R, Willibald G & Papen, H. 2002. Exchange of N-gases at the Höglwald Forest—A summary. *Plant and Soil* 240: 117-123
- Cai GX, Zhu ZL, Trevitt ACF, Freney JR & Simpson JR (1986) Nitrogen loss from ammonium bicarbonate and urea fertilizers applied to flooded rice. *Fert. Res.* 10: 203-215
- Capone DG (1983) Benthic nitrogen fixation. In: Carpenter EJ & Capone DG (Eds) *Nitrogen in the Marine Environment* (pp 105-137). Academic Press
- Capone D (2001) Marine nitrogen fixation: what's the fuss? *Curr. Opinions Microbiol.* 4: 341-348
- Capone DG & Carpenter EJ (1999) Nitrogen fixation by marine cyanobacteria: historical and global perspectives. *Bull. Inst. Oceanogr. Monaco* 19: 235-256
- Capone DG, Subramaniam A, Montoya J, Voss M, Humborg C, Johansen A, Siefert R & Carpenter EJ (1998) An extensive bloom of the N₂-fixing cyanobacterium, *Trichodesmium erythraeum*, in the Central Arabian Sea. *Mar. Ecol. Prog. Ser.* 172: 281-292
- Capone DG, Zehr J, Paerl H, Bergman B & Carpenter EJ (1997) *Trichodesmium*: A globally significant marine cyanobacterium. *Science* 276: 1221-1229
- Capone D, JA Burns, JP Montoya, AF Michaels, A Subramaniam, EJ Carpenter (submitted) New nitrogen input to the tropical North Atlantic Ocean by nitrogen fixation by the cyanobacterium, *Trichodesmium* spp. *Science*
- Carpenter E. J. and Price C. C. (1977) Nitrogen fixation, distribution, and production of *Oscillatoria (Trichodesmium)* in the northwestern Atlantic Ocean and Caribbean Sea. *Limnology & Oceanography* 22, 60-72.
- Carpenter EJ, Subramaniam A, Capone DG (in press) Biomass and primary productivity of the cyanobacterium, *Trichodesmium* spp., in the southwestern tropical N Atlantic Ocean. *Deep-Sea Research I*
- Carpenter EJ (1983a) Nitrogen fixation by marine *Oscillatoria (Trichodesmium)* in the world's oceans. In: Carpenter EJ & Capone DG (Eds) *Nitrogen in the Marine Environment* (pp 65-103). Academic Press, Inc., New York
- Carpenter EJ (1983b) Physiology and ecology of marine *Oscillatoria (Trichodesmium)*. *Marine Biology Letters* 4: 69-85
- Carpenter EJ & Capone DG (1992) Nitrogen fixation in *Trichodesmium* blooms. In: Carpenter EJ (Ed) *Marine Pelagic Cyanobacteria: Trichodesmium and other Diazotrophs* (pp. 211-217). Kluwer Academic Publishers, Netherlands
- Carpenter E, Montoya JP, Burns J, Mulholland M, Subramaniam A & Capone DG (1999) Extensive bloom of a N₂ fixing symbiotic association in the tropical Atlantic Ocean. *Mar. Ecol. Prog. Ser.* 188: 273-283
- Cassman KG, Dobermann AD & Walters D (2002) Agroecosystems, nitrogen management and economics. *Ambio* 31: 132-140
- Chameides WL, Kasibhatla PS, Yienger J & Levy II H (1994) The growth of continental-scale metro-agro-plexes, regional ozone pollution, and world food production. *Science* 264: 74-77

- Christensen JP, Murray JW, Devol AH & Codispoti LA (1987) Denitrification in continental shelf sediments has a major impact on the oceanic nitrogen budget. *Global Biogeochem. Cycles* 1: 97-116
- Cleveland CC, Townsend AR, Schimel DS, Fisher H, Howarth RW, Hedin LO, Perakis SS, Latty EF, Von Fischer JC, Elseroad A & Wasson MF (1999) Global patterns of terrestrial biological nitrogen (N₂) fixation in natural ecosystems. *Global Biogeochem. Cycles* 13: 623-645
- Codispoti LA, Richards FA (1976) Analysis of horizontal regime of denitrification in Eastern Tropical North Pacific. *Limnology & Oceanography* 21:379-388
- Codispoti L, Brandes J, Christensen J, Devol A, Naqvi S, Paerl H & Yoshinari T (2001) The oceanic fixed nitrogen and nitrous oxide budgets: Moving targets as we enter the anthropocene? *Scientia Marina* 65(Supp. 2): 85-105
- Cornell S, Rendell A, Jickells T (1995) Atmospheric input of dissolved organic nitrogen in the oceans. *Nature* 376:243-246
- Conrad R & Dentener FJ (1999) The application of compensation point concepts in scaling of fluxes. In: Bouwman AF (Ed) *Approaches to scaling of trace gas fluxes in ecosystems* (pp. 203-216). Elsevier
- Crews TE (1999) The presence of nitrogen fixing legumes in terrestrial communities: Evolutionary vs. ecological considerations. *Biogeochem.* 46: 233-246
- Craswell ET & Martin AE (1975a) Isotopic studies of the nitrogen balance in a cracking clay. I. Recovery of added nitrogen from soil and wheat in the glasshouse and gas lysimeter. *Aust. J. Soil Res.* 13: 43-52
- Craswell ET & Martin AE (1975b) Isotopic studies of the nitrogen balance in a cracking clay. II. Recovery of nitrate 15N added to columns of packed soil and microplots growing wheat in the field. *Aust. J. Soil Res.* 13: 53-61
- Davidson EA & Ackerman IL (1993) Changes in soil carbon inventories following cultivation of previously untilled soils. *Biogeochem.* 20: 161-193
- Davidson EA & Kinglee W (1997) A global inventory of nitric oxide emissions from soils. *Nutrient Cycling in Agroecosystems* 48, 37-50.
- De Datta SK, Trevitt ACF, Freney JR, Obcemea WN, Real JG & Simpson JR (1989) Measuring nitrogen losses from lowland rice using bulk aerodynamic and nitrogen-15 balance methods. *Soil Sci. Soc. Am. J.* 53: 1275-1281
- Delwiche CC (1970) The nitrogen cycle. *Sci. Am.* 223: 137-146
- Dentener FJ & Crutzen PJ (1994) A three-dimensional model of the global ammonia cycle. *J. Atmos. Chem.* 19: 331-369
- Dentener FJ & Raes F (2002) Greenhouse gases and atmospheric chemistry: Towards integration of air pollution and climate change policies. *Third International Symposium on Non-CO₂ Greenhouse Gases (NCGG-3)*, Maastricht, the Netherlands
- Deutsch C, Gruber N, Key RM, Sarmiento JL & Ganachaud A (2001) Denitrification and N₂ fixation in the Pacific Ocean. *Global Biogeochem. Cycles* 15: 483-506
- Devol A, Codispoti L & Christensen J (1997) Summer and winter denitrification rates in western Arctic shelf sediments. *Cont. Shelf Res.* 17: 1029-1050
- Devol AH (1991) Direct measurement of nitrogen gas fluxes from continental shelf sediments. *Nature* 349: 319-322

- Dise NB & Wright RF (1995) Nitrogen leaching from European forests in relation to nitrogen deposition. *For. Ecol. Manage.* 71: 153-161
- Dise NB & Wright RF (Eds) (1992) *The NITREX Project*, Commission of the European Communities; Ecosystems Research Rep. No. 2, Brussels
- Dore JE, Brum JR, Tupas LM, & Karl DM (2002) Seasonal and interannual variability in sources of nitrogen supporting export in the oligotrophic subtropical North Pacific Ocean. *Limnol. Oceanogr.* 47: 1595-1607
- EDC (2000) Global Land Cover Characterization. Accessible through the Eros Data Center Distributed Active Archive Center at <http://edcdaac.usgs.gov/>
- Emmer E. and Thunell R. C. (2000) Nitrogen isotope variations in Santa Barbara Basin sediments: Implications for denitrification in the eastern tropical North Pacific during the last 50,000 years. *Paleoceanography* 15(4), 377-387.
- Erisman JW, de Vries W, Kros H, Oenema O, van der Eerden L, van Zeijts H & Smeulders S (2001) An outlook for a national integrated nitrogen policy. *Environ. Sci. Policy* 4: 87-95
- ESRI (1993) Digital Chart of the World. Environmental Systems Research Institute, Inc., Redlands, California
- Falkowski P (1997) Evolution of the nitrogen cycle and its influence on biological sequestration of CO₂ in the oceans. *Nature* 387: 272-273
- Falkowski P (2000) Rationalizing elemental ratios in unicellular algae. *J. Phycol.* 36: 3-6
- Falkowski P. G., Barber R. T., and Smetacek V. (1998) Biogeochemical controls and feedbacks on ocean primary production. *Science* 281, 200-206.
- FAO (2000) Fertilizer requirements in 2015 and 2030. Published by Food and Agriculture Organization of the United Nations, Rome. ISBN 92-5-104450-3, 29 pp
- FAO (2002) World agriculture: towards 2015/2040, Summary Report. Published by Food and Agriculture Organization of the United Nations, Rome. ISBN 92-5-104761-8, 97 pp
- FAOSTAT (2000) FAO Statistical Databases. Accessible through the Food and Agriculture Organization of the United Nations at <http://apps.fao.org>
- Farrell JW, Pedersen TF, Calvert SE & Nielsen B (1995) Glacial-interglacial changes in nutrient utilization in the equatorial Pacific Ocean. *Nature* 377: 514-518
- Fenn ME, Poth MA, Aber JD, Baron JS, Bormann BT, Johnson DW, Lemly AD, McNulty SG, Ryan DF & Stottlemyer R (1998) Nitrogen excess in North American ecosystems: predisposing factors, ecosystem responses and management strategies. *Ecol. Appl.* 8: 706-733
- Febre Domene L, Ayres RU (2001) Nitrogen's role in industrial systems. *J. Industrial Ecology* 5: 77-103.
- Fekete BM, Vorosmarty CJ & Grabs W (2002) High-resolution fields of global runoff combining observed river discharge and simulated water balances. *Global Biogeochem. Cy.* 16: (3) art. no. 1042.
- Fixen PE & West FB (2002) Nitrogen fertilizers • meeting the challenge. *Ambio* 31: 169-176
- Follett JR, Follett RF (2001) Utilization and metabolism of nitrogen by humans. In: Follett R, Hatfield JL (Eds) *Nitrogen in the Environment: Sources, Problems and Management* (pp 65-92). Elsevier Science B.V. Pub

- Freney JR, Chen DL, Mosier AR, Rochester IJ, Constable GA & Chalk PM (1993) Use of nitrification inhibitors to increase fertilizer nitrogen recovery and lint yield in irrigated cotton. *Fert. Res.* 34: 37-44
- Freney JR, Keerthisinghe DG, Phongpan S, Chaiwanakupt P & Harrington K (1995) Effect of urease, nitrification and algal inhibitors on ammonia loss, and grain yield of flooded rice in Thailand. *Fert. Res.* 40: 225-233
- Freney JR, Smith CJ & Mosier AR (1992) Effect of a new nitrification inhibitor (wax coated calcium carbide) on transformations and recovery of fertilizer nitrogen by irrigated wheat. *Fert. Res.* 32: 1-11
- Freney JR, Trevitt ACF, De Datta SK, Obcemea WN & Real JG (1990) The interdependence of ammonia volatilization and denitrification as nitrogen loss processes in flooded rice in the Philippines. *Biol. Fert. Soils* 9: 31-36
- Fuhrman JA & Capone DG (1991) Possible biogeochemical consequences of ocean fertilization. *Limnol. Oceanogr.* 36: 1951-1959
- Galbally IE, Freney JR, Muirhead WA, Simpson JR, Trevitt ACF & Chalk PM (1987) Emission of nitrogen oxides (NO_x) from a flooded soil fertilized with urea: Relation to other nitrogen loss processes. *J. Atmos. Chem.* 5: 343-365
- Galloway JN (1998) The global nitrogen cycle: Changes and consequences. *Environ. Pollut.* 102(S1): 15-24
- Galloway JN (2000) Nitrogen mobilization in Asia. *Nutrient Cycling in Agroecosystems* 57: 1-12
- Galloway JN & Cowling EB (2002) Reactive nitrogen and the world: Two hundred years of change. *Ambio* 31: 64-77
- Galloway JN, Aber JD, Erisman JW, Seitzinger SP, Howarth RH, Cowling EB & Cosby BJ (2003) The nitrogen cascade. *BioScience*, 53, 341-356
- Galloway JN, Cowling EB, Seitzinger SJ & Socolow R (2002) Reactive nitrogen: Too much of a good thing? *Ambio* 31: 60-63
- Galloway JN, Howarth RW, Michaels AF, Nixon SW, Prospero JM & Dentener FJ (1996) Nitrogen and phosphorus budgets of the North Atlantic Ocean and its watershed. *Biogeochem.* 35: 3-25
- Galloway JN, Schlesinger WH, Levy II H, Michaels A & Schnoor JL (1995) Nitrogen fixation: Anthropogenic enhancement--environmental response. *Global Biogeochem. Sci.* 9: 235-252
- Ganeshram RS, Pedersen TF, Calvert SE & Murray JW (1995) Large changes in oceanic nutrient inventories from glacial to interglacial periods. *Nature* 376: 755-758
- Gao Y, Kaufman J, Tanre D, Kolber D & Falkowski PG (2001) Seasonal distribution of aeolian iron fluxes to the global ocean. *Geophys. Res. Lett.* 28: 29-32
- Goering J. J., Dugdale R. C., and Menzel D. (1966) Estimates of in situ rates of nitrogen uptake by *Trichodesmium* sp. in the tropical Atlantic Ocean. *Limnol. Oceanogr.* 11, 614- 620.
- Goodale CL, Lajtha K, Nadelhoffer KJ, Boyer EA & Jaworski NA (2002) Forest nitrogen sinks in large eastern U.S. watersheds: Estimates from forest inventory and an ecosystem model. *Biogeochem.* 57/58: 239-266
- Goolsby DA, Battaglin WA, Lawrence GB, Artz RS, Aulenbach BT, Hooper RP, Keeney DR & Stensland GJ (1999) Flux and sources of nutrients in the Mississippi-Atchafalaya River Basin; Topic 3. Report for the integrated Assessment on Hypoxia

- in the Gulf of Mexico. NOAA Coastal Ocean Program Decision Analysis Series No. 17. NOAA, Silver Spring, MD
- Gorham E, Vitousek PM, & Reiners WA (1979) The regulation of chemical budgets over the course of terrestrial ecosystem succession. *Ann. Rev. Ecol. Sys.* 10: 53-84
- Green PA, Vörösmarty CJ, Meybeck M, Galloway JN, Peterson B (2003) Pre-Industrial and Contemporary Fluxes of Nitrogen through Rivers: A Global Assessment Based on Typology, in press
- Green PA, Vorosmarty CJ, Meybeck M, Galloway JN, Peterson BJ, Boyer EW. 2004. Pre-Industrial and Contemporary Fluxes of Nitrogen through Rivers: A Global Assessment Based on Typology. *Biogeochemistry*, in press
- Gruber N. (in press) The dynamics of the marine nitrogen cycle and its influence on atmospheric CO₂ variation. In *Carbon Climate Interactions* (ed. M. Follows and T. Oguz). J. Wiley & Sons.
- Gruber N & Sarmiento J (1997) Global patterns of marine nitrogen fixation and denitrification. *Global Biogeochem. Cycles* 11: 235-266
- Gunderson K., Corbin J., Hanson C., Hanson M., Hanson R., Russell D., Stollar A., and Yamada O. (1976) Structure and biological dynamics of the oligotrophic ocean photic zone off the Hawaiian Islands. *Pacific Science* **30**, 45-68.
- Gundersen P (1991) Nitrogen deposition and the forest nitrogen cycle: role of denitrification. *For. Ecol. Manage.* 44: 15-28
- Haines JR, Atlas RM, Griffiths RP, Morita RY. 1981 Denitrification and nitrogen-fixation in Alaskan continental –shelf sediments. *Applied and Environmental Microbiology.* 41 (2): 412-421.
- Hansell D, Bates N, Olson D (in press) Excess nitrate and nitrogen fixation in the North Atlantic Ocean. *Marine Chemistry*
- Haug G, Pedersen T, Sigman D, Calvert S, Nielsen B and Peterson L (1998) Glacial/interglacial variations in production and nitrogen fixation in the Cariaco Basin during the last 580K years. *Paleoceanogr.* 13: 427-3432
- Hedges JJ and Keil RG (1995) Sedimentary organic matter preservation: an assessment and speculative synthesis. *Mar. Chem.* 49: 81-115
- Hedin, L. O., J. J. Armesto, and A. H. Johnson. 1995. Patterns of nutrient loss from unpolluted, old-growth temperate forests: Evaluation of biogeochemical theory. *Ecology* 76: 493-509
- Holland EA, Dentener FJ, Brasswell BH & Sulzman JM (1999) Contemporary and pre-industrial global reactive nitrogen budgets. *Biogeochem.* 46: 7-43
- Holland EA, Braswell BH, Lamarque JF, Townsend A, Sulzman JF, Müller J-F, Dentener F, Brasseur G, Levy H II, Penner JE, Roelofs GJ (1997) Variations in the predicted spatial distribution of atmospheric nitrogen deposition and their impact on carbon uptake by terrestrial ecosystems. *J. Geophys. Res.* 102: 15,849-15,866
- Houghton RA & Hackler JL (2002) Carbon flux to the atmosphere from land-use changes. In: *Trends: A Compendium of Data on Global Change. Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, U.S. Department of Energy, Oak Ridge, Tenn., USA*
- Howarth R.W (1998) An assessment of human influences on fluxes of nitrogen from the terrestrial landscape to the estuaries and continental shelves of the North Atlantic Ocean. *Nutrient Cycling in Agroecosystems* 52(2-3): 213-223

- Howarth RW, Billen G, Swaney D, Townsend A, Jaworski N, Lajtha K, Downing JA, Elmgren R, Caraco N, Jordan T, Berendse F, Freney J, Kudeyarov V, Murdoch P & Zhu Zhao-Liang (1996) Regional nitrogen budgets and riverine N & P fluxes for the drainages to the North Atlantic Ocean: Natural and human influences. *Biogeochem.* 35: 75-139
- Howarth RW, Boyer E, Pabich W & Galloway JN (2002) Nitrogen use in the United States from 1961 - 2000, and estimates of potential future trends. *Ambio* 31: 88-96
- Howarth RW et al. (1996) Regional nitrogen budgets and riverine N & P fluxes for the drainages to the North Atlantic Ocean: Natural and human influences. *Biogeochem.* 35: 75-139
- Howarth RW, Marino R, Lane J & Cole JJ (1988) Nitrogen fixation in freshwater, estuarine, and marine ecosystems. 2. Biogeochemical controls. *Limnology and Oceanography* 33: 688-701
- Humphreys E, Freney JR, Constable GA, Smith JWB, Lilley D & Rochester IJ (1990) The Fate of Your N Fertilizer (pp 161-164). *Porch. 5th Aust. Cotton Conf.*
- IMAGE (2001) The IMAGE 2.2 implementation of the SRES scenarios. A comprehensive analysis of emissions, climate change and impacts in the 21st Century. CD-ROM publication 481508018. National Institute for Public Health and the Environment. Bilthoven, The Netherlands
- IPCC (1996) Climate change 1995: The science of climate change, contribution of working group I to the Second Assessment report of the intergovernmental panel on climate change. Houghton JT, Meira Filho LG, Callander BA, Harris N, Kattenberg A, & Maskell K (Eds) Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 572 pp
- IPCC (2001) Climate Change 2001: The Scientific Basis. Contributions of Working Group I to the Third Assessment Report of the Intergovernmental Panel on Climate Change. Houghton JT, Griggs DJ, Noguera M, van der Linden PJ, Dai X Maskell K & Johnson CA (Eds). Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 881 pp
- Joos, F., G. K. Plattner, T. F. Stocker, O. Marchal and A. Schmittner (1999) Global warming and marine carbon cycle feedbacks and future atmospheric CO₂. *Science* 284(5413): 464-467
- Johnson HB & Mayeux HS (1990) *Prosopis glandulosa* and the nitrogen balance of rangelands: extent and occurrence of nodulation. *Oecologia* 84: 176-185
- Karl DM, Letelier R, Hebel DV, Bird DF, & Winn CD (1992) *Trichodesmium* blooms and new nitrogen in the north Pacific gyre. In: Carpenter, EJ, Capone DG & Rueter JG (Eds) *Marine Pelagic Cyanobacteria: Trichodesmium and other Diazotrophs* (pp. 219-237). Kluwer Academic, Dordrecht
- Karl D (1999) A sea of change: Biogeochemical variability in the North Pacific Subtropical Gyre. *Ecosystems* 2: 181-214
- Karl DM (2002) Nutrient dynamics in the deep blue sea. *Trends Microbiol.* 10: 410-418
- Karl D, Letelier R, Tupas L, Dore J, Christian J & Hebel D (1997) The role of nitrogen fixation in biogeochemical cycling in the subtropical north Pacific ocean. *Nature* 386: 533-538

- Keerthisinghe DG, Freney JR & Mosier AR (1993) Effect of wax-coated calcium carbide and nitrapyrin on nitrogen loss and methane emission from dry-seeded flooded rice. *Biol. Fertil. Soils*. 16: 71-75
- Khalil MA & Rasmussen RA (1982) The global sources of nitrous oxide. *J. Geophys. Res.* 97: 14651-14660
- Klein Goldewijk CGM, Battjes JJ. 1997. A hundred year (1890-1990) database for integrated environmental assessments (HYDE, version 1.1)rep. 422514002, Natl. Inst. of Pub. Health and the Environ. Bilthoven, Netherlands.
- Kossinna E (1921) Die tiefen des weltmeeres. *Inst. Meersekunde, Veroff., Geogr.-naturwiss* 9, 70 pp
- Kramer DA (1999) Minerals Yearbook. Nitrogen. U.S. Geological Survey Minerals Information. <http://minerals.usgs.gov/minerals/pubs/commodity/nitrogen/>
- Koike I. and Hattori A. 1979. Estimates of denitrification in sediments of the Bering Sea shelf. *Deep Sea Research*. 26 (4): 409-415.
- Kroeze C & Seitzinger SP (1998) Nitrogen inputs to rivers, estuaries and continental shelves and related nitrous oxide emissions in 1990 and 2050: A global model. *Nutrient Cycling in Agroecosystems* 52: 195-212
- Kroeze C, Aerts R, van Breemen N, van Dam D, van der Hoek K, Hofschreuder P, Hoosbeek M, de Klein J, Kros H, van Oene H, Oenema O, Tietema A, van der Veeren R & de Vries W (2003) Uncertainties in the fate of nitrogen. I: an overview of sources of uncertainty illustrated with a Dutch case study. *Nutrient Cycling in Agroecosystems*, in press
- Kroeze C, Mosier A & Bouwman L. (1999) Closing the global N₂O budget: A retrospective analysis 1500 – 1994. *Global Biogeochem. Cycles* 13: 1-8.
- Laursen AE and Seitzinger SP. 2002. Measurement of denitrification in rivers: an integrated, whole reach approach. *Hydrobiologia*. 485 (1-3): 67-81.
- Lee K, Karl DM, Wannikhof R & Zhang J-Z (2002) Global estimate of net carbon production in the nitrate-depleted tropical and sub-tropical ocean. *Geophys. Res. Lett.* 29: 10.1029/2001GL014198
- Lelieveld J & Dentener F (2000) What controls tropospheric ozone? *J. Geophys. Res.* 105: 3531-3551
- Letelier RM & Karl DM (1996) Role of *Trichodesmium* spp in the productivity of the subtropical north Pacific ocean. *Mar. Ecol. Prog. Ser.* 133: 263-273
- Letelier RM & Karl KM (1998) *Trichodesmium* spp physiology and nutrient fluxes in the North Pacific subtropical gyre. *Aquat. Microb. Ecol.* 15: 265-276
- Levitus S, Antonov JI, Boyer TP & Stephens C (2000) Warming of the world ocean. *Science* 287(5461): 2225-2229
- Li C, Zhuang Y, Froelking S, Galloway J, Harriss R, Moore III B, Schimel D & Wang X (2003) Soil organic matter and the sustainability of Chinese agriculture: A modeling study of coupled C and N biogeochemical cycles. *Ecol. Appl.*, in press
- Longhurst A (1998) *Ecological Geography of the Sea*. Academic Press, San Diego
- Mackenzie FT (1994) Global climatic change: climatically important biogenic gases and feedbacks. In: Woodwell GM & Mackenzie FT (Eds) *Biotic Feedbacks in the Global Climatic System: Will the Warming Feed the Warming* (pp 22-46). Oxford University Press

- Mackenzie FT (1998) *Our Changing Planet: An Introduction to Earth System Science and Global Environmental Change*. 2nd Edition. Prentice-Hall USA, 486 pp
- Martinelli LA, Piccolo MC, Townsend AR, Vitousek PM, Cuevas E, McDowell W, Robertson GP, Santos OC & Treseder K (1999) Nitrogen stable isotopic composition of leaves and soil: Tropical versus temperate forests. *Biogeochem.* 46: 45-65
- Matson P, Lohse K & Hall S (2002) The globalization of nitrogen: consequences for terrestrial ecosystems. *Ambio* 31: 113-119.
- Matson PA, McDowell WH, Townsend AR & Vitousek PM (1999) The globalization of N deposition: ecosystem consequences in tropical environments. *Biogeochem.* 46: 67-83.
- Melillo JM & Cowling EB (2002) Reactive nitrogen and public policies for environmental protection. *Ambio* 31: 150-158
- Menard H & Smith S (1966) Hypsometry of ocean basin provinces. *Journal of Geophysical Research* 71: 4305-4325
- Metz B., Davidson O., Swart R., and Pan J. (2001) *Climate Change 2001: Mitigation. Contribution of Working Group III to the Third Assessment Report of the Intergovernmental Panel on Climate Change*. CUP.
- Michaels A, Karl D & Capone D (2001) Element stoichiometry, new production and nitrogen fixation. *Oceanography* 14: 68-77
- Michaels AF, Olson D, Sarmiento JL, Ammerman JW, Fanning K, Jahnke R, Knap AH, Lipschultz F & Prospero JM (1996) Inputs, losses and transformations of nitrogen and phosphorus in the pelagic North Atlantic Ocean. *Biogeochem.* 35: 181-226
- Middleburg J, Soetaert K, Herman P & Heip C (1996) Denitrification in marine sediments: A model study. *Global Biogeochem. Cycles* 10: 661-673
- Moomaw WR (2002) Energy, industry and nitrogen: strategies for reducing reactive nitrogen emissions. *Ambio* 31: 184-189
- Mosier AR, Bleken MA, Chaiwanakupt P, Ellis EC, Freney JR, Howarth RB, Matson PA, Minami K, Naylor R, Weeks K, Zhu Z-L (2001) Policy implications of human accelerated nitrogen cycling. *Biogeochem.* 52: 281-320
- Mosier AR, Guenzi WD & Schweizer EE (1986) Soil losses of dinitrogen and nitrous oxide from irrigated crops in Northeastern Colorado. *Soil Sci. Soc. Am. J.* 50: 344-348
- Mosier AR, Doran JW & Freney JR (2002) Managing soil denitrification. *J. Soil Water Conserv.* 57: 505-513
- Nadelhoffer KJ (2001) The impacts of nitrogen deposition on forest ecosystems. In: Follett RF & Hatfield JL (Eds) *Nitrogen in the Environment: Sources, Problems and Management* (pp. 311-331). Elsevier Science
- Nadelhoffer KJ, Downs MR & Fry B (1999) Sinks for ¹⁵N-enriched additions to an oak forest and a red pine plantation. *Ecol. Appl.* 9: 72-86
- Nadelhoffer KJ, Aber JD, Downs MR, Fry B & Melillo JM (1992) Biological sinks for nitrogen additions to a forested catchment. *EPA/600/A-92/292*
- Nadelhoffer KJ, Downs MR, Fry B, Aber JD, Magill AH & Melillo JM (1995) The fate of ¹⁵N-labeled nitrate additions to a northern hardwood forest in eastern Maine, USA. *Oecologia* 103: 292-301

- Naqvi SWA, Noronha RJ, Shailaja MS, Somasundar K & Sen Gupta R (1992) Some aspects of the nitrogen cycle in the Arabian Sea. In: Desai BN (Ed) *Oceanography of the Indian Ocean* (pp 285-311). Oxford and IBH
- Naqvi S., Jayakumar D., Narvekar P., Naik H., Sarma V., D'Souza W, Joseph S., and George M. (2000) Increased marine production of N₂O due to intensifying anoxia on the Indian continental shelf. *Nature* **408**, 346-349.
- National Research Council (NRC) (2000) *Clean Coastal Waters: Understanding and Reducing the Effects of Nutrient Pollution*. National Academy Press, Washington, DC
- Neff JC, Holland EA, Dentener FJ, McDowell WH & Russell KM (2002) The origin, composition and rates of organic nitrogen deposition. *Biogeochem.* 57/58: 99-136
- Nevison C, Weiss R & Erikson D (1995) Global oceanic emissions of nitrous oxide. *J. Geophys. Res.* 100: 15809-15820
- Nixon SW, Ammerman JW, Atkinson P, Berounsky VM, Billen G, Boicourt WC, Boynton WR, Church TM, Ditoro DM, Elmgren R, Garber JH, Giblin AE, Jahnke RA, Owens NJP, Pilson MEQ & Seitzinger SP (1996) The fate of nitrogen and phosphorus at the land-sea margin of the North Atlantic Ocean. *Biogeochem.* 35: 141-180
- Oenema O & Pietrzak S (2002) Nutrient management in food production: Achieving agronomic and environmental targets. *Ambio* 31: 159-168
- Orcutt KM et al. (2001) A seasonal study of the significance of N₂ fixation by *Trichodesmium* spp. at the Bermuda Atlantic Time-series Study (BATS) site. *Deep-Sea Research Part II* 48:1583-1608
- Paerl HW (1993) Emerging role of atmospheric nitrogen deposition in coastal eutrophication : biogeochemical and trophic perspectives. *Can. J. Fish. Aquat. Sci.* 50:2254-2269
- Paul EA & Clark FE (1997) *Soil Microbiology and Biochemistry*. Academic Press
- Peoples MB, Herridge DF & Ladha JK(1995) Biological nitrogen fixation: an efficient source of nitrogen for sustainable agricultural production? *Plant and Soil* 174: 3-28
- Peterjohn WT & Schlesinger WH (1991) Factors controlling denitrification in a Chihuahuan Desert ecosystem. *Soil Science Society of America Journal.* 55, 1694-1701.
- Prather M, Ehalt D, Dentener F, Derwent R, Dlugokencky E, Holland E, Isaksen, I, Katima, J, Kirchhoff V, Matson P, Midgley P, Wang M (2001) Atmospheric chemistry and greenhouse gases. In: Houghton JT, Ding Y, Griggs DJ, Noguer M, van der Linden PJ, Dai X, Maskell K & Johnson CA (Eds). *Climate Change 2001: The Scientific Basis. Contributions of Working Group I to the Third Assessment Report of the Intergovernmental Panel on Climate Change*, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 881 pp
- Preston CM & Mead DJ (1994) Growth response and recover of ¹⁵N-fertilizer one and eight growing seasons after application to lodgepole pine in British Columbia. *For. Ecol. Manage.* 65: 219-229
- Prospero JM, Barrett K, Church T, Dentener F, Duce RA, Galloway JN, Levy H, Moody J & Quinn P (1996) Atmospheric deposition of nutrients to the North Atlantic Basin. *Biogeochem.* 35: 27-73

- Quinn PK, Barrett KJ, Dentener FJ, Lipschultz F, Six KD (1996) Estimation of the air/sea exchange of ammonia for the North Atlantic basin. *Biogeochem.* 35: 275-304
- Rabalais N (2002) Nitrogen in aquatic ecosystems. *Ambio* 31: 102-112
- Rasmussen L, Beier C, Van Breemen N, De Visser P, Kreutzer K, Schierl R, Matzner E, Farrel EP (1990) Study on acid deposition effects by manipulating forest ecosystems, New Title: EXMAN—EXperimental MANipulation of forest ecosystems in Europe. Air pollution Research Report 24, Commission of the European Communities, Brussels, 42 pp
- Redfield AC (1958) The biological control of chemical factors in the environment. *Am. Sci.* 46: 205-221
- Ridgwell A. J., Maslin M. A., and Watson A. J. (2002) Reduced effectiveness of terrestrial carbon sequestration due to antagonistic response of ocean productivity. *Geophysical Research Letters* 29(6), 10.1029/2001GL014304.
- Robertson, LA, Dalsgaard, T, Revsbech, N-P, Kuenen, JG (1995) Confirmation of 'aerobic denitrification' in batch cultures, using gas chromatography and ¹⁵N mass spectrometry. *FEMS Microbiol. Ecol.* 18: 113-120
- Rolston DE, Hoffman DL & Toy DW (1978) Field measurement of denitrification: I. Flux of N₂ and N₂O. *Soil Sci. Soc. Am. J.* 42:863-869
- Rolston DE, Sharpley AM, Toy DW & Broadbent FE. 1982. Field measurement of denitrification: III. Rates during irrigation cycles. *Soil Sci. Soc. Am. J.* 46:289-296.
- Roy RN, Misra RV & Montanez A (2002) Reduced reliance on mineral nitrogen, yet more food. *Ambio* 31: 177-183
- Sachs JP and Repeta DJ (1999) Oligotrophy and nitrogen fixation during eastern Mediterranean sapropel events. *Science* 286, 2485-2488.
- Saino T. (1977) Biological nitrogen fixation in the ocean with emphasis on the nitrogen fixing blue green alga, *Trichodesmium*, and its significance in the nitrogen cycle in the low latitude sea areas. *Ph.D. Dissertation, Tokyo University.*
- Sañudo-Wilhelmy SA, Kustka AD, Gobler CJ, Hutchins DA, Yang M, Lwiza K, Burns J, Capone DG, Raven JA & Carpenter EJ (2001) Phosphorus limitation of nitrogen fixation by *Trichodesmium* in the central Atlantic Ocean. *Nature* 411: 66-69
- Sarmiento JL, Hughes TMC, Stouffer RJ & Manabe S (1998) Simulated response of the ocean carbon cycle to anthropogenic climate warming. *Nature* 393: 245-249
- Schlesinger WH (1991) *Biogeochemistry: An Analysis of Global Change.* Academic Press, 443 p
- Schlesinger WH and Hartley AE (1992). A global budget for atmospheric ammonia. *Biogeochemistry* 15, 191-211.
- Schlesinger WH and Andrews JA (2000) Soil respiration and the global carbon cycle. *Biogeochemistry* 48, 7-20.
- Seely B & Lajtha K (1997) Application of a ¹⁵N tracer to simulate and track the fate of atmospherically deposited N in the coastal forests of the Waquoit Bay Watershed, Cape Cod, Massachusetts. *Oecologia* 112: 393-402
- Seitzinger SP (1988) Denitrification in freshwater and coastal marine ecosystems: ecological and geochemical importance. *Limnology Oceanography* 33: 702-724
- Seitzinger SP & Giblin AE (1996) Estimating denitrification in North Atlantic continental shelf sediments. *Biogeochem.* 35: 235-260

- Seitzinger SP & Kroeze C (1998) Global distribution of nitrous oxide production and N inputs in freshwater and coastal marine ecosystems. *Global Biogeochem. Cycles* 12: 93-113
- Seitzinger SP, Kroeze C, Styles RV. 2000. Global Distribution of N₂O Emissions from Aquatic Systems: Natural Emissions and Anthropogenic Effects. *Chemosphere: Global Change Science* 2: 267-279
- Seitzinger SP, Styles RV, Boyer E, Alexander RB, Billen G, Howarth R, Mayer B & van Breemen N (2002) Nitrogen retention in rivers: model development and application to watersheds in the eastern U.S. *Biogeochem.* 57/58:199-237
- Shaffer G and Ronner U, 1984. Denitrification in the Baltic Proper deep-water. *Deep-Sea Research Part A- Oceanographic Research Papers* 31 (3): 197-220.
- Simpson JR & Freney JR (1988) Interacting processes in gaseous nitrogen loss from urea applied to flooded rice fields. In: Pushparajah E, Husin A, & Bachik AT (Eds) *Int. Symp. on Urea Technol. and Utilization* (pp 281-290). Malaysian Society of Soil Science; Kuala Lumpur
- Simpson JR, Freney JR., Wetselaar R., Muirhead WA, Leuning R & Denmead OT (1984) Transformations and losses of urea nitrogen after application to flooded rice. *Aust. J. Agric. Res.* 35: 189-200
- Smil V (1994) *Energy in World History*. Westview Press, Boulder, CO, 300 pp
- Smil V (1995) Who will feed China? *The China Quarterly* 143: 801-813
- Smil V (1999) Nitrogen in crop production: An account of global flows. *Global Biogeochem. Cycles* 13: 647-662
- Smil V (2000) *Feeding the World. A Challenge for the Twenty-First Century*. MIT Press, Cambridge, MA, 359 pp
- Smil V (2001) *Enriching the Earth*. MIT Press, Cambridge, MA
- Smil V (2002) Nitrogen and food. *Ambio* 31: 126-131
- Smith BD (1995) *The Emergence of Agriculture*. Scientific American Library
- Socolow R (1999) Nitrogen management and the future of food: Lessons from the management of energy and carbon. *Proc. Natl. Acad. Sci. USA*, 96: 6001-600
- Soderland R & Rosswall T (1982) The nitrogen cycles. In: Hutzinger O (Ed) *The Natural Environment and Biogeochemical Cycles* (pp 61-81). Springer Verlag
- Somasundar K, Rajendran A, Dileep Kumar M and Sen Gupta R (1990) Carbon and nitrogen budgets of the Arabian Sea. *Mar. Chem.* 30: 363-377
- Stedman DH, & Shetter R (1983) The global budget of atmospheric nitrogen species. In: Schwartz SS (Ed) *Trace Atmospheric Constituents: Properties, Transformations and Fates* (pp 411-454) John Wiley
- Steinhart GS, Likens GE, Groffman PM (2000) Denitrification in stream sediments of five northeastern (USA) streams. *Verh. Internat. Verein. Limnol.* 27: 1331-1336
- Suthhof A., Ittekkot V., and Gaye-Haake B. (2001) Millennial-scale oscillation of denitrification intensity in the Arabian Sea during the late Quaternary and its potential influence on atmospheric N₂O and global climate. *Global Biogeochemical Cycles* 15(3), 637-649.
- Sverdrup H, Johnson M & Fleming R (1942) *The Oceans*. Prentice-Hall
- Tartowski S, Howarth RW (2000) Nitrogen, nitrogen cycling. *Encyclopedia of Biodiversity* 4: 377-388

- Thamdrup B & Dalsgaard T (2002) Production of N₂ through anaerobic ammonium oxidation coupled to nitrate reduction in marine sediments. *Appl. Environ. Microbiol.* 68: 1312-1318
- Tilman D, Fargione J, Wolff B, D'Antonio C, Dobson A, Howarth R, Schindler D, Schlesinger WH, Simberloff D & Swackhamer D (2001) Forecasting agriculturally driven global change. *Science* 292: 281-284
- Townsend AR, Howarth RH, Bazzaz FA, Booth MS, Cleveland CC, Collinge SK, Dobson AP, Epstein PR, Holland EA, Keeney DR, Mallin MA, Rogers CA, Wayne P, Wolfe AH 2003. Human health effects of a changing global nitrogen cycle. *Front Ecol Environ* 1: 240–246
- van Aardenne JA, Dentener FJ, Klijn Goldewijk CGM, Lelieveld J & Olivier JGJ (2001) A 1°-1° resolution dataset of historical anthropogenic trace gas emissions for the period 1890-1990. *Global Biogeochem. Cycles* 15: 909
- van Breemen N, Boyer EW, Goodale CL, Jaworski NA, Paustian K, Seitzinger S, Lajtha LK, Mayer B, Van Dam D, Howarth RW, Nadelhoffer KJ, Eve M & Billen G (2002) Where did all the nitrogen go? Fate of nitrogen inputs to large watersheds in the northeastern USA. *Biogeochem.* 57/58: 267-293
- van Drecht G, Bouwman AF, Knoop JM, Meinardi C & Beusen A (2001) Global Pollution of Surface Waters from Point and Nonpoint Sources of Nitrogen. *The Scientific World*, 1(S2), 632-641, DOI 10.1100/tsw.2001.326
- Van Egmond ND, Bresser AHM, Bouwman AF (2002) The European nitrogen case. *Ambio* 31: 72–78
- Vitousek PM (1994) Potential nitrogen fixation during primary succession in Hawaii Volcanoes National Park. *Biotropica* 26: 234-240
- Vitousek PM, Howarth RW, Likens GE, Matson PA, Schindler D, Schlesinger WH & Tilman GD (1997) Human alteration of the global nitrogen cycle: Causes and consequences. *Issues in Ecol.* 1: 1-17
- Vitousek PM, Hattenschwiler S, Olander L & Allison S (2002) Nitrogen and nature. *Ambio* 31: 97-101
- Vörösmarty CJ, Fekete BM, Meybeck M & Lammers RB (2000) Geomorphometric attributes of the global system of rivers at 30-minute spatial resolution. *J. Hydrol.* 237: 17-39
- Wang L (1987) Soybeans--The miracle bean of China. In: Wittwer S, Yu Y, Sun H & Wang L (Eds) *Feeding a Billion: Frontiers of Chinese Agriculture*. Michigan State University Press, 462 pp
- Whelpdale, DM, Dorling SR, Hicks BB & Summers PW (1996) Atmospheric Processes. In: WMO Global Atmospheric Watch: Global Acid Deposition Assessment, 106
- Whelpdale, DM, Summers PW & Sanhueza E (1997) A global overview of atmospheric acid deposition fluxes. *Environ. Monit. Assess.* 48: 217-227
- Williams MW, Baron JS, Caine N, Sommerfeld R & Sanford R (1996) Nitrogen saturation in the Rocky Mountains. *Environ. Sci. Tech.* 30: 640-646
- Wilson T. 1978. Evidence for denitrification in aerobic pelagic sediments. *Nature.* 274: 354-356.
- Wittwer S, Yu Y, Sun H, & Wang L (1987) *Feeding a Billion: Frontiers of Chinese Agriculture*. Michigan State University Press, 462 pp

- Wolfe A, & Patz JA (2002) Nitrogen and human health: direct and indirect impacts. *Ambio* 31: 120-125
- Wright RF & Rasmussen L (1998) Introduction to NITREX and EXMAN projects. *For. Ecol. Manage.* 101: 1-7
- Wu J, Sunda W, Boyle E and Karl D (2000) Phosphate depletion in the western North Atlantic Ocean. *Science* 289: 759-762
- Xing GX & Zhu ZL (2002) Regional nitrogen budgets for China and its major watersheds. *Biogeochem.* 57-58: 405-427
- Yavitt, JB and Fahey TJ (1993) Production of methane and nitrous oxide by organic soils within a northern hardwood forest ecosystem. Pp. 261-277. In *Biogeochemistry of Global Change*. R. S. Oremland (ed). Chapman and Hall, New York.
- Zehr, J. and B.B. Ward. 2002. Nitrogen cycling in the ocean. New perspective on processes and paradigms. *Applied and Environmental Microbiology* 68(3):1015-1024.
- Zehr J, Waterbury J, Turner P, Montoya J, Omoregie E, Steward G, Hansen A & Karl D (2001) Unicellular cyanobacteria fix N₂ in the subtropical North Pacific Ocean. *Nature* 412: 635-638
- Zehr JP, Carpenter EJ & Villareal TA (2000) New perspectives on nitrogen-fixing microorganisms in subtropical and tropical open oceans. *Trends in Microbiol.* 8: 68-73
- Zhang WL, Tian ZX, Zhang N & Li XQ (1996) Nitrate pollution of groundwaters in northern China. *Agric. Ecosyst. & Environ.* 59: 223-231
- Zheng X, Fu C, Xu X, Yan X, Chen G, Han S, Huang Y & Hu F (2002) The Asian nitrogen case. *Ambio* 31: 79-87
- Zhu ZL, Cai GX, Simpson JR, Zhang SL, Chen DL, Jackson AV & Freney JR (1989) Processes of nitrogen loss from fertilizers applied to flooded rice fields on a calcareous soil in North-Central China. *Fert. Res.* 18: 101-115
- Zobell C & Anderson DQ (1943) Observations on the multiplication of bacteria in different volumes of stored sea water and the influence of oxygen tension and solid surfaces. *J. Bacteriol.* 46: 324-342.

Appendix I

Uncertainties in the modeling of reduced and oxidized nitrogen deposition fluxes.

The main uncertainties in the modeling of N deposition fluxes are:

- Emission inventories
- Chemical transformations
- Wet and dry removal processes
- Atmospheric transport and resolution of the model.

The emission inventories for NO and NH₃ used in this study are based on the widely used EDGAR2.0 database for NO and the Bouwman et al. [1997] compilation for NH₃. The inventories provide data on 1x1 degree which were aggregated in the model resolution.

The inaccuracies of the NO emissions are of the order of 30% in the industrialized regions of N. America, Europe and Japan. In other regions a subjective estimate is of the order 50%. Due to cancellation of errors the uncertainties are smallest for larger regions, and become larger on smaller temporal and spatial scales. Natural emissions (e.g. lightning and soils). A typical global uncertainty of NO_x from lightning is 2-10 Tg N yr⁻¹. For soils 2-20 Tg N yr⁻¹.

The NH₃ emissions are more uncertain, since essentially European emission factors were used -with some corrections- in other regions. Also the activity data (e.g. amount of animals) is subject to large fluctuations in many countries (since it is market driven). The seasonal emissions are even more uncertain- e.g. for rice paddy emissions. Overall an uncertainty of 50% is estimated. Natural emissions- from soils, vegetation and oceans are poorly known with a large uncertainty range.

There are still large uncertainties in the chemistry of NO_x. However the influence of these uncertainties on oxidized nitrogen deposition is probably not so large. The distribution between e.g. nitrate and other forms of organic oxidized nitrogen deposition is an issue- but the influence on over-all oxidized nitrogen deposition should be limited.

The transformation of NH₃ into ammonium and subsequent deposition is relatively straightforward since it essentially reacts with sulfate and nitrate aerosol in the atmosphere. Some uncertainties arise from inadequate treatment of sulfur chemistry (e.g. underestimate of sulfate aerosol formation in winter) and sub-grid phenomena in ammonium nitrate formation. The efficiency of transformation is closely linked to the dry deposition of NH₃, which is much faster than the dry deposition of ammonium aerosol.

Whereas wet deposition in models can be tested with some independent variables (e.g. rainfall can be compared to measurement, ²¹⁰Pb is a radionuclide which behaves as aerosol. However there are very few possibilities to test the dry deposition parameterizations in the model. The latter is very important for the removal of gas phase HNO₃ and NH₃. It is likely that the aggregation of dry plus wet deposition in models leads to cancellation of errors, especially on larger spatial scales. E.g. an underestimate of wet deposition automatically to higher dry deposition. This results in a locally higher N deposition flux, and less N deposition away from the source. In a larger region the sum is balancing.

The underlying meteorological data are calculated by a 0.5-0.5 weather forecast model, which data were interpolated to the model resolution, except in the tropics these

|

data are probably not a major source of errors. The transport scheme ('slopes') is designed to correctly represent strong gradients, such as seen in emission regions. However, one can still expect problems in e.g. coastal regions, where the model resolution is not high enough to capture the land-sea gradients. A problem is that with a coarse resolution model tracers are emitted and partly deposited in a grid box that contains both land as well as sea surface.

An overall impression of the accuracy of the model comes from comparison with measurements. Except in Europe and North America there are not many measurements available on wet deposition and (surface) aerosol concentrations. In general the model depositions are indeed within plus/minus 50% of the measurements- and many times better.

Appendix II The NTNI Model of Riverine Nr Transport

In their analysis of the flux of Nr from large regions to the North Atlantic Ocean, Howarth et al. (1996) put forth an empirical model relating net anthropogenic Nr inputs per landscape area (NANI) to the total flux of Nr discharged in rivers to the coastal zone, yielding a strong positive linear relationship between the two. The approach considers new inputs of Nr to a region that are human controlled, including inputs from fossil-fuel derived atmospheric deposition, fixation in cultivated croplands, fertilizer use, and the net import (or export) in food and feed. Subsequent studies have found that the form of the relationship holds when considering other temperate regions of the world over multiple scales (e.g., Boyer et al. 2002, Boyer et al. in prep). Further, a model intercomparison by Alexander et al. (2002) found the NANI model to be the most robust and least biased of the models used to estimate N fluxes from a variety of large temperate watersheds.

In a paper related to this effort, Boyer et al. (in prep.) describe that the relationship between anthropogenic Nr inputs and outputs has not held in tropical regions of the world and other areas where natural Nr inputs are substantial (Howarth et al. 1996, Boyer et al., in prep.). Addressing this issue, they put forth a modification to the NANI model, considering new inputs of Nr to a region from natural BNF in forests and other non-cultivated vegetated lands in addition to anthropogenic Nr inputs. This model relates net total nitrogen inputs per unit area of landscape (or NTNI, which includes anthropogenic plus natural Nr inputs) to riverine exports, and has been shown to hold over multiple spatial scales in contrasting ecoregions of the world including temperate and tropical areas. Using data for a variety of coastal watersheds throughout the world, riverine export was approximately 25% of the net total nitrogen inputs per unit area of landscape.

Aggregation of Nr input data for each region by Boyer et al. was based on the same data sources discussed in this paper, mapped spatially using GIS. They obtained modeled estimates of total (wet + dry) atmospheric deposition of $\text{NO}_y\text{-N}$ from fossil fuel combustion from the global chemistry transport model (TM3) of the University of Utrecht (F. Dentener pers. comm.). This model has been widely used and validated extensively for N species (e.g., Dentener and Crutzen 1994; Prospero et al. 1996; Holland et al. 1999). We used TM3 model simulations for 1860, 1990, and 2050 (F. Dentener pers. comm.). To quantify net imports of N in food and feed, we used the continental values presented in this paper for 1860 and 1990, and disaggregated to the scale of water regions based on their fraction of area within each continent. We assumed that net food and feed exports would stay the same between 1990 and 2050, which is known to be a conservative estimate regardless of future population and diet scenarios (Howarth et al. 2002; Galloway et al. 2002).

To describe natural BNF in forests and other non-cultivated vegetated lands of the world, Boyer et al. used modelled estimates put forth by Cleveland et al. (1999) and modified by Cleveland and Asner (pers. comm.). Their modified model is based on estimates of plant N requirement simulated with the TerraFlux biophysical-biogeochemical process model to constrain estimates of BNF in vegetation across biomes of the world (Asner et al., 2001; Bonan, 1996).. Fixation rates encompassed in the model are based on a synthesis of rates reported in the literature. (G. Asner and C. Cleveland,

pers. comm.). We used simulations for 1860 and for 1990, where cultivated areas of the landscape under human control were excluded. To quantify BNF due to human cultivation of crops, we used the continental values presented in this paper for 1860 and 1990, and disaggregated to the scale of water regions based on vegetated area within each continent. We assumed that both natural and cultivated BNF rates will stay at 1990 levels in 2050, which we consider to be a conservative assumption given the increase in world population and the associated need to increase food production.

No commercial fertilizers were used in 1860, but to describe the pattern of N fertilizer use in 1990, Boyer et al. used a gridded map of world fertilizers prepared by the Water Systems Analysis Group, UNH (Green et al. 2003) which is based on FAO data. Country-level nitrogenous fertilizer consumption totals for 1995 taken from the FAOSTAT Statistical Databases (FAOSTAT 2000) were evenly distributed among a 1 km resolution cropland dataset derived from the 2000 EDC global land cover dataset (EDC 2000) and resampled to the 30-minute resolution to create this dataset. To extrapolate a future fertilizer use scenario, we used the projected fertilizer trends for world regions presented by FAO (2000) in their analysis based on scenarios of population growth and human behavior. Taking their “baseline” scenario for N fertilizer use, we used the data presented for each world region to calculate change per year between 1990 and 2030, and assumed this rate of increase from 2030 until 2050. We assumed that fertilizers were applied only to agricultural lands within each world region, and disaggregated the 2050 fertilizer use estimates from the scale of the world regions to the scale of the water regions within them based on the fraction of agricultural area in each watershed.

Boyer et al. calculated riverine N_r exports to both inland receiving waters and coastal waters. Estimates of riverine flux to the coastal zone exclude the N_r fluxes to inland catchments that do not drain to the coasts and inactive areas of the landscape that do not transmit water to the coast due to insufficient surface water runoff. Inactive areas were determined using potential river flow paths and a threshold of 3 mm/yr representing the minimum upstream runoff required to sustain perennial discharge in river channels. The inland and inactive areas are mapped according to a global scale watershed delineation (Vörösmarty, et al. 2000) in conjunction with a global composite runoff dataset (Fekete, et al 2002).

Table Legends

1. Global Nr creation and distribution, Tg N yr⁻¹
2. Global atmospheric emissions of (a) NH₃ and NO_x, Tg N yr⁻¹ and (b) N₂O, Tg N yr⁻¹
3. Nr creation by region, Tg N yr⁻¹
4. Cultivation-induced Nr creation by region, Tg N yr⁻¹ (1995)
5. Exchange of N (import and export) in commodities among regions, Tg N yr⁻¹
6. Regional atmospheric emissions and deposition for the early-1990s, Tg N yr⁻¹
7. Average areal rate of marine BNF. Values are means from indicated studies. Standard errors of the means are given for the individual studies, where available. For recent cruise based studies of Capone et al., stations are chosen arbitrarily along a cruise track at 0600 each morning.
8. BNF and denitrification by ocean basin
9. Some estimates of marine sedimentary denitrification

Figure Legends

1. Components of the global nitrogen cycle for (a) 1860, (b) early-1990s and (c) 2050, Tg N yr^{-1} . All colored boxes represent reservoirs of nitrogen species in the atmosphere.. Creation of Nr is depicted with bold arrows from the N_2 reservoir to the Nr reservoir (depicted by the dotted box). ‘N-BNF’ is biological nitrogen fixation within natural ecosystems; ‘C-BNF’ is biological nitrogen fixation within agroecosystems. Denitrification creation of N_2 from Nr within the dotted box is also shown with bold arrows. All arrows that do not leave the dotted box represent inter-reservoir exchanges of Nr . The dashed lines within the dotted box associated with NH_x represent natural emissions of NH_3 that are re-deposited on fast time-scales to the oceans and continents.
2. Spatial patterns of total inorganic nitrogen deposition in (a) 1860, (b) early-1990s and (c) 2050, $\text{mg N m}^{-2} \text{ yr}^{-1}$.
3. Nr creation and distribution by geopolitical region for the early-1990s, Tg N yr^{-1} .
4. Nr creation and distribution by ocean basin, Tg N yr^{-1} in the early-1990s
5. Riverine Nr export to the coastal zone (Tg N yr^{-1}) in the past (1860), present (1990) and future (2050). Dry and inland watershed regions that do not transmit to coastal areas are shown in gray.
6. Nr creation in 1860, early-1990s and 2050, as a function of atmospheric emissions of $\text{NO}_x + \text{NH}_3$ and riverine Nr discharge to the coastal zone, Tg N yr^{-1} .