

**Science Advisory Board Panel Draft Report (dated April 8, 2011) - Do not Cite or Quote**

This draft is a work in progress, does not reflect consensus advice or recommendations, has not been approved by the chartered SAB, and does not represent EPA policy.

1  
2  
3  
4  
5  
6  
7  
8  
9

10 EPA-SAB-11-xxx

11  
12 The Honorable Lisa P. Jackson  
13 Administrator  
14 U.S. Environmental Protection Agency  
15 1200 Pennsylvania Avenue, N.W.  
16 Washington, D.C. 20460

17  
18  
19  
20  
21

Subject: Review of EPA's draft Approaches for Deriving Numeric Nutrient  
Criteria for Florida's Estuaries, Coastal Waters, and Southern Inland  
Flowing Waters

22 Dear Administrator Jackson:

23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34

Nitrogen (N) and phosphorus (P) inputs from urban and agricultural sources are known to impact water quality, and nutrient pollution has been identified as a source of impairment for estuarine, marine and fresh waters in Florida. In 2009, EPA determined that numeric nutrient criteria were needed to protect aquatic life in Florida, and initiated a process to develop such criteria for categories of state waters. The SAB was asked to review and provide advice on the proposed approaches for estuarine, coastal and South Florida waters, as described in the draft EPA document, *Methods and Approaches for Deriving Numeric Criteria for Nitrogen/Phosphorus Pollution in Florida's Estuaries, Coastal Waters, and Southern Inland Flowing Waters*. An *ad hoc* SAB panel, the Nutrient Criteria Review Panel, was formed for this task.

35  
36  
37  
38  
39  
40  
41

The Charge to the Panel included questions about the conceptual model used to select assessment endpoints, data sources, and possible approaches to define criteria for each of the categories of waters: estuaries, coastal waters, inland flowing waters (including canals) in South Florida, and South Florida marine waters. The Panel held a public review meeting on December 13-14, 2010 and a follow-up public teleconference on February 7, 2011. For both meetings, the Panel received numerous technical comments from Florida state and local agencies and interested members of the public.

42  
43  
44  
45  
46

Nitrogen and phosphorus may be limiting in different portions of the fresh-to-marine continuum, and in some cases may be co-limiting. Thus, a dual nutrient (N and P) strategy is warranted, and we agree with EPA's decision to take this approach. The Panel acknowledges the substantial effort that already has been made by EPA to get the work to this stage, much of it

**Science Advisory Board Panel Draft Report (dated April 8, 2011) - Do not Cite or Quote**

This draft is a work in progress, does not reflect consensus advice or recommendations, has not been approved by the chartered SAB, and does not represent EPA policy.

1 solid and thoughtful. However, much work remains to be done. Under a court-ordered consent  
2 decree, the Agency has committed to proposing nutrient criteria by November 14, 2011 and final  
3 criteria by August 14, 2012. Given this time frame and the extent of the task at hand, the Panel  
4 is concerned that the Agency not sacrifice quality work for the sake of a schedule.

5  
6 To guide its development of nutrient criteria, EPA proposes a conceptual model that links  
7 nitrogen and phosphorus levels in Florida waters to biological endpoints to be protected using  
8 one or more analytical approaches. Although the general conceptual model provides a starting  
9 point for choosing numeric criteria, the Panel has numerous concerns about how the driver  
10 variables—total nitrogen (TN) and total phosphorus (TP)—would be linked to measurable  
11 biological endpoints.

12  
13 EPA is proposing three general approaches to relate nutrient levels to balanced natural  
14 populations: (1) reference conditions; (2) predictive stressor-response relationships; and/or (3)  
15 numerical water quality models. The Panel had some concern with the uneven treatment of the 3  
16 approaches (i.e., the emphasis on water quality modeling), and encourages EPA to continue to  
17 develop all three. The Panel agrees that these approaches all have utility and that using a  
18 combination may be best where data and models are available. However, EPA should provide  
19 more detail on the adequacy of the data for applying each approach; how decisions would be  
20 made on which approaches to use; and how discrepancies in targets derived from different  
21 approaches would be resolved.

22  
23 The proposed biological endpoints (healthy seagrasses, balanced phytoplankton biomass,  
24 and balanced faunal communities) are appropriate, but EPA needs to define “balanced” for each  
25 of these endpoints, preferably in quantitative terms. The Panel agrees with EPA’s broad  
26 delineation of Florida coastal waters into four categories (estuaries, coastal waters, South Florida  
27 inland waters, and South Florida estuarine and coastal waters) for purposes of criteria  
28 development, but suggests some refinements to segmentation within the categories. EPA also  
29 proposes to use downstream protection value (DPV) criteria to ensure that upstream nutrient  
30 criteria will be set at levels that will protect downstream estuaries. While agreeing with the goal,  
31 the Panel has concern with the overlap between DPV and the Total Maximum Daily Load  
32 (TMDL) process. The Panel reached the following broad conclusions:

- 33  
34
- 35 • For estuaries, the Panel recommends that EPA adopt additional measures of seagrass  
36 health beyond the proposed use of chlorophyll-*a*, and encourages the use of direct  
37 measures of the faunal communities to be protected, rather than relying on a dissolved  
38 oxygen criterion.
  - 39 • For coastal waters, a criterion based on satellite-derived estimates of chlorophyll may be  
40 the only feasible approach for this large, poorly sampled region. However, the Panel is  
41 concerned that no direct measurements of nutrients are proposed to verify the strength of  
42 the relationship between pollutant loads from land and observed Chl-*a* concentrations.
  - 43 • For South Florida inland waters, the Panel was not convinced that nutrient criteria based  
44 on instream protection values were meaningful for man-made and managed canals, but  
45 does agree that nutrients in canal waters should be managed to ensure downstream,  
estuarine designated uses.

**Science Advisory Board Panel Draft Report (dated April 8, 2011) - Do not Cite or Quote**

This draft is a work in progress, does not reflect consensus advice or recommendations, has not been approved by the chartered SAB, and does not represent EPA policy.

- For South Florida coastal and estuarine waters, the Panel recommends that seagrass endpoints be considered in addition to chlorophyll-*a*.
- If the DPV approach is pursued, the Panel urges that apportionment strategies not preclude cross-tributary nutrient trading or other approaches to achieving the necessary load reductions.

In closing, the SAB encourages EPA to continue efforts to develop numeric nutrient criteria for Florida estuarine and coastal waters, using the best available scientific data and methods. We appreciate the opportunity to provide advice on this important endeavor, and look forward to your response.

Sincerely,

Dr. Deborah L. Swackhamer, Chair  
Science Advisory Board

Dr. Judith L. Meyer, Chair  
SAB Nutrient Criteria Review Panel

Enclosure

**Science Advisory Board Panel Draft Report (dated April 8, 2011) - Do not Cite or Quote**

This draft is a work in progress, does not reflect consensus advice or recommendations, has not been approved by the chartered SAB, and does not represent EPA policy.

**NOTICE**

1  
2  
3 This report has been written as part of the activities of the EPA Science Advisory Board (SAB),  
4 a public advisory group providing extramural scientific information and advice to the  
5 Administrator and other officials of the Environmental Protection Agency. The SAB is  
6 structured to provide balanced, expert assessment of scientific matters related to problems facing  
7 the Agency. This report has not been reviewed for approval by the Agency and, hence, the  
8 contents of this report do not necessarily represent the views and policies of the Environmental  
9 Protection Agency, nor of other agencies in the Executive Branch of the Federal government, nor  
10 does mention of trade names of commercial products constitute a recommendation for use.  
11 Reports of the SAB are posted on the EPA Web site at <http://www.epa.gov/sab>.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45

**U.S. Environmental Protection Agency  
Science Advisory Board  
Nutrient Criteria Review Panel**

**CHAIR**

**Dr. Judith L. Meyer**, Professor Emeritus, Odum School of Ecology, University of Georgia, Lopez Island, WA

**MEMBERS**

**Dr. Walter Boynton**, Professor, Chesapeake Biological Laboratory, Center for Environmental Science, University of Maryland, Solomons, MD

**Dr. Deborah Bronk**, Professor, Physical Science Dept, The College of William and Mary, Virginia Institute of Marine Science, Gloucester Point, VA

**Dr. Piers Chapman**, Professor and Head, Department of Oceanography, Texas A&M University, College Station, TX

**Dr. Robert Diaz**, Professor, Department of Biological Sciences, Virginia Institute of Marine Science, College of William and Mary, Gloucester Pt., VA

**Dr. Anne Giblin**, Senior Scientist, The Ecosystems Center, Marine Biological Laboratory, Woods Hole, MA, United States

**Dr. Kenneth L. Heck, Jr**, Chair, University Programs/Senior Marine Scientist III/Professor, Dauphin Island Sea Lab, Dauphin Island, AL

**Dr. Mark Noll**, Associate Professor, Earth Sciences & Special Assistant to the Provost, Academic Affairs, The College at Brockport, State University of New York, Brockport, NY

**Dr. Peter Ortner**, Research Professor, Marine Biology and Fisheries, Rosenstiel School of Marine and Atmospheric Science, University of Miami, Miami, FL

**Dr. Hans Paerl**, Professor of Marine and Environmental Sciences, Institute of Marine Sciences, University of North Carolina - Chapel Hill, Morehead City, NC

**Dr. Kenneth Reckhow**, Chief Scientist, Global Climate Change and Environmental Sciences, RTI International, Research Triangle Park, NC

**Dr. James Sanders**, Director and Professor, Skidaway Institute of Oceanography, Savannah, GA

**Science Advisory Board Panel Draft Report (dated April 8, 2011) - Do not Cite or Quote**

This draft is a work in progress, does not reflect consensus advice or recommendations, has not been approved by the chartered SAB, and does not represent EPA policy.

1 **Dr. David C. Schneider**, Professor, Ocean Sciences Centre, Memorial University, St. John's,  
2 NL, Canada

3  
4 **Dr. Andrew N. Sharpley**, Research Soil Scientist, Department of Crop, Soil and Environmental  
5 Sciences, University of Arkansas, Fayetteville, AR, U.S.A.

6  
7 **Dr. Andrew Solow**, Associate Scientist, and Center Director, Woods Hole Oceanographic  
8 Institution, Woods Hole, MA

9  
10 **Dr. Alan Steinman**, Director, Annis Water Resources Institute, Grand Valley State University,  
11 Muskegon, MI

12  
13 **Dr. Joseph C. Zieman**, Professor, Environmental Sciences, College and Graduate School for  
14 Arts and Sciences, University of Virginia, Charlottesville, VA

15  
16  
17  
18 **SCIENCE ADVISORY BOARD STAFF**

19 **Ms. Stephanie Sanzone**, Designated Federal Officer, U.S. Environmental Protection Agency,  
20 Science Advisory Board (1400R), 1200 Pennsylvania Avenue, NW, Washington, DC, Phone:  
21 202-564-2067, Fax: 202-565-2098, ([sanzone.stephanie@epa.gov](mailto:sanzone.stephanie@epa.gov))  
22

1 **U.S. Environmental Protection Agency**  
2 **Science Advisory Board**  
3  
4

5 **CHAIR**

6 **Dr. Deborah L. Swackhamer**, Professor and Charles M. Denny, Jr., Chair in Science,  
7 Technology and Public Policy- Hubert H. Humphrey School of Public Affairs and Co-Director  
8 of the Water Resources Center, University of Minnesota, St. Paul, MN  
9

10  
11 **MEMBERS**

12 **Dr. David T. Allen**, Professor, Department of Chemical Engineering, University of Texas,  
13 Austin, TX  
14

15 **Dr. Claudia Benitez-Nelson**, Full Professor and Director of the Marine Science Program,  
16 Department of Earth and Ocean Sciences , University of South Carolina, Columbia, SC  
17

18 **Dr. Timothy Buckley**, Associate Professor and Chair, Division of Environmental Health  
19 Sciences, College of Public Health, The Ohio State University, Columbus, OH  
20

21 **Dr. Patricia Buffler**, Professor of Epidemiology and Dean Emerita, Department of  
22 Epidemiology, School of Public Health, University of California, Berkeley, CA  
23

24 **Dr. Ingrid Burke**, Director, Haub School and Ruckelshaus Institute of Environment and Natural  
25 Resources, University of Wyoming, Laramie, WY  
26

27 **Dr. Thomas Burke**, Professor, Department of Health Policy and Management, Johns Hopkins  
28 Bloomberg School of Public Health, Johns Hopkins University, Baltimore, MD  
29

30 **Dr. Terry Daniel**, Professor of Psychology and Natural Resources, Department of Psychology,  
31 School of Natural Resources, University of Arizona, Tucson, AZ  
32

33 **Dr. George Daston**, Victor Mills Society Research Fellow, Product Safety and Regulatory  
34 Affairs, Procter & Gamble, Cincinnati, OH  
35

36 **Dr. Costel Denson**, Managing Member, Costech Technologies, LLC, Newark, DE  
37

38 **Dr. Otto C. Doering III**, Professor, Department of Agricultural Economics, Purdue University,  
39 W. Lafayette, IN  
40

41 **Dr. David A. Dzombak**, Walter J. Blenko Sr. Professor of Environmental Engineering ,  
42 Department of Civil and Environmental Engineering, College of Engineering, Carnegie Mellon  
43 University, Pittsburgh, PA  
44

**Science Advisory Board Panel Draft Report (dated April 8, 2011) - Do not Cite or Quote**

This draft is a work in progress, does not reflect consensus advice or recommendations, has not been approved by the chartered SAB, and does not represent EPA policy.

1 **Dr. T. Taylor Eighmy**, Vice President for Research, Office of the Vice President for Research,  
2 Texas Tech University, Lubbock, TX

3

4 **Dr. Elaine Faustman**, Professor and Director, Institute for Risk Analysis and Risk  
5 Communication, School of Public Health, University of Washington, Seattle, WA

6

7 **Dr. John P. Giesy**, Professor and Canada Research Chair, Veterinary Biomedical Sciences and  
8 Toxicology Centre, University of Saskatchewan, Saskatoon, Saskatchewan, Canada

9

10 **Dr. Jeffrey K. Griffiths**, Associate Professor, Department of Public Health and Community  
11 Medicine, School of Medicine, Tufts University, Boston, MA

12

13 **Dr. James K. Hammitt**, Professor, Center for Risk Analysis, Harvard University, Boston, MA

14

15 **Dr. Bernd Kahn**, Professor Emeritus and Associate Director, Environmental Radiation Center,  
16 Georgia Institute of Technology, Atlanta, GA

17

18 **Dr. Agnes Kane**, Professor and Chair, Department of Pathology and Laboratory Medicine,  
19 Brown University, Providence, RI

20

21 **Dr. Madhu Khanna**, Professor, Department of Agricultural and Consumer Economics,  
22 University of Illinois at Urbana-Champaign, Urbana, IL

23

24 **Dr. Nancy K. Kim**, Senior Executive, Health Research, Inc., Troy, NY

25

26 **Dr. Catherine Kling**, Professor, Department of Economics, Iowa State University, Ames, IA

27

28 **Dr. Kai Lee**, Program Officer, Conservation and Science Program, David & Lucile Packard  
29 Foundation, Los Altos, CA

30

31 **Dr. Cecil Lue-Hing**, President, Cecil Lue-Hing & Assoc. Inc., Burr Ridge, IL

32

33 **Dr. Floyd Malveaux**, Executive Director, Merck Childhood Asthma Network, Inc., Washington,  
34 DC

35

36 **Dr. Lee D. McMullen**, Water Resources Practice Leader, Snyder & Associates, Inc., Ankeny,  
37 IA

38

39 **Dr. Judith L. Meyer**, Professor Emeritus, Odum School of Ecology, University of Georgia,  
40 Lopez Island, WA

41

42 **Dr. James R. Mihelcic**, Professor, Civil and Environmental Engineering, University of South  
43 Florida, Tampa, FL

44

45 **Dr. Jana Milford**, Professor, Department of Mechanical Engineering, University of Colorado,  
46 Boulder, CO

**Science Advisory Board Panel Draft Report (dated April 8, 2011) - Do not Cite or Quote**

This draft is a work in progress, does not reflect consensus advice or recommendations, has not been approved by the chartered SAB, and does not represent EPA policy.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46

**Dr. Christine Moe**, Eugene J. Gangarosa Professor, Hubert Department of Global Health, Rollins School of Public Health, Emory University, Atlanta, GA

**Dr. Horace Moo-Young**, Dean and Professor, College of Engineering, Computer Science, and Technology, California State University, Los Angeles, CA

**Dr. Eileen Murphy**, Grants Facilitator, Ernest Mario School of Pharmacy, Rutgers University, Piscataway, NJ

**Dr. Duncan Patten**, Research Professor, Hydroecology Research Program, Department of Land Resources and Environmental Sciences, Montana State University, Bozeman, MT

**Dr. Stephen Polasky**, Fesler-Lampert Professor of Ecological/Environmental Economics, Department of Applied Economics, University of Minnesota, St. Paul, MN

**Dr. Arden Pope**, Professor, Department of Economics, Brigham Young University, Provo, UT

**Dr. Stephen M. Roberts**, Professor, Department of Physiological Sciences, Director, Center for Environmental and Human Toxicology, University of Florida, Gainesville, FL

**Dr. Amanda Rodewald**, Professor of Wildlife Ecology, School of Environment and Natural Resources, The Ohio State University, Columbus, OH

**Dr. Jonathan M. Samet**, Professor and Flora L. Thornton Chair, Department of Preventive Medicine, University of Southern California, Los Angeles, CA

**Dr. James Sanders**, Director and Professor, Skidaway Institute of Oceanography, Savannah, GA

**Dr. Jerald Schnoor**, Allen S. Henry Chair Professor, Department of Civil and Environmental Engineering, Co-Director, Center for Global and Regional Environmental Research, University of Iowa, Iowa City, IA

**Dr. Kathleen Segerson**, Philip E. Austin Professor of Economics, Department of Economics, University of Connecticut, Storrs, CT

**Dr. Herman Taylor**, Director, Principal Investigator, Jackson Heart Study, University of Mississippi Medical Center, Jackson, MS

**Dr. Barton H. (Buzz) Thompson, Jr.**, Robert E. Paradise Professor of Natural Resources Law at the Stanford Law School and Perry L. McCarty Director, Woods Institute for the Environment, Stanford University, Stanford, CA

**Dr. Paige Tolbert**, Professor and Chair, Department of Environmental Health, Rollins School of Public Health, Emory University, Atlanta, GA

**Science Advisory Board Panel Draft Report (dated April 8, 2011) - Do not Cite or Quote**

This draft is a work in progress, does not reflect consensus advice or recommendations, has not been approved by the chartered SAB, and does not represent EPA policy.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18

**Dr. John Vena**, Professor and Department Head, Department of Epidemiology and Biostatistics, College of Public Health, University of Georgia, Athens, GA

**Dr. Thomas S. Wallsten**, Professor and Chair, Department of Psychology, University of Maryland, College Park, MD

**Dr. Robert Watts**, Professor of Mechanical Engineering Emeritus, Tulane University, Annapolis, MD

**Dr. R. Thomas Zoeller**, Professor, Department of Biology, University of Massachusetts, Amherst, MA

**SCIENCE ADVISORY BOARD STAFF**

**Dr. Angela Nugent**, Designated Federal Officer, U.S. Environmental Protection Agency, Science Advisory Board (1400R), 1200 Pennsylvania Avenue, NW, Washington, DC, Phone: 202-564-2218, Fax: 202-565-2098, (nugent.angela@epa.gov)

**Science Advisory Board Panel Draft Report (dated April 8, 2011) - Do not Cite or Quote**  
This draft is a work in progress, does not reflect consensus advice or recommendations, has not been approved by  
the chartered SAB, and does not represent EPA policy.

**Table of Contents**

1			
2	<b>1.</b>	<b>EXECUTIVE SUMMARY.....</b>	<b>1</b>
3	<b>2.</b>	<b>INTRODUCTION.....</b>	<b>8</b>
4	2.1.	BACKGROUND.....	8
5	2.2.	CHARGE TO THE PANEL.....	8
6	<b>3.</b>	<b>RESPONSE TO CHARGE QUESTIONS.....</b>	<b>9</b>
7	3.1.	CONCEPTUAL APPROACH.....	9
8	3.2.	FLORIDA ESTUARIES.....	16
9	3.3.	FLORIDA COASTAL WATERS.....	22
10	3.4.	SOUTH FLORIDA INLAND FLOWING WATERS.....	25
11	3.5.	SOUTH FLORIDA MARINE WATERS.....	35
12	3.6.	DOWNSTREAM PROTECTION VALUES.....	38
13		<b>REFERENCES.....</b>	<b>R-1</b>
14		<b>APPENDIX A: CHARGE TO THE PANEL.....</b>	<b>A-1</b>
15		<b>APPENDIX B: PHOSPHORUS TRANSPORT AND FATE IN FRESHWATER SYSTEMS.....</b>	<b>B-1</b>
16		<b>APPENDIX C: OPTIONS FOR DISTRIBUTING LOADS.....</b>	<b>C-1</b>
17			

## 1. EXECUTIVE SUMMARY

Nitrogen (N) and phosphorus (P) inputs from urban and agricultural sources are known to influence water quality, and nutrient pollution has been identified as a source of impairment for estuarine, marine and fresh waters in Florida. The state of Florida has a narrative criterion for nutrients, which states that “in no case shall nutrient concentrations of a body of water be altered so as to cause an imbalance in natural populations of flora or fauna.” The criterion specifically refers to degradation from anthropogenic loads of total nitrogen (TN) or total phosphorus (TP). In 2009, EPA determined that numeric nutrient criteria were needed to protect aquatic life in Florida, and initiated a process to develop such criteria for categories of state waters.

The SAB was asked to review and provide advice on the proposed approaches for estuarine, coastal and South Florida waters, as described in the draft EPA document, *Methods and Approaches for Deriving Numeric Criteria for Nitrogen/Phosphorus Pollution in Florida’s Estuaries, Coastal Waters, and Southern Inland Flowing Waters*. An *ad hoc* SAB panel, the Nutrient Criteria Review Panel, was formed for this task. The Charge to the Panel included questions about the conceptual model used to select assessment endpoints, data sources, and possible approaches to define criteria for each of the categories of waters: estuaries, coastal waters, inland flowing waters (including canals) in South Florida, and South Florida marine waters.

The Panel acknowledges the substantial effort that already has been made by EPA to get the work to this stage, much of it solid and thoughtful. However, much work remains to be done. It is not clear what approaches will be selected, if multiple approaches will be used, and which approaches will provide useful information towards the goal of developing nutrient criteria. Under a court-ordered consent decree, the Agency has committed to proposing nutrient criteria for estuaries, coastal waters, and South Florida inland flowing and marine waters by November 14, 2011 and final criteria by August 14, 2012. Given this time frame, and the extent of the task at hand, the Panel is concerned that the Agency not sacrifice quality work for the sake of a schedule.

Different approaches are being considered for each of four major categories of Florida waters. Consistent with the Charge to the Panel, the Panel provides review and advice on the proposed approaches for developing nutrient criteria for each category of waters.

### Conceptual Model (Charge question 1a)

EPA proposes a conceptual model that links nitrogen and phosphorus levels in Florida waters to biological endpoints to be protected, using one or more analytical approaches. Although the general conceptual model provides a starting point for choosing numeric criteria, the Panel has numerous concerns about how TN and TP would be linked to measurable biological endpoints. TN and TP should be considered driver (rather than causal) variables; there are many factors other than nutrients that control Chl-*a*, and this is not acknowledged in the conceptual diagram. TN and TP loadings are likely to be better predictors of Chl-*a*, hypoxia and seagrass loss than TN or TP concentrations. TN and TP concentrations also can be considered

**Science Advisory Board Panel Draft Report (dated April 8, 2011) - Do not Cite or Quote**

This draft is a work in progress, does not reflect consensus advice or recommendations, has not been approved by the chartered SAB, and does not represent EPA policy.

1 response variables in that they reflect long-term loading rates; the temporal and spatial scales  
2 over which they would be measured should be clarified.

3 The conceptual diagram is a representation of important linkages, but Biological  
4 Endpoints and Objectives are not discussed in sufficient detail. The three biological endpoints  
5 (healthy seagrasses, balanced phytoplankton biomass, and balanced faunal communities) are  
6 appropriate. However, these endpoints need to be much better defined and, in some cases, more  
7 clearly connected to the explanatory variables that would be the basis for setting numeric criteria.  
8 It is critical that EPA define “balanced” for each of the three biological endpoints, preferably in  
9 quantitative terms. Simply relying on the state’s assessment of whether or not waters are  
10 “healthy” is not an adequate definition.

11 EPA proposes to identify levels of water quality variables that would protect the  
12 biological endpoints using reference condition, stressor-response, and/or modeling approaches  
13 for a particular water body. There would be a greater confidence in the criteria if all three  
14 approaches were applied, or as many as possible, to each of the systems where data and models  
15 are available. The EPA document should discuss how the results from multiple approaches  
16 would be integrated to develop the final numeric criteria. The three analytical approaches are  
17 being applied somewhat differently within the different categories of Florida waters, and each  
18 approach has different data requirements and—more importantly—different assumptions,  
19 limitations and uncertainties. EPA should describe the uncertainty associated with the various  
20 approaches to criteria development and discuss how this uncertainty might influence the use and  
21 appropriateness of specific numeric criteria.

22 The Panel agrees that Chl-*a* concentration in the water column is both sensitive to  
23 nutrient inputs and an important measure of ecosystem health and therefore a reasonable  
24 endpoint in itself. However, Chl-*a* alone, which measures phytoplankton biomass, cannot be  
25 used to infer anything about primary productivity or whether or not phytoplankton populations  
26 are “balanced” in terms of species composition or relative abundance/dominance. While we  
27 support using Chl-*a* as an endpoint, its limitations need to be explicitly recognized. Water  
28 column Chl-*a* also is linked to seagrass health because it is an indirect measure of water clarity,  
29 which influences seagrass photosynthesis. However, the Panel is concerned about relying upon  
30 Chl-*a* as the sole criterion to protect seagrasses because, in some systems, macroalgae or  
31 epiphyte growth can significantly impact seagrass communities even as water column Chl-*a*  
32 levels remain low. EPA should consider a stressor-response approach to link nutrient loading  
33 with seagrass areal extent for protecting seagrass communities.

34 The Panel is concerned that no direct measures of the faunal community are proposed to  
35 define whether a balanced community is being protected and maintained. Instead, EPA proposes  
36 to rely on attainment of the State of Florida dissolved oxygen (DO) standard as an indicator for  
37 the presence of balanced faunal communities. EPA proposes to look for relationships between  
38 TN and/or TP and DO, and use those relationships to determine numeric criteria for TN and TP  
39 that are protective. The Panel is concerned with the absence of any reference whatsoever to  
40 explicit faunal metrics.

1 **General Delineation of Florida Waters (Charge question 1b)**

2 In the document, EPA proposes an initial grouping of Florida waters into four categories:  
3 estuaries, coastal waters, South Florida inland flowing waters, and South Florida marine waters.  
4 Separation of estuarine and coastal waters is appropriate, and the separate consideration of South  
5 Florida also is warranted. The Panel recommends that the term “marine waters” be replaced  
6 with “estuarine and coastal waters.” South Florida inland flowing waters appears to be a default  
7 category without strong scientific rationale for this classification; for purposes of criteria  
8 development, these waters should be subdivided into managed canals and natural streams.  
9 Additional comments are provided on sub-delineation and classification within the four broad  
10 categories of waters, below.  
11

12 **Florida Estuaries**

13  
14 **Delineation and Data (Charge question 2a).** The geographic delineations of estuaries  
15 seem appropriate although it was not clear why a salinity of 2.7 psu rather than 0.5 psu was used  
16 to delineate the upper reaches of these systems. A finer classification based on degree of impact  
17 may be useful (e.g., to separate Caloosahatchee and St. Lucie estuaries from the others given  
18 their unique hydrologic relationship to Lake Okeechobee). EPA should consider adding tidal  
19 creeks as a separate ecosystem type because they have different characteristics than the open  
20 estuaries and therefore may require different nutrient criteria. The Panel has few issues with the  
21 data sets presented and has provided suggestions for additional sources of data.  
22

23 **Assessment Endpoints (Charge question 2b).** Healthy seagrass communities are an  
24 appropriate biological endpoint for Florida estuaries, given their widespread occurrence and  
25 ecological (and economic) significance. To protect seagrass communities, EPA should consider  
26 a measure of epiphyte abundance in addition to the proposed determination of Chl-*a* in the water  
27 column. Whether Chl-*a* is an appropriate endpoint for assessing “balanced” phytoplankton  
28 communities, depends on how EPA defines “balanced”, which has not been done. Direct  
29 indicators of faunal community balance should be considered in addition to DO. When hypoxic  
30 conditions are observed, impacts on the biota usually have already occurred. It is preferable to  
31 identify indicators that show stress on the faunal community before such degraded conditions  
32 develop. Diel variability in DO needs to be considered when establishing DO water quality  
33 targets. DO criteria may be better characterized by percent saturation than by concentration.  
34

35 **Approaches (Charge question 2c).** N and P may be limiting in different portions of the  
36 fresh-to-marine continuum, and in some cases may be co-limiting. Thus, a dual nutrient (N and  
37 P) strategy is warranted and we agree with EPA’s decision to take this approach. The Panel  
38 acknowledges the substantial effort made to date to collect data and evaluate possible approaches  
39 to criteria development for these waters, but is concerned that the timetable for completion of  
40 this work may be unrealistic. Previous experience suggests that if the reference condition  
41 approach can be implemented, it might be the most “time-efficient” pathway to developing  
42 nutrient criteria, although the stressor-response approach is worthy of more attention than it has  
43 been given. The Panel is aware of a separate EPA report on stressor-response approaches, but  
44 would have preferred additional discussion in the present document on how such approaches  
45 might be applied in Florida’s estuaries. We urge caution in EPA’s apparent emphasis on water  
46 quality modeling approaches; if this approach is selected, a reasonable representation of internal

**Science Advisory Board Panel Draft Report (dated April 8, 2011) - Do not Cite or Quote**

This draft is a work in progress, does not reflect consensus advice or recommendations, has not been approved by the chartered SAB, and does not represent EPA policy.

1 nutrient cycling needs to be included. In light of climate-related effects on hydrology and  
2 temperature regimes, as well as increases in freshwater withdrawals for human use, the Panel  
3 recommends that EPA consider the possibility that thresholds could be crossed, fundamentally  
4 changing these systems.

5 **Florida Coastal Waters (Charge question 3a)**

6 **Delineation and data sources.** EPA proposes to use remotely sensed chlorophyll  
7 ( $Chl_{RS-a}$ ) to develop a reference criterion associated with a balanced phytoplankton population in  
8 coastal waters. The approach is appropriate and sensible for this large, poorly sampled region,  
9 although the Panel is concerned that no direct measurements of nutrients are proposed to verify  
10 the strength of the relationship between pollutant loads from land and observed  $Chl-a$   
11 concentrations in coastal waters. EPA proposes to use chlorophyll data from coastal waters out  
12 to 3 miles. However, under the Clean Water Act, water quality criteria apply to state coastal  
13 waters, defined for Florida as waters out to 3 nautical miles on the Atlantic coast and 9 nautical  
14 miles on the Gulf coast. In developing coastal criteria, the Panel recommends that EPA consider  
15 remote-sensed chlorophyll in waters beyond the 3-mile zone because some blooms observed in  
16 coastal waters may form further offshore. In addition, data from the entire shelf should be used  
17 to improve the calibration of remotely sensed and field-measured (*in situ*) chlorophyll. The  
18 Panel further notes that the proposed coastal segments are a result of historical precedence, and  
19 EPA may wish to consider segments defined in terms of bathymetry.

20  
21 **Assessment Endpoints (Charge question 3 b).** While acknowledging that  $Chl_{RS-a}$  is the  
22 most feasible indicator of nutrient status for coastal waters, given available data and the lack of a  
23 better alternative, the Panel has several concerns: (1) there is a causal relationship between  $Chl_{RS-}$   
24  $a$  and nutrient loads but a number of other factors also influence  $Chl_{RS-a}$  levels; (2) few *in situ*  
25  $Chl-a$  data are available for calibration; (3) the ten-year remote-sensing dataset may not  
26 constitute an adequate baseline, given decadal-scale variability; and (4) the future availability of  
27  $Chl_{RS-a}$  data is in doubt, particularly given the loss of the SeaWiFS sensor.

28  
29 **Approaches (Charge question 3c).** In order to relate remotely sensed chlorophyll to  
30 water column chlorophyll levels, it is necessary to calibrate satellite sensor readings using field-  
31 measured chlorophyll data. The approach to calibration has been thorough, and the Panel agrees  
32 with use of *in situ* calibration data taken within 3 hours of the satellite overpass. The ratio  
33 between the chlorophyll concentrations in the upper two meters and the full euphotic zone needs  
34 to be established, and obvious antecedent bloom data points should be removed from analyses.  
35 Because available satellite-based sensors will change over time, as some platforms are retired  
36 and others are launched, the Panel recommends that EPA cross-calibrate data from existing  
37 sensors with future sensors as they become available.

38 **South Florida Inland Flowing Waters**

39 **Rationale for Criteria (Charge question 4).** The Panel recommends that EPA  
40 subdivide South Florida inland flowing waters into human-managed canals vs. natural flowing  
41 streams. Canals would be viewed as a source of nutrients to downstream, more oligotrophic,  
42 systems rather than for any valued ecological attributes that may be unique to them, and canal  
43 nutrient criteria would be in the form of DPV. Depending on data availability, criteria could be  
44 developed for natural flowing streams as instream protection values (IPV).

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46

**Delineation and Data Sources (Charge question 4a).** EPA proposes to derive numeric criteria for South Florida inland flowing waters, including canals, as instream protection values (IPV) for TN and TP using the reference condition approach. Canals apparently account for approximately 90 percent of waters in this category. The Panel is not convinced from the material provided that IPV nutrient criteria are appropriate for these uniquely artificial and highly managed ecosystems. The underlying problem is that the canals are classified as Class III waters (with a designated use of recreation and balanced population of fish and wildlife), although their primary purpose is flood control and management of water supply. The proposed inventory of inland flowing waters that catalogues and distinguishes natural streams and canals should provide very useful information, and could provide the basis for subdividing this category of waters as recommended above.

EPA is considering use of the Landscape Development Index (LDI) as an approach to identify areas with reference conditions. The LDI is a surrogate for stressors, including TN and TP loads, associated with various land uses. The 100-m stream/canal buffers proposed for use with the LDI may be too limited, and may not capture impacts from land uses in the full watershed. The proposed classification appears reasonable as it incorporates surface and subsurface flow regimes, soil types and land use; however legacy N and P loads from previous land management practices also must be considered. Use of the LDI is not recommended for canals.

**Assessment Endpoints (Charge question 4b).** If canal IPV's are to be developed, more information needs to be presented on how balanced natural populations would be assessed for these unique aquatic ecosystems. The Panel recommends further consideration and assessment of the response variables to be used and the form of the nutrients (i.e., those with short-term versus long-term bioavailability) that are most relevant. It was not clear that sufficient data are available to support either invertebrate or phytoplankton endpoints for canals.

**Approaches (Charge question 4c).** EPA proposes two approaches for determining numeric criteria for South Florida inland flowing waters. The first is based on reference conditions and the second is based on stressor-response relationships. If canal IPV's are developed, either of these approaches could work, although it was not clear if the available data would show interpretable patterns. Finding patterns in natural streams may be more likely than in the highly managed canals. For the reference condition approach, data on historical annual values of TN and TP from a set of least-disturbed sites (identified using  $LDI < 2$ ) would be used to develop lognormal distributions of TN and TP under least disturbance. Variability of nutrient levels at the least-disturbed sites will reflect heterogeneity in hydrology, geology, etc. Failure to account for such heterogeneity, which is also present in disturbed sites, may result in numeric criteria that are under- or over-protective. The Panel also notes that selecting "least disturbed sites" using an  $LDI < 2$  may not be feasible in this region that has been subject to active management for many years.

The stressor-response approach also should work if a suitable relationship between *Chl-a* and nutrient load can be demonstrated, but several of the same caveats apply here as for setting limits in coastal waters. As with the approach based on reference conditions, the relationship

**Science Advisory Board Panel Draft Report (dated April 8, 2011) - Do not Cite or Quote**

This draft is a work in progress, does not reflect consensus advice or recommendations, has not been approved by the chartered SAB, and does not represent EPA policy.

1 between Chl-*a* and TN or TP is likely to be modulated by the effects of hydrological, geological,  
2 and other covariates. Failure to account for such factors may lead to criteria that are over- or  
3 under-protective. The document needs to address how regression models will be used to  
4 determine numeric criteria, specifically, how they will determine the level of Chl-*a* considered to  
5 be protective of balanced phytoplankton and faunal communities.

6 **South Florida Estuarine and Coastal Waters**

7 **Delineation and Data (Charge question 5a).** South Florida estuarine and coastal waters  
8 have a different nutrient regime than other parts of the state, given their oligotrophic nature and  
9 susceptibility to upstream water management (versus nutrient regulatory) decisions. The Panel  
10 agrees that these waters should be considered separately for purposes of nutrient criteria  
11 development. The data identified for South Florida estuarine and coastal waters seem  
12 appropriate. However, the proposed subdivision/subclassification of these waters does not  
13 clearly relate to the oceanographic circulation and degree of connectivity in the region. EPA  
14 might consider an alternative segmentation scheme that incorporates alongshore and cross-shelf  
15 circulation patterns.

16  
17 **Approaches (Charge question 5b).** EPA proposes to use a reference condition  
18 approach using least-disturbed sites or a binomial test applied to a distribution of raw data. Both  
19 approaches have merit and the Panel encourages application of both to provide a more robust  
20 evaluation of criteria. However, if least-disturbed sites are those most distant from land-based  
21 sources, nutrient levels at these sites may reflect dilution by oligotrophic ocean water rather than  
22 a significant difference in nutrient loading. In addition, the coastal and estuarine waters of South  
23 Florida have experienced enormous changes over the last century in freshwater inflows, salinity  
24 and residence times, with associated changes in nutrient cycling and seagrass extent. These past  
25 alterations should be considered when defining reference conditions. The Panel recommends  
26 that seagrass coverage and extent of epiphytic colonization be considered as endpoints in  
27 addition to water column chlorophyll. The document should clarify which coastal and estuarine  
28 areas will be under the jurisdiction of the forthcoming nutrient criteria, versus other regulations  
29 (e.g., for federal and state protected waters).

30 **Downstream Protection Values (Charge question 6)**

31 EPA proposes to use DPV criteria to ensure that upstream N and P water quality criteria  
32 will be set at levels that will protect downstream estuarine designated uses. While agreeing with  
33 the goal, the Panel has concern with the overlap between DPV and the Total Maximum Daily  
34 Load (TMDL) process. For development of an estuarine TMDL, standard practice is to model  
35 the estuary and watershed to determine the additional pollutant load reduction needed, and then  
36 to allocate the load reduction based on input from state and local officials. This process thus  
37 allows consideration of socio-economic factors, natural background levels of nutrients in  
38 different tributaries, etc. in determining the load allocation. However, a significant drawback of  
39 relying on TMDLs rather than DPVs is the fact that the TMDL process is not triggered until  
40 impairment has occurred in a waterbody.

41  
42 The EPA document includes an example of a DPV developed using an equal load  
43 reduction for each tributary to an estuary. The Panel notes that this equal allocation approach  
44 ignores the possibility of different background nutrient concentrations in upstream segments, and

**Science Advisory Board Panel Draft Report (dated April 8, 2011) - Do not Cite or Quote**

This draft is a work in progress, does not reflect consensus advice or recommendations, has not been approved by the chartered SAB, and does not represent EPA policy.

1 other characteristics of segments that would make nutrient load reductions impractical. In  
2 response to a Panel request for further information, the Agency described four options for  
3 distributing loads for calculation of DPV, each of which presents some difficulties. If the DPV  
4 approach is pursued, we urge that selected apportionment strategies not preclude cross-tributary  
5 nutrient trading or other approaches to achieving the necessary load reductions.  
6

7 Whether developing a DPV or a TMDL, the modeling of load reduction apportionment  
8 for upstream segments is a valid approach, but watershed characteristics such as predominant  
9 land-use (especially urbanized area) should be considered. EPA should justify the choice of the  
10 LSPC model and explain why it is the most applicable model for this case. The timeframe of the  
11 modeling should be linked to the response of biological endpoints in the receiving waters; annual  
12 average values may grossly under-predict the impact of large storm events. In addition, the  
13 document needs to discuss the impact on criteria of P cycling and transformation.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32

## 2. INTRODUCTION

### 2.1. Background

Nitrogen (N) and phosphorus (P) inputs from urban and agricultural sources are known to influence water quality, and nutrient pollution has been identified as a source of impairment for estuarine, marine and fresh waters in Florida. The state of Florida has a narrative criterion for nutrients, and is in the process of developing numeric nutrient criteria for its estuaries and coastal waters. In 2009, EPA determined that numeric nutrient criteria were needed to protect aquatic life in Florida, and initiated a process to develop such criteria for categories of state waters. Criteria for total nitrogen (TN), total phosphorus (TP) and chlorophyll *a* (Chl-*a*)—a measure of water column algal biomass—were finalized for Florida lakes and inland flowing waters in 2010. Numeric nutrient criteria for estuarine and coastal waters, and South Florida inland flowing waters, are being developed separately, using a variety of approaches and ecological endpoints. The SAB was asked to provide review and advice on the proposed approaches for estuarine, coastal and South Florida waters, as described in the draft EPA document, *Methods and Approaches for Deriving Numeric Criteria for Nitrogen/Phosphorus Pollution in Florida’s Estuaries, Coastal Waters, and Southern Inland Flowing Waters* (November 17, 2010 draft; U.S. EPA 2010).

An *ad hoc* panel of the SAB, the Nutrient Criteria Review Panel, was formed for this task. The Panel met on December 13-14, 2010, to hear EPA technical presentations and public comments, and to discuss responses to the questions in the Charge to the Panel (Appendix A). A follow-up public teleconference of the Panel was held on February 7, 2011, to discuss an initial Panel draft report. The full SAB conducted a quality review of the Panel’s report on xxx.

### 2.2. Charge to the Panel

The Charge to the Panel included questions about the conceptual model used to select assessment endpoints, data sources for the various categories of waters, and possible approaches to define criteria for each of the categories of waters: estuaries, coastal waters, inland flowing waters (including canals) in South Florida, and South Florida marine waters. Relevant charge questions are included at the beginning of each section of the Panel’s report, and the full Charge to the Panel is included as Appendix A.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39

### 3. Response to Charge Questions

#### 3.1. Conceptual Approach

##### 3.1.1. Conceptual Model

*Charge Question 1(a). EPA has introduced a general conceptual model in Chapter 2, including the selection of assessment endpoint and indicator variables. What is your perspective of the general conceptual model?*

The State of Florida currently has a narrative criterion for nutrients that says, “in no case shall nutrient concentrations of a body of water be altered so as to cause an imbalance in natural populations of flora or fauna,” with specific reference to degradation from “man-induced nutrient enrichment (total nitrogen or total phosphorus)” (Florida Administrative Code 62-302.530). EPA has put forward a conceptual model, describing the relationships between nutrient levels (total nitrogen [TN] and total phosphorus [TP]) and biological responses, to guide its efforts to translate Florida’s current narrative nutrient criterion into numeric nutrient criteria. Although the general conceptual model (Figure 1, below) provides a starting point for choosing numeric criteria, the Panel has numerous concerns about how the causal variables will be linked to biological endpoints.

The conceptual model would translate Florida’s objective of “balanced natural populations of aquatic flora and fauna” into numeric criteria for three biological endpoints: seagrasses, phytoplankton, and faunal communities. While agreeing that these endpoints are appropriate, the Panel strongly felt that these endpoints need to be much better defined and, in some cases, connected to the explanatory variables that would be the basis for setting numeric criteria. As detailed below, the Panel did not always see a direct link between causal variables and the endpoints.

The term “balanced” is not defined in the document and is subject to a great range of interpretation. EPA needs to provide a definition of “balanced” early in the document. Simply relying on the state’s assessment of whether or not waters are “healthy” is not an adequate definition. EPA also should define how it will measure and interpret these three endpoints, preferably in quantitative terms. The Florida DEP has defined a balanced community as one that may have experienced “modest” change, but which still has: (1) reproducing populations of sensitive taxa; (2) a balanced distribution of all expected major groups; and (3) an ecosystem with functions that are largely intact due to redundant system attributes (FDEP 2010). If EPA chooses to use this definition, then the link between the water quality targets and these goals should be documented.

**Science Advisory Board Panel Draft Report (dated April 8, 2011) - Do not Cite or Quote**

This draft is a work in progress, does not reflect consensus advice or recommendations, has not been approved by the chartered SAB, and does not represent EPA policy.

1 More information is needed on the methodologies that will be used in order to determine  
2 if the general conceptual approach is workable within the time constraints<sup>1</sup>. The Panel  
3 recognizes that details on methods are to some extent specific to type of water body and  
4 appropriate for later chapters, but further information on the methods is needed in this chapter as  
5 well. EPA is proposing three general approaches to relate nutrient levels to balanced natural  
6 populations:

- 7
- 8 1. Identify reference conditions for a water body type based on available data or best  
9 professional judgment;
- 10 2. Use predictive stressor-response relationships and nutrient/algal thresholds; and/or
- 11 3. Use numerical water quality models to predict nutrient loadings that would be  
12 protective of system biology.
- 13

14 The Panel agrees that these approaches all have utility and that using a combination of  
15 approaches may be best where data and models are available. However, more detail is needed  
16 on: (1) the adequacy of the data for applying each approach; (2) how decisions would be made  
17 on which approaches to use; and (3) how discrepancies in targets derived from different  
18 approaches would be resolved.

19

20 Each of the three approaches has strengths and weaknesses. The use of **nutrient**  
21 **reference conditions** implies that nutrient concentrations and loadings to a system are known  
22 with enough certainty that target values protective of biological endpoints can be determined. In  
23 cases where data specific to a system are not sufficient, best professional judgment could be used  
24 to determine suitable target values. There has been extensive hydrologic modification of  
25 Florida's waters and extreme weather events, a number of which have occurred in the last 15  
26 years, which complicates defining reference conditions in the context of spatial and temporal  
27 variability. Current reference conditions may not represent historical conditions. EPA may need  
28 to state explicitly the general hydrologic ranges over which these targets will be useful and have  
29 clearly stated goals in cases where remediation is suggested. When using the reference condition  
30 approach, EPA also needs to evaluate whether data sets used to set values are representative of  
31 long-term averages or (in the case of relatively short-term data sets) are heavily influenced by  
32 recent hurricanes. Comments by Briceño et al. (2010) may be useful in this regard.

33 The use of predictive **stressor-response relationships** and thresholds assumes that data  
34 on nutrient-organism interactions from Florida waters and other regions, or countries, could be  
35 appropriately applied to setting protective target values. The use of numerical **water quality**  
36 **models** assumes that models would be a useful and realistic representation of nutrients and other  
37 water quality parameters. For practical application of numerical models, there still remain  
38 questions as to the appropriateness of selected models, availability of data, and level of detail  
39 required to adequately populate each model approach. For example, the EPA document states  
40 that a watershed model will be run with all anthropogenic sources removed to determine  
41 background TN and TP levels. More information and justification is needed to provide  
42 assurances that the models being used can adequately accomplish this with the stated degree of

---

<sup>1</sup> Under a consent decree, EPA is required to issue proposed criteria by November 14, 2011, and final criteria by August 14, 2012 (U.S. District Court, Northern District of Florida, Tallahassee Division, Case No. 4:08-cv-00324-RH-WCS).

**Science Advisory Board Panel Draft Report (dated April 8, 2011) - Do not Cite or Quote**

This draft is a work in progress, does not reflect consensus advice or recommendations, has not been approved by the chartered SAB, and does not represent EPA policy.

1 certainty. Most water quality models have been developed to assess and predict fate and  
2 transport processes as a result of anthropogenic activities and not for determining pristine  
3 conditions. Detailed validation of such uses is needed, which means calibration with non-  
4 impacted watershed loads. However, there are few non-impacted watersheds with conditions  
5 that reflect baseline concentrations in relation to determining water impairment. A key factor  
6 involved in using numerical models would be their validation and some analysis of uncertainty  
7 for each of the systems where they are applied. It may not be feasible to apply these models to a  
8 large number of estuaries in a short period of time.

9 The three methodological approaches listed above are being applied somewhat differently  
10 within the different categories of Florida waters. Since each approach has different data  
11 requirements (and more importantly, different assumptions, limitations and uncertainties), it is  
12 critical that the reasoning behind applying or not applying a given method be provided. The  
13 EPA document notes that EPA may use one, two or all three of these approaches for a particular  
14 water body. There would be a greater confidence in the criteria if all three approaches were  
15 applied, or as many as possible, to each of the systems for which data and models are available.  
16 This would provide an ensemble approach and a range of values for setting numeric criteria.  
17 However, this could result in three different answers as to what numeric values would be  
18 protective. This is understandable given the different conceptual bases for each approach, but  
19 the EPA document should discuss how the results from multiple approaches would be integrated  
20 to develop the final numeric criteria.

21  
22 Specific suggestions on the conceptual model for different ecosystem types follow.  
23 Further discussion of EPA's proposed approaches can be found in the responses to the charge  
24 questions for specific categories of waters.

25  
26 ***Conceptual Diagram***


27 The conceptual diagram (Figure 1) gives a general overview of how the conceptual  
28 model will be linked to three biological endpoints and how the approaches would be applied in  
29 each of the categories of Florida waters. The upper three levels of the diagram (Causal Variable,  
30 Response Variable, and Water Quality Targets) are dealt with at great length, but the bottom two  
31 levels (Biological Endpoints, Objective) are not discussed in sufficient detail. We recommend  
32 that terminology that implies TN and TP are causal variables be dropped in favor of terms such  
33 as driver variables. While there is a cause/effect relationship between nutrients and Chl-*a*, there  
34 are many other factors that control Chl-*a*, and these factors are not adequately acknowledged in  
35 the diagram.

36  
37 In Figure 1, Chl-*a* also is shown to be a water quality target that relates to balanced faunal  
38 communities. There are mechanisms by which these two are linked, in addition to changes in  
39 DO (e.g., through increased sediment loading), but we did not find any discussion of mechanistic  
40 links in the EPA document. Also, while low DO is closely linked with eutrophication, it is not  
41 the only mechanism of nutrient impacts, which is what is implied in the diagram. The Panel  
42 suggests that EPA alter the diagram or include an explanation on how numeric criteria for Chl-*a*  
43 will be linked to balanced faunal communities. EPA also should provide more background and  
44 theory on the relationships between biological endpoints and water quality targets. There are  
45 many factors that regulate "balanced" ecosystem functions in addition to the few listed in Figure

**Science Advisory Board Panel Draft Report (dated April 8, 2011) - Do not Cite or Quote**

This draft is a work in progress, does not reflect consensus advice or recommendations, has not been approved by the chartered SAB, and does not represent EPA policy.

1 1, including predation and other food web interactions, harvest, salinity, substrate, species  
2 turnover, and N:P ratios.



3  
4 **Figure 1. EPA's Proposed Conceptual Diagram Relating TN/TP Criteria to Florida's Narrative Nutrient**  
5 **Criterion (Source: U.S. EPA 2010, Fig. 2-1)**

6  
7 An additional concern is that the conceptual model does not include any discussion about  
8 how seasonal and inter-annual variability will be addressed. Average annual values for  
9 parameters may not be protective if there are wide seasonal changes. Seasonal variability (e.g.,  
10 warm vs. cold, wet vs. dry) should be evaluated to determine whether multiple criteria are  
11 required.

12  
13 The goal of the criteria development effort is protection of biological populations, and so  
14 the Panel begins its comments with the biological endpoints to be protected and works through  
15 the conceptual diagram from the bottom up.

16  
17 ***Protection and Restoration of Healthy Seagrass Communities***

18 Chapter 3 of the EPA document describes in more detail how a healthy seagrass  
19 population might be determined using historical data and colonization depth. This is a specific  
20 and quantifiable parameter (see section 3.2.2). A brief explanation of this is needed in Chapter 2

**Science Advisory Board Panel Draft Report (dated April 8, 2011) - Do not Cite or Quote**

This draft is a work in progress, does not reflect consensus advice or recommendations, has not been approved by the chartered SAB, and does not represent EPA policy.

1 to outline the approach. We did not find specific decision criteria for determining when  
2 management objectives have been met for impaired water bodies or what magnitude of changes  
3 would be considered a significant change (i.e., what percent of historical seagrass coverage  
4 would be set as a target for restoration?). Data tables provided in Appendix B of the EPA  
5 document (Estuarine and Marine Assessment Endpoints and Water Quality Indicator Variables  
6 Literature Review) may be used as a starting point. A thorough presentation of key parameters  
7 and breakpoints needs to be included if numeric criteria are to be applied.  
8

9 The Panel is concerned about relying upon water column Chl-*a* as the sole criterion to  
10 protect seagrasses. No numeric criteria directly related to macroalgae or epiphytes are being  
11 proposed. In systems where the nutrients are largely taken up by the phytoplankton, Chl-*a* will  
12 reflect the major impact of nutrient loading. However, there are systems where macroalgae  
13 proliferate in response to increased nutrient loadings, but water column Chl-*a* remains low due to  
14 short water residence times. In these systems, water column Chl-*a* is a poor measure of nutrient  
15 effects. Hauxwell et al. (2001, 2003) found that light levels in benthic macroalgal mats  
16 prevented young eelgrass shoots from being established. Epiphytes also can increase in response  
17 to nutrient loading in systems even though water column Chl-*a* levels remain fairly low.  
18

19 EPA could consider an approach linking nutrient loading with changes in seagrass areal  
20 extent for protecting seagrass communities. This approach has been successful in Tampa Bay  
21 (Greening 2010). It was also applied to a range of systems in New England (Latimer and Rego  
22 2010), with data on eelgrass loss for a number of estuaries being compared to calculated nutrient  
23 loadings. The Latimer and Rego study found eelgrass loss began to occur at N loads above 50  
24 Kg ha<sup>-1</sup> y<sup>-1</sup> and eelgrass disappeared at 100 Kg ha<sup>-1</sup> y<sup>-1</sup>. It may be possible to develop a similar  
25 relationship for Florida seagrass systems. The Panel recommends that EPA consider this  
26 approach and also ways to back-calculate concentrations from specific loadings that impair  
27 seagrasses.  
28

29 ***Phytoplankton Production and Biomass***

30 The Panel agrees that Chl-*a* concentration is both sensitive to nutrient inputs and an  
31 important measure of ecosystem health and therefore a reasonable endpoint in itself. However,  
32 Chl-*a*, which measures biomass, cannot be used to infer whether or not populations are  
33 “balanced” in terms of species composition or relative abundance/dominance. In testimony,  
34 EPA provided examples where toxic blooms are known to occur at high Chl-*a* values. Although  
35 undesirable species may be more prevalent in areas with higher nutrient loading (and higher algal  
36 biomass), low biomass does not ensure that a toxic species will not occur or that species  
37 composition has not changed. Similarly, while Chl-*a* is a measure of biomass (standing stock), it  
38 is not a measure of production (a rate) and cannot be used to assess the biological endpoint of  
39 production. In sum, the Panel supports using Chl-*a* as an endpoint, but its limitations need to be  
40 recognized.  
41

42 ***Balanced Faunal Communities***

43 The conceptual model does not include a direct metric for balanced faunal communities,  
44 but instead proposes that healthy faunal communities rely upon sufficient concentrations of DO.  
45 While this is true and the document cites studies where low DO (hypoxia and anoxia) causes  
46 mortality and impairment of marine life, low DO usually is a secondary response to excess

**Science Advisory Board Panel Draft Report (dated April 8, 2011) - Do not Cite or Quote**

This draft is a work in progress, does not reflect consensus advice or recommendations, has not been approved by the chartered SAB, and does not represent EPA policy.

1 nutrients in a system. The primary response of excess nutrients is eutrophication, the excess  
2 production and accumulation of organic matter in a system. For Florida, in particular, there are  
3 also many swamp and backwater systems that feed into tidal waters that are naturally low in DO  
4 and not related to TN or TP concentrations or loadings.

5  
6 EPA proposes to use the State of Florida’s DO standard to maintain the biological  
7 endpoint of balanced faunal communities. Specifically, EPA proposes to look for relationships  
8 between TN and/or TP and DO, and use those relationships to determine numeric criteria for TN  
9 and TP that are protective (i.e., that are associated with attainment of the existing DO standard).  
10 How linkages between these variables and faunal metrics will be assessed needs to be clarified.  
11 Chapter 3 (p. 49) of the EPA document implies that the absence of hypoxia will be the indicator  
12 for the presence of balanced communities, so that numeric criteria would be based on ambient  
13 nutrient levels where hypoxia is absent. The Panel is concerned with the absence of any  
14 reference whatsoever to explicit faunal metrics (see also response to charge question 2).

15  
16 Chapter 3 also notes that DO can be computed in water quality models from TN and TP  
17 loading. However, in determining if hypoxia is present in any particular system, seasonality  
18 needs to be considered as hypoxia is a seasonal phenomenon and not present all year round

19  
20 ***Dissolved Oxygen Targets***

21 Additional concerns arose during the discussions and submitted testimony about using a  
22 single DO standard. Some seagrass meadows routinely exhibit low oxygen conditions at night  
23 even in the absence of any nutrient impairment. This diel cycling of oxygen—from  
24 supersaturated during daylight hours to undersaturated, and at times hypoxic, at nighttime—has  
25 recently been found to be common in shallow vegetated and unvegetated habitats (Moore et al.  
26 2004; Verity et al. 2006; Tyler et al. 2009). Similar conditions appear to occur in some Florida  
27 waters (see Boyer and Briceño 2010). Another issue is that oxygen is less soluble under higher  
28 temperatures and higher salinity, conditions seen in many of Florida’s warm temperate and  
29 subtropical waters. Hence, low DO criteria may be better characterized by percent saturation.  
30 Although the Florida DO numeric standard is not a subject of the current review, the Panel raises  
31 these issues to point out some of the challenges in relying on a DO standard to protect healthy  
32 biological communities. There are factors other than nutrients involved in development of  
33 hypoxia and anoxia such as water column stratification and water residence time.

34  
35 ***TN and TP Criteria***

36 In the EPA document, TN and TP are listed as “causal variables” and defined (p. 39) as  
37 concentrations (mg/L) of total (organic and inorganic) N and P. This may lead to confusion. As  
38 the table on page 36 of the document points out, TP and TN loadings are normally considered to  
39 be the ultimate driver of ecosystem changes while TP and TN water column concentrations are  
40 “associated with influent loading over the long term”. Hence this would make water column  
41 concentrations of both TP and TN explanatory or response variables. The narrative (p. 53) also  
42 refers to loading as the causal variable and water column concentrations as a response variable.  
43 This distinction is important when considering using TP or TN to predict other parameters.

44  
45 Many assessments have been based upon loading. Loading data, when available, would  
46 be expected to be a better predictor of Chl-*a*, hypoxia and seagrass loss than concentration.

**Science Advisory Board Panel Draft Report (dated April 8, 2011) - Do not Cite or Quote**

This draft is a work in progress, does not reflect consensus advice or recommendations, has not been approved by the chartered SAB, and does not represent EPA policy.

1 Furthermore, TN and TP concentrations may co-vary with Chl-*a*, since both are contained in  
2 phytoplankton, so they also are not completely independent from Chl-*a*. This presumes TN and  
3 TP are measured on unfiltered samples; yet the document does not clearly state whether that is  
4 what is proposed. Given the availability of data, there may be excellent reasons to use TN and  
5 TP concentrations as numeric criteria but they should be considered response variables. It also  
6 would be useful to characterize these variables in more detail, including the temporal and spatial  
7 scales over which they would need to be measured (i.e., weekly or monthly averages, surface  
8 values, depth-integrated samples, discrete depths).

9  
10 The issue remains of whether TN and TP or “reactive N and P” (i.e., dissolved inorganic  
11 N [DIN] and P [DIP]) are the most relevant variables to link nutrient enrichment to specific  
12 effects on biological endpoints (i.e., primary production, biomass as Chl-*a*, and cascading effects  
13 such as food web alterations and hypoxia). This issue has been the subject of considerable  
14 research, discussion and controversy for decades. Much of the uncertainty regarding whether to  
15 use TN and TP or more “reactive” dissolved forms of these nutrients revolves around the  
16 bioreactivity and roles of organic forms of these nutrients. Bioreactivity may be system-specific  
17 (or even system-component-specific), adding to the complexity and uncertainty of measuring  
18 responses and impacts on water quality and habitat condition. It is important for EPA to discuss  
19 this issue in the context of developing numeric nutrient criteria for nutrient-sensitive waters, both  
20 in Florida and nationally.

21  
22 Regardless of form, some consideration is needed for what portion of TN and TP loading  
23 in a system is from natural sources versus anthropogenic sources. This is particularly important  
24 for open coastal waters where conditions may be influenced by non-anthropogenic nutrient  
25 sources from outside the geographic boundaries of the coastal zone. While we know that  
26 reducing nutrients is key to restoring ecosystems in general, careful consideration needs to be  
27 given to all sources of N and P that in combination affect the biological endpoints for a system.

28  
29 ***Uncertainty***

30 Throughout the document, uncertainty is mentioned only briefly, and that in the context  
31 of uncertainty introduced because some environmental variables can covary with the explanatory  
32 variable of interest. It would be easier to convey the concept of statistical uncertainty by  
33 presenting confidence intervals. However, the level of uncertainty or size of confidence intervals  
34 associated with particular numeric criteria should be described further. Predictions developed  
35 using stressor-response models should explicitly state the level of confidence and be “ground-  
36 truthed” using site-specific data. The possibility that multiple covarying factors are involved in  
37 the cause-effect relationship for a particular system should be assessed. For example, the  
38 morphology of a system, habitat, and biological interactions within the water body may modify  
39 the relationship between nutrient concentrations (both N and P) and observed biological  
40 endpoints. These covarying factors need to be better documented, so that the uncertainty of the  
41 relationship can be reduced. In other words, a relationship between nutrient levels and a  
42 biological metric could be derived that is statistically significant, yet which accounts for only a  
43 small portion of the total variability in the data. Inclusion of covarying factors would strengthen  
44 statistical relationships, reduce uncertainty, and provide greater confidence in the final criterion.

1 **3.1.2. Categories of Florida Waters**

2 *Charge Question 1(b). EPA has delineated the State of Florida into 4 general categories*  
3 *of waters—Florida estuaries, Florida coastal waters, South Florida inland flowing*  
4 *waters, and South Florida marine waters—for purposes of considering approaches to*  
5 *numeric nutrient criteria development. Are these categories appropriate and*  
6 *scientifically defensible?*

7 Separation of estuarine and coastal waters is appropriate given the differences in natural  
8 populations of aquatic flora and fauna between higher salinity coastal systems and lower salinity  
9 estuarine and inland systems. Freshwater management in the region is complex and the separate  
10 consideration of South Florida also is warranted, although the Panel recommends that the term  
11 “marine waters” be replaced with “estuarine and coastal waters” for clarity and consistency. A  
12 finer classification based on degree of impact may be useful; for example, to separate the  
13 Caloosahatchee and St. Lucie estuaries from the other Florida estuaries, given their unique (i.e.,  
14 strong human influence) hydrological relationship to Lake Okeechobee. While nutrients clearly  
15 influence the biota in these systems, salinity levels play a stronger role than is typically the case  
16 in other Florida estuarine systems (Kraemer et al. 1999; Doering et al. 1999; Steinman et al.  
17 2002a).

18 The category of South Florida inland flowing waters seems to be a grab bag for waters  
19 that are primarily man-made canals, but which also include a small percent of natural streams.  
20 These canals present a unique challenge for setting numeric nutrient standards because of their  
21 morphology and management regime. It would be preferable to have a strong scientific rationale  
22 for this classification, and further information on canal management could be helpful in this  
23 regard.

24 **3.2. Florida Estuaries**

25 **3.2.1. Delineation and Data Sources**

26 *Charge Question 2(a). Are the data sources identified appropriate for use in deriving*  
27 *numeric criteria in Florida’s estuaries (as discussed in Sections 2.4 and 3.2)? Is the SAB*  
28 *aware of additional available, reliable data that EPA should consider in delineating*  
29 *estuaries or deriving criteria for estuarine waters? Please identify the additional data*  
30 *sources.*

31  
32 In general the geographic delineations of estuaries seem appropriate. We were unclear  
33 why a salinity of 2.7 psu was used to delineate the upper reaches of systems. Traditionally, such  
34 salinity would denote an oligohaline region of an estuary. Why not use “head of tide” or salinity  
35 of less than 0.5 psu? In any case, sediment nutrient dynamics change in this salinity transition  
36 zone (from approximately 0.0 to 5.0 psu). For example, at the toe of the salinity front, P-releases  
37 from sediments can increase sharply. Wherever the upper boundary is fixed, such issues need to  
38 be considered.

39  
40 EPA should consider adding another unit to the estuary delineation that would focus on  
41 tidal creeks. A case was made that these systems are common but have different characteristics  
42 than the open estuaries and therefore should have different nutrient criteria.

**Science Advisory Board Panel Draft Report (dated April 8, 2011) - Do not Cite or Quote**

This draft is a work in progress, does not reflect consensus advice or recommendations, has not been approved by the chartered SAB, and does not represent EPA policy.

1  
2 The Panel has few, if any, issues with the data sets presented. The summary tables in the  
3 EPA document indicate a careful review of data sources, including attention to time-series data.  
4 We encourage continued searching for appropriate data. In public comments to the Panel, one  
5 researcher (Dr. Tom Frazer, University of Florida) indicated that additional data are available for  
6 some estuarine areas and have yet to be utilized. It may be that County agencies have data sets  
7 not yet considered. This effort could be especially useful in the Big Bend area, where offshore  
8 seagrass beds are extensive, satellite data on Chl-*a* are not useful, and existing data sets from  
9 prior studies are rare. All data sets would need to meet EPA requirements for QA/QC, but the  
10 Panel encourages EPA to continue consultations with state and local agencies and researchers to  
11 access additional data and local knowledge where possible.  
12

13 **3.2.2. Assessment Endpoints**

14  
15 *Charge Question 2(b). Are the assessment endpoints identified in Sections 2.3 and 3.2*  
16 *(healthy seagrass communities; balanced phytoplankton biomass and production; and*  
17 *balanced faunal communities) appropriate to translate Florida's narrative nutrient*  
18 *criterion into numeric criteria for Florida's estuaries, given currently available data?*  
19 *Does the SAB suggest modification or addition to these assessment endpoints?*  
20

21 ***Healthy Seagrass Communities***

22 Florida seagrass beds are an extremely valuable natural resource, and the two largest  
23 contiguous seagrass beds in the continental United States are found in the Florida Keys and  
24 Florida's Big Bend region. Approximately 2.2 million acres of seagrass have been mapped in  
25 estuarine and near-shore Florida waters by researchers at the Florida Fish and Wildlife Research  
26 Institute in St. Petersburg (Carlson and Madley 2007). However, when seagrass beds growing in  
27 water too deep to easily map are included, the total area of seagrasses within Florida waters and  
28 adjacent federal waters is likely over 3 million acres. Florida's seagrass beds improve water  
29 quality and reduce shoreline erosion, but their most important ecological role is to provide food  
30 and shelter for many economically important finfish and shellfish species (Carlson and Madley  
31 2007). Estimates of the ecological services provided by Florida seagrass beds are approximately  
32 \$20,500 per acre per year (Handley et al. 2007), and it is entirely appropriate that EPA use  
33 healthy seagrass communities as one of its assessment endpoints.  
34

35 There are, however, several issues relating to seagrasses that deserve further  
36 consideration. First, as acknowledged in the draft EPA document, Chl-*a* usually explains a  
37 significant amount of variation in water clarity, but frequently does not explain the majority of  
38 this variation. Water clarity is often greatly influenced by colored dissolved organic matter  
39 (CDOM) and inorganic suspended material in the water column. Of greater importance,  
40 seagrasses in the shallow waters of Florida are typically shaded more by epiphytes growing on  
41 their leaves and by associated macroalgae (see Dixon 1999, and review by Burkholder et al.  
42 2007) than by Chl-*a* in the water column. Thus, EPA should consider a measure of epiphyte  
43 abundance in addition to the proposed determination of Chl-*a* in the water column. Epiphyte  
44 abundance is also often controlled to a significant extent by the animals that feed on these  
45 epiphytes (Hughes et al. 2004; Burkepile and Hay 2006; Heck and Valentine 2007; Baden et al.  
46 2010). This control by consumers, often referred to as top-down control, is beyond the scope of

**Science Advisory Board Panel Draft Report (dated April 8, 2011) - Do not Cite or Quote**

This draft is a work in progress, does not reflect consensus advice or recommendations, has not been approved by the chartered SAB, and does not represent EPA policy.

1 the draft document, but it is notable that both nutrients and herbivorous consumers affect the  
2 condition of seagrass meadows. As pointed out in Section 3.4.3, similar consumer effects can  
3 also be important in controlling Chl-*a* in the water column (Cloern 1982; Cohen et al. 1984;  
4 Thompson et al. 2008).

5  
6 It might also be possible to develop direct seagrass targets. In comments to the Panel, Dr.  
7 Paul Carlson suggested submerged aquatic vegetation (SAV), which includes seagrasses, as a  
8 management endpoint in support of the EPA plan but went further, arguing that SAV occurrence  
9 (abundance, density, etc) might be thought of as an integrative numeric criterion. In support of  
10 this suggestion, he states the following: (a) measures of SAV occurrence over time and space can  
11 integrate water quality issues of hypoxia, water clarity and nutrient conditions; (b) Florida has  
12 many areas dominated by SAV and thus these plant communities are representative of many  
13 estuaries; and (c) some beds are surveyed every two years, so there is a very good temporal  
14 record of their condition. The Panel suggests that serious consideration be given to Dr. Carlson's  
15 suggestion.

16  
17 ***Balanced Phytoplankton Biomass and Production***

18 As noted earlier (3.1.1), EPA needs to provide a clear definition of “balanced”. If EPA is  
19 not referring to species composition and relative abundance, but rather the entire phytoplankton  
20 or benthic microalgal communities, then Chl-*a* or other indicators of biomass (i.e., dry weight,  
21 particulate C, total cell counts) will suffice. If EPA is referring to species diversity or some other  
22 index of biological diversity, then more specific techniques will have to be employed, including  
23 microscopic species identification, photopigment analyses, molecular analyses, etc. We  
24 recommend community-level biomass metrics, using Chl-*a* or other indicators of biomass, as this  
25 is best related to nutrient and C-flux, hypoxia, and other drivers/indicators of impacts of and  
26 responses to nutrient inputs. Endpoints that require taxonomic-level resolution (e.g., to  
27 characterize harmful algal blooms, HAB) will need more specific suites of indicators to identify,  
28 quantify and characterize factors such as toxicity and food web effects. Taxonomic  
29 identification of HABs can provide diagnostic value, but the Panel acknowledges that such  
30 taxonomic analysis may not be possible with current monitoring programs in the systems and  
31 regions of interest.

32  
33 ***Balanced Faunal Communities***

34 There is little discussion in Chapter 3 of how “balanced faunal communities” are defined,  
35 and this is a concern for several reasons. First, given the generally shallow nature of Florida  
36 estuaries and our general impression that water clarity is (or was) high, it is likely that these  
37 systems were (or are) benthic-dominated. If this is the case, a variety of benthic heterotrophs  
38 could provide strong metrics for estuarine health, as a complement to seagrass community  
39 measures. Second, a strong shift from one common benthic species to another (e.g., a pollution  
40 tolerant species) can provide a good indicator of benthic habitat condition or deterioration,  
41 although the “species pair” might differ among estuaries. Has this approach, using indicator  
42 species, been considered as a measure of the health of faunal communities for Florida estuaries?  
43

44 Chapter 3 indicates that hypoxia will be used as an indicator of compromised benthic  
45 habitat condition. As a first pass, this will certainly tell us something about these habitats but not  
46 all that is needed. Unfortunately, when hypoxic conditions are observed, impacts on the biota

**Science Advisory Board Panel Draft Report (dated April 8, 2011) - Do not Cite or Quote**

This draft is a work in progress, does not reflect consensus advice or recommendations, has not been approved by the chartered SAB, and does not represent EPA policy.

1 usually have already occurred. It would be useful to have indicators of stress on the faunal  
2 community before such degraded conditions develop. In addition, DO in Florida seagrass  
3 meadows during the early morning hours is often below the levels considered to be hypoxic in  
4 unimpaired Florida waters, owing to the low amounts of oxygen that can be dissolved in the high  
5 temperature and high salinity waters characteristic of Florida and the high rates of nighttime  
6 respiration in the biomass-rich seagrass meadows. Thus, as suggested in comments provided by  
7 Boyer and Briceño (2010), a percent saturation criterion may be more useful than an absolute  
8 measure of oxygen concentration.

9 **3.2.3. Approaches**

10 *Charge Question 2(c). EPA describes potential approaches in Section 3.3 (reference*  
11 *conditions, stressor response relationships, and water quality simulation models) for*  
12 *deriving numeric criteria in Florida’s estuaries. Compare and contrast the ability of each*  
13 *approach to ensure the attainment and maintenance of natural populations of aquatic*  
14 *flora and fauna for different types of estuaries, given currently available data?*  
15  
16

17 As noted previously, the Panel recommends that EPA provide a more quantitative  
18 description of the concept of balanced phytoplankton and faunal communities, and remove the  
19 word “production” in the description of phytoplankton unless measures of production are added.  
20 Nutrient criteria development should take into account the natural diversity of Florida estuarine  
21 systems. For example, in some systems having low N/P nutrient ratios, blue-green algae may be  
22 the normal dominant species. Recognition of special system features will prevent systems from  
23 failing to meet criteria on the basis of natural background conditions. In addition, the estuarine  
24 continuum, from freshwater to the sea, often involves a transition from P- to N-limitation and  
25 possibly zones where co-limitation occurs. Thus, a dual nutrient strategy is warranted and we  
26 agree with EPA’s decision to take this approach. Similar strategies have been adopted in the  
27 Chesapeake Bay, Neuse River Estuary and Baltic Sea (e.g., see Elmgren and Larsson 2001;  
28 Conley et al. 2009; Paerl 2009).  
29

30 The Panel has a general concern that the timetable for completion of this work is  
31 unrealistic. A substantial effort already has been made to get the work to this stage, much of it  
32 solid and thoughtful. However, much work remains to be done. In the case of Florida estuaries,  
33 it is not clear what approaches will be selected, if multiple approaches will be used, and which  
34 approaches will provide useful information towards the goal of developing nutrient criteria. So,  
35 we are concerned that the Agency not sacrifice quality work for the sake of a schedule.  
36

37 The Panel emphasizes that there is no single approach that is ideal for developing nutrient  
38 criteria. This being the case, we support using multiple approaches where possible. If results for  
39 two or more approaches converge, then there is increased confidence in the results, and EPA  
40 needs to provide guidance on how to use this information to develop a criterion. If different  
41 approaches yield conflicting results, then EPA needs to have a defensible methodology to resolve  
42 the differences and move forward.  
43

1 **Reference Condition**

2 Philosophically, the reference condition approach is the most satisfying, although making  
3 it operational is often difficult because sufficient data are lacking to define “reference  
4 conditions”, and because of the problem of “shifting baselines” (Pauly 1995)—in other words,  
5 many ecosystems have been impacted by human activities for some time and we run the risk of  
6 using degraded coastal environments as reference conditions when the true (unimpacted)  
7 reference conditions have long since ceased to exist. The Panel is aware of at least one other  
8 State (New Hampshire) using the reference approach for developing nutrient criteria and that  
9 effort yielded some useful results. Experience suggests that this approach, if it can be  
10 implemented, might be the most “time-efficient” pathway to developing nutrient criteria.  
11

12 **Stressor-Response Models**

13 The Panel was disappointed that more attention was not given to the stressor-response  
14 approach. One of the Nation’s best estuarine restoration examples is Tampa Bay, where a  
15 stressor/response approach was used to develop local criteria. The Tampa Bay success suggests  
16 that this approach should be utilized more in the State-wide effort. The Panel is aware of a  
17 separate EPA report on the stressor-response approach (U.S. EPA 2010), but would have  
18 preferred to see more discussion in the current document of how this approach might be applied  
19 to development of nutrient criteria in Florida. Limnologists have had great success with this  
20 approach. Recently, EPA staff in New England published results of an analysis relating nutrient  
21 loads to seagrass health in a variety of small coastal systems (Latimer and Rego 2010). These  
22 sorts of studies suggest that the stressor-response approach needs to be more seriously  
23 considered. The EPA document simply refers to “regression models,” leaving many readers with  
24 the impression that EPA is considering only the simplest forms of regression analysis. (In  
25 contrast, the discussion of simulation modeling packages is presented in considerable detail.)  
26 Statistical models are correlative and the amount of variance explained by the correlations can be  
27 less than that needed for criteria development. However, we encourage EPA to give thorough  
28 consideration to the stressor-response approach when developing criteria for Florida waters.  
29

30 **Water Quality Simulation Models**

31 The level of detail on simulation modeling in the EPA document suggests that EPA has  
32 decided to use modeling as the primary tool for development of nutrient criteria. This may not  
33 be the case but we urge some caution here. The description of the model(s) sounds great, which  
34 can be quite seductive, and some issues can only be addressed with simulation models (e.g.,  
35 forecasting, understanding highly interactive processes). However, using simulation models  
36 would be a major undertaking, requiring considerable time and money, and useful results are not  
37 guaranteed. (Note that the Chesapeake Bay model has been under development for about 25  
38 years and still does not predict inter-annual hypoxic volumes well.) There also are very  
39 considerable issues related to data needed for calibration and verification of model results.  
40

41 If the simulation modeling approach is used, a reasonable representation of internal  
42 nutrient cycling needs to be included. In the generally shallow Florida systems, benthic  
43 processes will be especially important. In addition, these processes will interact with  
44 temperature and flow changes. Ultimately, nutrient concentrations reflect the net effect of these  
45 biogeochemical processes, as well as loadings.

1

## 2 ***Hydrologic Forcing***

3 From a spatial perspective, the location of phytoplankton production and biomass  
4 responses to nutrient inputs is strongly influenced by freshwater inflow and its impacts on  
5 estuarine residence time. Under drought conditions, the biomass peak, or Chl-*a* maximum ( $C_{\max}$ )  
6 will tend to be in the most upstream portion of estuaries (Valdes-Weaver et al. 2006). Under  
7 moderate freshwater discharge and flow conditions,  $C_{\max}$  will form in mid-estuarine locations,  
8 while under high flow conditions,  $C_{\max}$  will tend to predominate downstream (Valdes-Weaver et  
9 al. 2006; Paerl et al. 2007). Under extreme hydrologic conditions resulting from tropical  
10 cyclones,  $C_{\max}$  may not form at all, but rather the maximum phytoplankton biomass response will  
11 be in the sounds and coastal waters (Paerl et al. 2006a, b). These conditions represent a special  
12 challenge, because it may be difficult to evaluate and assign numeric criteria for nutrient loads to  
13 estuaries, as the response may not occur in estuarine waters.

14

15 The earth is experiencing a sustained period of elevated tropical cyclone activity and  
16 intensity (Webster et al. 2005). Florida is particularly impacted, because it experiences more  
17 tropical cyclone strikes than any other state in the U.S. Therefore, this aspect of climate change  
18 needs to be factored into the anticipated/predicted responses to nutrient inputs and the  
19 development of nutrient criteria. Conversely, periods of extreme (and record) droughts require  
20 additional attention and consideration in the context of the development of nutrient criteria, as  
21 the location and amounts of phytoplankton biomass responses to nutrient inputs will be  
22 dramatically affected.

23

## 24 ***Groundwater and Surface Water Withdrawals***

25 Florida is undergoing significant increases in freshwater withdrawal (for drinking water  
26 and agricultural irrigation purposes) from its lakes and rivers (Steinman et al. 2004). In addition,  
27 continued (and increasing) groundwater withdrawal will reduce freshwater recharge from springs  
28 and other natural seepage sources. These withdrawals will impact freshwater discharge to  
29 estuarine and coastal waters, which in turn will affect the location and magnitude of  
30 phytoplankton (including HABs such as cyanobacteria and dinoflagellates) responses to nutrient  
31 inputs, as well as the responses of benthic microalgae and macrophytes. Growing human  
32 demand for freshwater will need to be factored into the formulation of nutrient criteria as it will  
33 influence freshwater discharge, nutrient loads, nutrient concentrations and microalgal responses  
34 in impacted estuaries. Changes in freshwater flows add further uncertainty to derivation of  
35 nutrient criteria based on historical relationships and will need to be considered in triennial  
36 reviews of nutrient standards

37

## 38 ***Climate and Temperature Changes***

39 In addition to climate-related hydrologic effects, changes in temperature need to be  
40 considered. Changes in the range of 1.5 °C have been noted in some systems during the past 60-  
41 70 years. Temperature changes on this order can influence phytoplankton community  
42 composition (i.e., “cyanobacteria like it hot”; Paerl and Huisman 2008), as well as key  
43 biogeochemical nutrient and organic matter transformation processes (e.g., nitrification,  
44 denitrification, and sediment oxygen demand). These changes may alter relationships between

**Science Advisory Board Panel Draft Report (dated April 8, 2011) - Do not Cite or Quote**

This draft is a work in progress, does not reflect consensus advice or recommendations, has not been approved by the chartered SAB, and does not represent EPA policy.

1 nutrients and chlorophyll based on previous climatic conditions and will need to be considered in  
2 triennial reviews of nutrient standards.

3  
4 ***Threshold Changes***

5 When setting nutrient criteria, the Panel recommends that EPA recognize that threshold  
6 changes could occur in these systems. We include here non-linear responses, lags relative to  
7 input changes and general “state changes”. These changes will in part be a result of changing  
8 nutrient loading and freshwater discharge dynamics due to changing anthropogenic activities in  
9 watersheds and airsheds. They will also be modulated by climatic changes, including changes in  
10 rainfall (and conversely drought) intensities, frequencies and geographic patterns, as well as  
11 temperature changes (i.e., warming, which will favor the growth and proliferation of nuisance  
12 taxa such as cyanobacteria). These changes need to be considered during future triennial water  
13 quality standards reviews.

14 **3.3. Florida Coastal Waters**

15 **3.3.1. Delineation and Data Sources**

16 *Charge Question 3(a). Are the data sources identified in Sections 2.4, 4.1.1 and 4.2*  
17 *appropriate for use in deriving numeric criteria in Florida’s coastal waters? Is the SAB*  
18 *aware of additional available, reliable data that EPA should consider in delineating*  
19 *coastal waters or deriving criteria for coastal waters? Please identify the additional data*  
20 *sources.*

21  
22 The EPA document defines the outer boundary of the coastal zone as extending 3 nautical  
23 miles from shore. However, Florida’s territorial waters extend out to 3 nautical miles on the  
24 Atlantic coast and 9 nautical miles on the Gulf coast. Thus, additional data points could be  
25 included in the analysis and the Panel recommends that EPA also consider monitoring remotely  
26 sensed chlorophyll in waters further from shore. Given the dynamic nature of algal blooms in  
27 the Gulf of Mexico in particular, it is possible, and perhaps even likely, that blooms that form  
28 further than 3 miles offshore will migrate toward the coastline, thus eventually "appearing" in the  
29 state’s coastal waters. It will be important to understand the source of such patches of elevated  
30 chlorophyll, and to determine whether they are found in close proximity to the shoreline because  
31 of land- and estuary-derived nutrients or formed offshore.

32  
33 Restricting the offshore boundary to 3 nautical miles greatly reduced the number of  
34 calibration samples compared to the available data. As there is no clear boundary in water types  
35 at three nautical miles, it is appropriate to use data from the entire shelf. Extending the outer  
36 boundary to the shelf break in this way will improve the quality of the dataset. According to  
37 EPA personnel, adding these additional data increased both the correlation and the slope of the  
38 calibration graph (Fig. 4.5 in the EPA document) considerably (i.e.,  $r^2$  increased from 0.52 to ~  
39 0.8). EPA might consider using anomalies relative to either seasonal or annual means, rather  
40 than absolute Chl-*a* concentrations, in their estimates (see Stumpf et al. 2003, 2009; Tomlinson  
41 et al. 2004). This approach will mitigate problems inherent in working close to the coast, as  
42 bottom backscatter reflectance, for example, will be constant and therefore disappear from the  
43 equation.

**Science Advisory Board Panel Draft Report (dated April 8, 2011) - Do not Cite or Quote**

This draft is a work in progress, does not reflect consensus advice or recommendations, has not been approved by the chartered SAB, and does not represent EPA policy.

1 The coastal segmentation scheme suggested in the EPA document apparently is a result  
2 of historical precedence, rather than any underlying scientific rationale. Given the general  
3 alongshore flow that creates anisotropy with strong gradients perpendicular to the coast and  
4 weaker gradients parallel to it, EPA may wish to consider segments defined in terms of  
5 bathymetry.  
6

7 Another recurrent topic in panel discussions was the “missing kilometer” at the coast  
8 where Chl<sub>RS-a</sub> data are not being used because the satellite chlorophyll estimate is corrupted by  
9 the presence of land within the pixels and because of backscatter from shallow water. A  
10 potential solution may be to use turbidity data to connect conditions in the estuary proper with  
11 the coastal system just offshore, thus bridging the km gap. Another potential solution would be  
12 to collect airborne spectrographic imagery, but this would require a new data collection scheme.  
13

14 The Panel agrees that the use of remotely sensed data to develop a reference criterion for  
15 Chl-*a* is appropriate and sensible for this large, poorly sampled region. The use of these data,  
16 however, requires calibration with *in situ* chlorophyll samples, of which there are few. The  
17 Panel accepts that these sources are limited, but felt that additional sampling, including  
18 opportunistic sampling (using ferries, fishing and charter boats, etc.), where feasible, would  
19 improve the dataset. While the use of a reference criterion (Chl-*a*) is reasonable, the Panel is  
20 concerned with the sole reliance on a surrogate (see below) with no direct measurements of  
21 nutrients being made.  
22

23 The question also arises as to what reference level is applicable in this region. Historic  
24 nutrient concentrations were likely very different from today (although few data are available to  
25 provide quantitative information), yet the document assumes that these areas are currently  
26 supporting a balanced phytoplankton community. Although the Panel recognizes that a longer  
27 data record is not available, it is not clear that the ten-year dataset available from satellite  
28 observations constitutes an adequate baseline, given decadal-scale variability.  
29

30 The Panel notes that reliance on satellite observations may not be as feasible in the future,  
31 as existing sensors near the end of their lifetime. The Panel’s concerns about the continued  
32 availability of SeaWiFS data were born out by the announcement in February 2011 that after  
33 several months of attempts to communicate with the spacecraft, GeoEye has determined that the  
34 SeaWiFS mission is no longer recoverable (per a notice posted on NASA’s Ocean Color website,  
35 at [http://www.nasa.gov/vision/earth/lookingatearth/seawifs\\_10th\\_feature.html](http://www.nasa.gov/vision/earth/lookingatearth/seawifs_10th_feature.html)). While VIIRS  
36 may be launched in time to avoid disruption in the collection of ocean color data, there is also  
37 question about that sensor’s capability to produce high quality data for chlorophyll. Therefore,  
38 the Panel recommends that the EPA ensure that data from the existing U.S. and European  
39 satellites, as well as future sensors, be cross-calibrated to ensure as complete a data record as  
40 possible.  
41

1 **3.3.2. Assessment Endpoints**

2 *Charge Question 3(b). Is the assessment endpoint identified in Section 4.2 (chlorophyll-a*  
3 *to measure balanced phytoplankton biomass and production) appropriate to translate*  
4 *Florida's narrative nutrient criteria (described above) into numeric criteria for Florida's*  
5 *coastal waters, given currently available data? Does the SAB suggest modification or*  
6 *addition to this assessment endpoint?*

7  
8 EPA's suggested reference-based approach with satellite remote sensing of Chl-*a* (Chl<sub>RS</sub>-  
9 *a*) to derive numeric values is likely to be effective in Florida coastal waters, because they are  
10 optically amenable to remote sensing of chlorophyll, color (CDOM) and turbidity (Hu et al.  
11 2005; Muller-Karger et al. 2005; Palandro et al. 2004). Remote sensing technology has evolved  
12 sufficiently to begin using calibrated imagery for estimating chlorophyll.

13  
14 The Panel acknowledges that Chl<sub>RS</sub>-*a* is the most feasible indicator of nutrient status for  
15 coastal waters, given available data and the lack of a better alternative (i.e., direct monitoring of  
16 nitrogen and phosphorus offshore). However, we caution that Chl-*a* levels in these waters also  
17 are influenced by seasonal water temperatures, circulation and mixing, and influx of nutrient-rich  
18 waters from advection or upwelling. Thus, there may be only a weak relationship between  
19 nutrient concentrations and chlorophyll concentrations. There is certainly a potential relationship  
20 between nutrients and organic carbon production, but the strength of this relationship can vary  
21 depending on factors such as season or relative availability of N and P, as shown clearly in Fig.  
22 4.4 of the EPA document. Walker and Rabalais (2006), cited in the EPA document, found only  
23 about 40% of the variance in phytoplankton production could be ascribed to nutrient  
24 concentration, and this was in an area of the northern Gulf of Mexico known to be affected  
25 strongly by nutrient inputs from the Mississippi River. Also, the carbon:chlorophyll ratio within  
26 phytoplankton can vary by an order of magnitude (Banse 1977), while *Trichodesmium* blooms  
27 can arise in low N regimes because these organisms are nitrogen fixers. The Panel notes that  
28 Chl-*a* will not be useful as an indicator of species composition, as has been discussed earlier.

29  
30 **3.3.3. Approaches**

31 *Charge Question 3(c). Does the approach EPA describes in Section 4.2 appropriately*  
32 *apply remote sensing data to ensure attainment and maintenance of balanced natural*  
33 *populations of aquatic flora and fauna in Florida's coastal waters? If not, please*  
34 *provide an alternate methodology utilizing available reliable data and tools, and*  
35 *describe the corresponding advantages and disadvantages.*

36  
37 EPA is considering a reference condition approach for coastal waters. The Panel  
38 recognizes the problems with other approaches, but notes that the reference condition approach is  
39 being undertaken with no linkage to nutrient input into the coastal zone. The Panel recommends  
40 that a preliminary assessment of nutrient inputs be undertaken to better understand chlorophyll  
41 levels in the coastal zone (i.e., to relate observed chlorophyll levels in coastal waters to TN/TP  
42 concentrations or loadings from land). A first pass calculation might consist of an estimate of  
43 the total nitrogen and total phosphorus released into the coastal zones from all sources.

**Science Advisory Board Panel Draft Report (dated April 8, 2011) - Do not Cite or Quote**

This draft is a work in progress, does not reflect consensus advice or recommendations, has not been approved by the chartered SAB, and does not represent EPA policy.

1 The Panel notes the thorough approach to calibration, but has several recommendations  
2 for consideration:  
3

- 4 • According to the document, EPA used a 3 x 3 km pixel matrix and used only *in situ*  
5 calibration data taken within 3 hours of the satellite overpass. This should be sufficient,  
6 as tidal current velocities, particularly off the Florida panhandle and over the wide West  
7 Florida shelf, are generally small (< 10 cm/s; He and Weisberg 2002). Largest values are  
8 about 20 cm/s in the vicinity of the Big Bend and Florida Bay. Tidal ellipses here tend to  
9 be perpendicular to the bathymetry except very close to the coast, where they tend to  
10 parallel it (Koblinsky 1981; He and Weisberg 2002).
- 11 • The Panel recommends that satellite data within a larger, “coastal” context be used for  
12 the calibration, i.e., including data from outside the 3-mile zone. Because the calibration  
13 presented was not strong ( $r^2 = 0.52$ ), this inclusion of additional data should improve the  
14 predictive power of the model. The Panel also recommends that EPA cross-calibrate its  
15 imagery from multiple existing (and future) sensors so that  $Chl_{RS-a}$  values can be  
16 compared over time.
- 17 • Another issue on calibration concerns the relation between remotely sensed chlorophyll  
18 and water column measurements. EPA calibrated the satellite data to chlorophyll  
19 measured in the uppermost two meters of the water column. The ratio between the  
20 chlorophyll concentrations in the upper two meters and the full euphotic zone needs to be  
21 established for the subtropical waters around Florida. In many cases, there is not a strong  
22 relationship between total water column chlorophyll and near-surface values, particularly  
23 in areas with upwelling and other circulation complexity.
- 24 • The Panel recommends that obvious antecedent bloom data points be removed from  
25 analyses as these are likely not representative of desired “reference conditions” (p. 83,  
26 paragraph 1, regarding *Karenia* blooms).

27 **3.4. South Florida Inland Flowing Waters**

28 **3.4.1. Rationale for Criteria**

29 The Panel recognizes the considerable time and effort that has been put into identifying  
30 data sources, assessing endpoints, and developing two approaches for deriving nutrient criteria  
31 for inland flowing waters of South Florida. However, waters included in this category are  
32 dominated by man-made canals and the Panel is not convinced from the material provided that  
33 nutrient criteria are appropriate for these uniquely artificial and highly managed ecosystems. We  
34 identify five specific concerns in this introductory section before addressing the specific charge  
35 questions. We acknowledge that these comments and questions may not have explicit answers;  
36 however, they deserve thought and consideration.  
37

- 38 • The category of South Florida inland flowing waters includes a variety of different water  
39 types, including both human-created/human-controlled canals and natural streams (the  
40 latter of which are largely confined to the southwest region of South Florida; Tom  
41 DeBusk, personal communication; Hecker and Young 2011). A rough estimate from the  
42 SFWMD (Garth Redfield, personal communication) indicated that approximately 90  
43 percent of the waters in this region are classified as canals, and the remaining 10 percent

**Science Advisory Board Panel Draft Report (dated April 8, 2011) - Do not Cite or Quote**

This draft is a work in progress, does not reflect consensus advice or recommendations, has not been approved by the chartered SAB, and does not represent EPA policy.

1 are natural streams. Although canals dominate in number, the range of water types  
2 included in this category make it unlikely that there will be a consistent response to  
3 nutrients.  
4

- 5 • South Florida's inland flowing waters have a long history of being highly manipulated  
6 and managed, and in this regard they represent a special challenge to developing numeric  
7 nutrient criteria. The underlying problem is that the canals are classified as Class III  
8 waters, although their primary purpose is management of water quantity to alleviate  
9 flooding potential. Specific concerns include the difficulty in determining what  
10 constitutes an appropriate reference condition for these systems, and the related issue of  
11 whether or not appropriate data are available to help define reference conditions.  
12
- 13 • Internal nutrient loading from sediment accumulated in canals may confound the  
14 relationship between nutrient criteria and system response if sediments are a major source  
15 of nutrients (and based on SFWMD [2010], sediment accumulation and P mass are quite  
16 variable).  
17
- 18 • It is unclear how nutrient sources from past activities (i.e., legacy nutrients), including  
19 legacy losses or inputs of N and P to water bodies, will be accounted for in the proposed  
20 approach. If no water quality improvement or indicator biological response is seen after  
21 numeric nutrient criteria are put in place, is this because (1) the nutrient criteria are not  
22 stringent enough, (2) legacy nutrient inputs are an increasingly significant contributor, or  
23 (3) the monitoring interval is not long enough to capture the response of dynamic  
24 ecosystems and watersheds? How will continued legacy inputs of N and P be  
25 distinguished from decreased loadings related to management changes? Internal  
26 recycling of nutrients can mask water quality improvements brought about by nutrient  
27 reductions resulting from land management changes. Given the role of legacy nutrients  
28 in influencing water quality in these systems, an adaptive management approach (e.g., as  
29 part of the triennial review of water quality standards) is needed to incorporate new  
30 monitoring data and revise criteria or loading targets as appropriate.
- 31 • South Florida inland flowing waters involve a spatially and temporally dynamic  
32 interaction between surface and groundwater flows and as such, biological condition of  
33 these waters may be more responsive to hydrology than to nutrients. For instance, N and  
34 P loadings at different times of the year can influence biotic responses in different ways.  
35 In other words, it is not just how much or in what concentration, but at what time these  
36 loadings occur. In dry years, ground water will greatly influence surface water  
37 chemistry/quality compared with wet years. There also is concern that cross watershed /  
38 ecoregion / system transfers of water and nutrients in ground waters could confound the  
39 ability to relate ecological response to water column nutrient concentrations or loadings.  
40

1 **3.4.2. Delineation and Data Sources**

2 *Charge Question 4(a). Are the data sources identified in Section 2.4 and 5.4 appropriate*  
3 *for use in deriving numeric criteria in South Florida's inland flowing waters (as*  
4 *discussed in Chapters 2 and 5)? Is the SAB aware of additional available, reliable data*  
5 *that EPA should consider in delineating or deriving criteria for South Florida's inland*  
6 *flowing waters? Please identify the additional data sources.*  
7

8 The Panel recommends that the category of South Florida inland flowing waters be  
9 subdivided into two subclasses, each with separate approaches for nutrient criteria development:  
10 (1) human-managed canal systems (or some equivalent term) and (2) natural flowing lotic  
11 systems. EPA should identify characteristics to separate these two classes of water bodies, with  
12 one suggested characteristic being whether their flow regime is natural or managed.  
13

14 The human-managed canal systems should be viewed as connecting channels from inland  
15 water sources to downstream estuaries, rather than for any valued ecological attributes that may  
16 be unique to them. This would be consistent with the canal science summary document  
17 (SFWMD 2010), which describes the aquatic life in the canals (macroinvertebrates, fish,  
18 alligators), but acknowledges that the ecological value of the canals is secondary to their use for  
19 water conveyance. These canals serve as a water conduit, but also transport considerable  
20 nutrient loads to downstream receiving areas. Hence, the nutrient content in the canals can serve  
21 as a proxy for "potential impact" to the more natural wetlands and water bodies adjacent to and  
22 downstream from canals and may be best addressed through downstream protection values  
23 (DPVs) for estuaries, rather than by establishing canal nutrient IPV's. (However, see section  
24 3.6.2 on DPV).  
25

26 The Panel recommends that EPA evaluate the appropriateness of developing nutrient  
27 criteria for South Florida natural flowing lotic systems, given their greater potential for valued  
28 ecological attributes relative to canal systems. Suggestions on approaches for nutrient criteria  
29 development are provided in the next section.  
30

31 There is considerable debate as to whether or not the data in Sections 2.4 and 5.4 of the  
32 EPA document are sufficient to derive numeric criteria for South Florida's inland flowing  
33 waters. These data sources are certainly the most logical beginning point. However, EPA  
34 should look into datasets potentially available from the local water/drainage districts (not water  
35 management districts), such as Lake Worth; these local regulatory authorities can control water  
36 regulation and were created for the purpose of reclaiming the lands within their boundaries and  
37 to provide water control and water supply for settlement and agriculture (see [www.lwdd.net](http://www.lwdd.net)). In  
38 addition, agricultural interests that border the canals (e.g., U.S. Sugar Corp.) may have useful  
39 data, although these data may be proprietary.  
40

41 The proposed inventory of inland flowing waters that catalogues and distinguishes  
42 natural streams and canals should provide very useful information. This inventory could provide  
43 the basis for dividing inland flowing waters into two subcategories for purposes of criteria  
44 development, as recommended above.  
45

**Science Advisory Board Panel Draft Report (dated April 8, 2011) - Do not Cite or Quote**

This draft is a work in progress, does not reflect consensus advice or recommendations, has not been approved by the chartered SAB, and does not represent EPA policy.

1 The EPA document explores the use of the Landscape Development Index (LDI) as a  
2 potential approach (and data source) for determining reference conditions in inland flowing  
3 waters (reference conditions where  $LDI < 2$ , p. 105). It is well established that surrounding land  
4 use can have substantial impacts on receiving water bodies, so this approach has conceptual and  
5 intuitive appeal. However, insufficient information is available in the EPA document to  
6 determine the appropriateness of the LDI approach for South Florida's inland flowing waters.  
7 The Panel does not recommend that the LDI be applied to South Florida canals. Further concerns  
8 include the use of a 100-m buffer along canals; would not the canal's water quality to be  
9 determined by the entire area that drains into it? The document cites a study by Fore (2004) to  
10 justify this approach. However, Fore (2004) was based on streams throughout the state and not  
11 just canals; there are considerable differences in hydrology and land-water interactions between  
12 canals and natural stream channels. The 100-m buffers proposed for use with the LDI (p. 105-  
13 106) may be too limited, particularly where stormwater pipes convey runoff from distances much  
14 further than 100 meters.

15  
16 The condition of inland flowing waters is highly influenced by geology and  
17 anthropogenic activity. In this regard, there is logic to subdividing these waters according to  
18 basin and sub-basin soil types and land uses. An additional challenge is incorporating  
19 groundwater hydrologic/nutrient dynamics, which also have been altered, but are likely to be  
20 very important in determining nutrient sources and impacts. The proposed classification scheme  
21 appropriately incorporates surface and subsurface flow regimes and flow lines, as well as soil  
22 types and human agricultural and urban impacts (i.e., land use). Inland water regions should be  
23 classified according to soil order, color of water, or a combination of several such characteristics.

24  
25 As mentioned above, legacy N and P loads from past management and from natural  
26 sources also must be considered. The EPA document appears to minimize the importance of  
27 legacy effects of past management (e.g., the statement on the top of page 40 in reference to  
28 Huang and Hong 1999). However, soil nutrient levels vary greatly as a function of past  
29 management, within and among fields and within areas with uniform LDIs for which specific  
30 criteria concentrations will be set. Additionally, geologic rock deposits vary within areas  
31 assumed to have similar nutrient concentrations. There is a wealth of data on soil nutrient levels  
32 (particularly P) available from NRCS and land-grant university extension offices. EPA should  
33 evaluate these, and other, data to better support the regional classification of South Florida inland  
34 flowing waters.

35  
36 **3.4.3. Assessment Endpoints**

37 *Charge Question 4(b). Are the assessment endpoints identified in Section 5.4 (balanced*  
38 *faunal communities, i.e., aquatic macroinvertebrates, and balanced phytoplankton biomass*  
39 *and production) appropriate to translate Florida's narrative nutrient criteria (described*  
40 *above) into numeric criteria for South Florida's inland flowing waters, given currently*  
41 *available data? Does the SAB suggest modification or addition to these assessment*  
42 *endpoints?*

43  
44 Philosophically (but with practical implications), one can question whether any  
45 assessment endpoint is appropriate for systems that have been artificially created. How does one  
46 establish an appropriate reference condition for such systems, especially when they are heavily

**Science Advisory Board Panel Draft Report (dated April 8, 2011) - Do not Cite or Quote**

This draft is a work in progress, does not reflect consensus advice or recommendations, has not been approved by the chartered SAB, and does not represent EPA policy.

1 managed? There are no easy answers for these questions, although this has certainly been done  
2 for reservoirs. Furthermore, nutrient inputs are not the main drivers of habitat quality in the  
3 canals of South Florida.  
4

5 South Florida canals have been constructed continuously over the last century, so it is not  
6 clear how reference conditions can be assessed for these highly variable systems that are  
7 designed to be dynamic and flashy during wet periods (to get water off the landscape quickly)  
8 but are managed during dry periods to be slow or non-flowing. Least-disturbed sites tend to be  
9 in one region only and may not be transferable to other identified regions. Because canals are  
10 unique aquatic ecosystems, more information needs to be presented on how balanced natural  
11 populations are to be assessed. An initial inventory of science for South Florida canals, provided  
12 by SFWMD (2010), summarizes data on water quality and biological conditions in the canals.  
13 The closest analog to South Florida canals would be in The Netherlands, where much of the  
14 inland waters flow through canals (locally called ditches). There is some literature on  
15 assessment endpoints from Netherland ditches (e.g., see Verdonshot 1987) that may be of use in  
16 developing methods for assessing the status of flora and fauna in Florida canals.  
17

18 If nutrient criteria are to be developed for South Florida inland flowing waters, the Panel  
19 recommends further consideration and assessment of both the response variables and the form of  
20 the nutrients being assessed. For example, distinction is needed among nutrient forms that are  
21 immediately availability for biological uptake —i.e., short-term bioavailability and growth  
22 response, such as inorganic nitrogen ( $\text{NO}_3$  and  $\text{NH}_4$ ) and phosphorus ( $\text{PO}_4$ )—compared with  
23 losses, such as particulate and organic forms of N and P (i.e., long-term availability).  
24

25 For assessment endpoints, both Chl-*a* and invertebrate measures have conceptual  
26 appeal, but their utility is not straightforward. Some freshwater and estuarine ecosystem studies  
27 have shown that Chl-*a* can be a function of grazing pressures rather than, or in addition to,  
28 nutrient concentrations. For example, increasing nutrient concentrations in inland flowing  
29 waters can increase the number of grazers, which can lead to a lower Chl-*a* concentration; i.e., a  
30 top-down regulation of primary production (Cloern 1982; Cohen et al. 1984; Hilton et al. 2006;  
31 Thompson et al. 2008; Lewis and McCutchan 2010). Aquatic macroinvertebrate community  
32 structure and/or traits have been shown to be reliable bioindicators in other aquatic ecosystems.  
33 Hence, they are a reasonable starting point for South Florida's inland flowing waters. However,  
34 these systems are poorly understood, highly managed, and heavily modified. As a consequence,  
35 it is unclear at present if these proposed assessment endpoints can be applied effectively.  
36

37 The Panel identified areas of uncertainty that need further attention before a reasonable  
38 level of scientific confidence can be applied to the use of balanced faunal communities and/or  
39 balanced phytoplankton biomass and production. We elaborate on these below.  
40

#### 41 ***Faunal Communities***

42 The macroinvertebrate index used by Snyder et al. (1998), provided as Figure 5-8 in the  
43 EPA document, shows a good relationship between land use and macroinvertebrate community  
44 structure. However, macroinvertebrate data provided in a presentation to the Panel reveal a  
45 much more tenuous stressor-response relationship between total P concentrations and  
46 macroinvertebrate indices (DeBusk 2010). It is important that EPA examine possible reasons for

**Science Advisory Board Panel Draft Report (dated April 8, 2011) - Do not Cite or Quote**

This draft is a work in progress, does not reflect consensus advice or recommendations, has not been approved by the chartered SAB, and does not represent EPA policy.

1 the lack of correspondence in these two data sets. Possible explanations include the use of  
2 different measures of stressor (land use vs. total P), different types of indices (e.g., emphasizing  
3 different taxa), or inclusion of inland flowing waters from different parts of south Florida (e.g.,  
4 that experience different pressures). For example, the relatively high Stream Condition Index  
5 (SCI) score for the wetland sites shown in the Snyder et al. data may have more to do with  
6 habitat quality than nutrients. The summary of canal science prepared by the SFWMD (2010)  
7 notes that “additional research is needed to select sensitive (macroinvertebrate) metrics and a  
8 quality threshold applicable to low gradient streams and canals within the peninsula and  
9 Everglades bioregions.” The Panel agrees with this statement. If the different macroinvertebrate  
10 patterns in these data sets can be explained and related to nutrients, aquatic macroinvertebrates  
11 may be a very useful assessment endpoint and one that the Panel recommends be given more  
12 attention. However, that relationship has not yet been established.

13 ***Phytoplankton Biomass***

14 There is a relative paucity of phytoplankton data (either as Chl-*a*, species composition, or  
15 productivity) in these inland flowing waters. Data from SFWMD (2010) show geometric mean  
16 Chl-*a* concentrations ranging from 2 µg/L (Lower East Coast) to 8.0 µg/L (Everglades  
17 Agriculture Area) in canals within the South Florida region. However, these concentrations are  
18 not paired with data on hydrologic conditions, and it is impossible to assess if they represent  
19 actively growing algae populations (as might be expected in a non-flowing canal) or algae being  
20 transported downstream (i.e., in a flowing canal) and therefore not representative of local  
21 conditions. The hydrologic status of the canal (non-flowing, slow-flowing, fast-flowing) has  
22 enormous implications for the plankton community, and this needs to be accounted for in EPA’s  
23 assessment. In addition, both present and antecedent hydrologic conditions need to be  
24 considered in any analysis, as the flow regimes are quite variable (and can be artificially  
25 controlled) in these systems. Hence, a canal that may be flowing when sampled may have been  
26 stagnant for weeks prior to sampling, and vice versa. These kinds of lags make relationship  
27 interpretation (and accurate modeling) complex. At this point, it is unclear if there are sufficient  
28 data to know what a “protective” level of Chl-*a* should be for these systems; as a consequence, it  
29 is currently not possible to assess whether or not phytoplankton can be used as an effective  
30 assessment endpoint.

31  
32 The inventory of the inland flowing waters, and subsequent screening of water bodies, is  
33 an important step and may help in the selection of appropriate assessment endpoints. The  
34 approach provided in the technical document is a good starting point, but the Panel has identified  
35 some issues and suggestions with respect to the classification procedure. The Panel identified  
36 several factors that EPA may want to consider with respect to the chlorophyll endpoint:

- 37
- EPA proposes a classification of inland water regions according to soil order, land management systems or color of water (see below). Some combination of these factors should be considered, taking into account covariates.
  - Currently, EPA does not appear (at least explicitly) to consider the potential influence of humic soils in their classification of inland flowing water types, with respect to their role in discoloration of waters; phytoplankton response will be very different in waters that are naturally colored (i.e., influenced by humics) vs. those that are not.
- 38  
39  
40  
41  
42  
43

**Science Advisory Board Panel Draft Report (dated April 8, 2011) - Do not Cite or Quote**

This draft is a work in progress, does not reflect consensus advice or recommendations, has not been approved by the chartered SAB, and does not represent EPA policy.

1 Dissolved oxygen concentration reflects the relative amount of photosynthesis (DO  
2 production) and respiration (DO consumption) in aquatic ecosystems, so DO might be an  
3 alternative endpoint in this geographic domain; however, new studies would be needed to  
4 determine if DO levels are linked to nutrient loads or concentrations, and not to other factors  
5 (such as light), and if groundwater influx (low DO) confounds the use of this assessment  
6 endpoint.  
7

8 **3.4.4. Approaches**

9 *Charge Question 4(c). EPA describes two approaches in Section 5.4 (reference conditions*  
10 *and stressor-response relationships) for deriving numeric criteria in South Florida inland*  
11 *flowing waters. Compare and contrast the ability of each approach to ensure attainment and*  
12 *maintenance of balanced natural populations of aquatic flora and fauna in different types of*  
13 *flowing water or geographical areas, given currently available data?*  
14

15 The two approaches that EPA is considering for determining numeric criteria for South  
16 Florida inland flowing waters are discussed in Section 5.4 of the EPA document. The first is  
17 based on reference conditions and the second is based on stressor-response relationships. The  
18 Panel recommends against the application of either approach for the human-controlled canal  
19 systems, as these inland waters are more appropriately thought of as hydrologic conduits  
20 designed to rapidly remove storm water from the landscape, while at the same time delivering  
21 nutrients to downstream estuaries. The Panel does recommend that EPA evaluate both  
22 approaches for the natural flowing lotic systems, but recognizes that data availability may  
23 severely limit their ability to develop rigorous criteria at present. As such, the Panel  
24 recommends that, at least for the present, South Florida inland flowing waters be addressed  
25 through downstream protective values. We comment on the problems with EPA's proposed  
26 approaches below.  
27

28 ***Reference Conditions***

29 Briefly, under the approach based on reference conditions, a set of least-disturbed sites  
30 would be identified using the Land Development Intensity (LDI) index. The total LDI for each  
31 site would be calculated as an area-weighted sum of the LDI coefficients for all land uses within  
32 an area of influence. Sites with total LDI below 2.0 or another specified threshold would be  
33 classified as least-disturbed and would form the reference set. The historical annual values of  
34 TN and TP for these sites would be used to fit lognormal distributions of TN and TP under least  
35 disturbance and specified quantiles of these fitted distributions – the EPA document mentions the  
36 0.75 and 0.90 quantiles – would be used as the numeric criteria.  
37

38 On the bottom of page 107, the EPA document discusses the question of the frequency  
39 with which these numeric criteria could be exceeded. This discussion is difficult to follow, but  
40 the general point appears to be this: Consider that the estimated 0.75 quantile for one nutrient is  
41 exceeded  $k$  or more times in  $n$  years. Commonly used values are 1 in 3 and 2 in 5. Under the  
42 assumption that values in different years are independent and have the same distribution as the  
43 reference set (and ignoring any error in the estimation of the 0.75 quantile), the probability of  
44 this event is given by:  
45

**Science Advisory Board Panel Draft Report (dated April 8, 2011) - Do not Cite or Quote**

This draft is a work in progress, does not reflect consensus advice or recommendations, has not been approved by the chartered SAB, and does not represent EPA policy.

$$p(k, n) = \sum_{j=k}^n \binom{n}{j} 0.25^j 0.75^{n-j}$$

For fixed  $k$  and  $n$ , this formula essentially provides a one-sided significance level for testing the null hypothesis that the nutrient distribution is the same as that of the least-disturbed sites. So, for example, under this null hypothesis, the probability  $p(1, 3)$  of at least 1 exceedance of the 0.75-quantile in 3 years is 0.58, while the probability  $p(2, 3)$  of at least 2 exceedances in 3 years is 0.16.

The Panel notes the following:

- The choice of quantile,  $k$ , and  $n$  can have a profound effect on the performance of criteria derived in this way and some discussion is needed about how this choice will be made.
- The probability calculations sketched here pertain to exceedances of a single nutrient criterion. If the same rule is applied to both nutrients (and assuming that nutrient levels are independent) then, under the null hypothesis that both nutrient distributions are the same as those for least-disturbed sites, the probability that either or both nutrients exceed their respective 0.75-quantiles in at least  $k$  of  $n$  years is  $1 - (1 - p(k, n))^2$ . Thus for both nutrients, the probability of at least 1 exceedance of the 0.75-quantile of either or both nutrients in 3 years is 0.82, while the probability of at least 2 exceedances in 3 years is 0.29, i.e., much greater than for each nutrient considered individually

As noted, these calculations assume that the relevant quantile of the annual nutrient levels in least-disturbed sites are estimated without error. An assessment of the impact of estimation error – including non-normality of the log of annual nutrient levels – on the accuracy of these calculations is needed.

Although these calculations provide information about the rate of Type I error (i.e., the exceedance of the criterion when the underlying distribution is the same as that for least-disturbed sites), they provide no information about the rate of Type II error (i.e., the non-exceedance of the criterion when the underlying distribution is different from that for least-disturbed sites). In the jargon of hypothesis-testing, this analysis provides no information about power. To gain such information, it is necessary to consider also the distribution of nutrients in disturbed sites.

To some extent, variability of nutrient levels in the least-disturbed sites will reflect heterogeneity in hydrology, geology, etc. Failure to account for such heterogeneity, which is also present in disturbed sites, may result in numeric criteria that are under- or over-protective for some sites. It would, therefore, seem preferable to develop criteria that account for such factors. The EPA document briefly notes (on p. 108) that EPA also is considering following the reference conditions approach using all sites as a reference set (and not only least-disturbed sites as discussed above). With the exception of the identification of least-disturbed sites, the mechanics of these approaches are the same. However, the underlying logic seems rather different – loosely speaking, one approach aims to reproduce conditions in least-disturbed sites

**Science Advisory Board Panel Draft Report (dated April 8, 2011) - Do not Cite or Quote**

This draft is a work in progress, does not reflect consensus advice or recommendations, has not been approved by the chartered SAB, and does not represent EPA policy.

1 and the other aims to maintain conditions within a specified quantile of the distribution of all  
2 sites, whatever their level of disturbance – and this needs to be discussed.

3  
4 ***Another Possible Approach***

5 The Panel briefly discussed another statistical approach to determining numeric criteria  
6 for TN and TP. The ultimate goal of this determination is to ensure that waterbodies meet their  
7 designated uses. If information is available about whether or not waterbodies with the same  
8 designated use are, in fact, meeting this use, then it should be possible to construct logistic or  
9 other binary regression models relating the probability that designated use is met to levels of  
10 nitrogen and phosphorus (and possibly to other regressors such as rainfall).

11  
12 Consider that for years Florida DEP scientists examined water quality data on Florida  
13 surface waters and then made the judgment as to whether or not a waterbody was in compliance  
14 with the narrative nutrient criteria. In effect, these scientists created a binary response variable  
15 (compliance or noncompliance) for each “data row” of water quality data for each Florida  
16 waterbody. These data rows for each waterbody type (e.g., lakes, rivers, estuaries) could be the  
17 basis for development of logistic or other binary regression models with the water quality  
18 variables as predictors and the DEP scientists’ classifications as the response.

19  
20 For example, if initial analyses indicated that TN and TP were clearly the best predictors  
21 of compliance response in estuaries, then a binary regression model could be fitted with TN and  
22 TP as the two predictors of the compliance response. Then new TN and TP data could be  
23 inserted into this binary regression model to determine the probability of compliance with the  
24 nutrient criteria. So, for example, with a 50:50 cutoff for compliance as the transition point, if an  
25 estuary’s TN and TP result in greater than 50% probability of compliance with the nutrient  
26 criteria, the estuary would be classified as not impaired.

27  
28 This approach can be undertaken with little cost and time, and it makes direct use of the  
29 State’s interpretation of the existing narrative criteria. This approach is similar in spirit to the  
30 one based on reference conditions. However, unlike the approach based on reference conditions,  
31 the statistical analysis discussed here would use information about both unimpaired and impaired  
32 cases. Although the results of this statistical analysis will be only as good as the underlying  
33 categorization of waters as impaired or unimpaired, the approach may offer additional insight for  
34 interpreting proposed criteria.

35  
36 ***Stressor-Response Relationships***

37 The second approach that EPA is considering for developing numeric criteria for South  
38 Florida inland flowing waters is based on stressor-response relationships. This involves  
39 developing a statistical model relating the level of Chl-*a* to TN or TP. The EPA document  
40 presents examples involving linear, nonparametric, and quantile regressions of log Chl-*a* (as  
41 response) and log TN or log TP (as stressor).

**Science Advisory Board Panel Draft Report (dated April 8, 2011) - Do not Cite or Quote**

This draft is a work in progress, does not reflect consensus advice or recommendations, has not been approved by the chartered SAB, and does not represent EPA policy.

1 The Panel notes the following:  
2

- 3 • A fundamental question that the EPA document leaves unanswered is how such fitted  
4 regression models will be used to determine numeric criteria; i.e., how they will  
5 determine the level of Chl-*a* that will be considered protective of balanced phytoplankton  
6 and faunal communities. This is a serious shortcoming that needs to be addressed.
- 7 • As the EPA document notes, it has not been possible to develop stressor-response  
8 relationships in which a biological endpoint other than chlorophyll serves as the response.  
9 However, if there is a clear relationship between Chl-*a* and TN, say, and a clear  
10 relationship between another biological endpoint (e.g., a faunal community metric) and  
11 Chl-*a*, then there would ordinarily be a clear relationship between that biological  
12 endpoint and TN. The fact that it is difficult to identify this latter relationship may reflect  
13 limitations of the statistical models considered so far. For example, the effect of TN or  
14 TP on a biological endpoint may be modulated by other factors. This effect could be  
15 obscured by omitting these factors from the regression model.
- 16 • As with the approach based on reference conditions, the relationship between Chl-*a* and  
17 TN or TP is likely to be modulated by the effects of hydrological, geological, and other  
18 covariates. Failure to account for such factors may lead to criteria that are over- or  
19 under-protective at some sites, and it would again seem preferable to include such  
20 covariates in developing numeric criteria. Furthermore, should TN and TP be considered  
21 simultaneously (i.e., is a multiple or simple regression most appropriate)?
- 22 • Some of the variability in the stressor-response relationship could be a result of season.  
23 This should be investigated, and it may lead to the formulation of different criteria for  
24 different seasons.
- 25 • A substantial amount of effort will be put into identifying and quantifying stressor-  
26 response relationships in these waters using correlative/regression analysis. Considering  
27 the difficulty of working across the surface-subsurface interfaces in deriving nutrient  
28 loading estimates, as well as effects of these loads, the authors have done a good job of  
29 addressing these challenges. This section could however benefit from closer  
30 process/response connections (including applying modeling approaches) to receiving  
31 estuarine and coastal waters.
- 32 • The results of a distribution approach (p. 108) are sensitive to the distribution of sites  
33 along the disturbance gradient. If a larger proportion of the samples are from more  
34 disturbed sites, then using the lower percentile to set the criteria will result in a higher  
35 number than if a larger proportion of the samples are from less disturbed sites. Some  
36 requirements for the distribution of sites along the disturbance gradient should be  
37 specified.

1 **3.5. South Florida Marine Waters**

2 **3.5.1. Delineation and Data Sources**

3 *Charge Question 5(a). Are the data sources identified in Section 2.4 and 5.5 appropriate*  
4 *for use in deriving numeric criteria in South Florida’s marine waters (as discussed in*  
5 *Chapters 2 and 5)? Is the SAB aware of additional available, reliable data that EPA*  
6 *should consider in delineating or deriving criteria for South Florida’s marine waters?*  
7 *Please identify the additional data sources.*

8  
9 One general recommendation is that the waters currently termed “marine waters” in the  
10 EPA document be changed to “South Florida coastal and estuarine waters” to be consistent with  
11 the use of the terms throughout the rest of the document.

12  
13 The southern part of Florida has a different nutrient regime than other parts of the state: it  
14 is highly oligotrophic and conditions can be rapidly altered by upstream water management  
15 (versus nutrient regulatory) decisions. In addition, embayments with calcareous sediments are  
16 highly phosphorus (not nitrogen) limited. The Panel agrees that these waters should be  
17 considered separately for purposes of nutrient criteria development. However, the proposed  
18 subdivision/subclassification of South Florida estuarine and coastal waters does not clearly relate  
19 to the oceanographic circulation and degree of connectivity in the region, particularly on the  
20 seaward side of the Keys and the southeast and southwest interior coasts. Alongshore circulation  
21 and interconnection is much more vigorous than across shelf except for highly restricted  
22 intertidal exchange passages (Lee and Mayer 1977; Lee et al. 2002).

23  
24 The data identified in the report seem appropriate for use in this exercise. There also are  
25 water quality data from NOAA’s Atlantic Oceanographic Meteorological Laboratory (AOML)  
26 that have been collected for Florida Bay, Biscayne Bay, the Florida Keys and SW Florida Shelf  
27 for more than a decade as part of the NOAA South Florida Program ([www.aoml.noaa.gov/sfp](http://www.aoml.noaa.gov/sfp)).  
28 There are some possibly significant differences between these data and the Southeast  
29 Environmental Research Center (SERC) data, which covers the same domains. For some  
30 periods in some subregions, the NOAA data were temporally more dense (bimonthly versus  
31 quarterly) in larger domains and the nutrient methodologies were more sensitive (long path  
32 length liquid wave guide) in accordance with oceanographic practice for oligotrophic open ocean  
33 waters (as established in JGOFS, GLOBEC and other international programs).

34  
35 **3.5.2. Approaches**

36 *Charge Question 5(b). EPA describes two methods in Section 5.6 for using a reference*  
37 *condition approach for deriving numeric criteria in South Florida marine waters (least-*  
38 *disturbed sites or bionomial test). Compare and contrast the ability of each approach to*  
39 *ensure attainment and maintenance of balanced natural populations of aquatic flora and*  
40 *fauna in South Florida marine waters, given currently available data?*

41  
42 Two approaches to nutrient criteria are being considered for South Florida coastal and  
43 estuarine waters. The first is to identify criteria that are inherently protective based on a  
44 statistical evaluation of data from least-disturbed sites. In some of these zones, least-disturbed

**Science Advisory Board Panel Draft Report (dated April 8, 2011) - Do not Cite or Quote**

This draft is a work in progress, does not reflect consensus advice or recommendations, has not been approved by the chartered SAB, and does not represent EPA policy.

1 sites may be those most distant from land-based sources; this is not a result of a significant  
2 different in nutrient loading but rather a result of dilution with naturally highly oligotrophic  
3 waters. Hence, by including sites being diluted by highly oligotrophic ocean waters, the criteria  
4 may be overly protective. Moreover at such sites it is not atypical that the limiting nutrient can  
5 change. A second proposed approach is also based on a statistical evaluation, but in this case  
6 raw data are analysed using a binomial test and two criteria are generated – an average  
7 concentration and an upper percentile concentration that is more sensitive to higher  
8 concentrations. Both approaches have merit and the Panel encourages the application of both to  
9 provide a more robust evaluation of criteria.

10  
11 A third alternative has been suggested by Briceño et al. (2010), based on statistical  
12 analysis of long-term monitoring data for Chl-*a* and nutrient concentrations in South Florida  
13 coastal and estuarine waters. While that approach has some merit in subregions where sufficient  
14 time-series data are available, it is not applicable in many other areas. Moreover, an inherent  
15 lagtime is introduced in that analysis which may make it insensitive to ecologically significant  
16 deleterious changes if they occur too rapidly.

17  
18 It is also critical to address how the statistical approaches would be applied. For  
19 example, if a baseline (i.e., reference) condition is established using the median or geometric  
20 mean of a decade of data for the undisturbed condition, there still remains the major issue of how  
21 concentrations that exceed the criteria will be determined. Will each year be assessed against the  
22 baseline (the approach taken with the CERP System Status Report and the South Florida  
23 Ecosystem Restoration Task Force, SFERTF, Scientific Indicators) or will five years of data be  
24 required to determine if 2 (or 3) had “exceeded” the baseline? How would the variability in the  
25 two data sets (baseline and evaluation) be incorporated? Given how variable some of these  
26 numbers are, it is a lot less powerful (less chance of seeing a change) to ask if the means of the  
27 two datasets (in the example above, 10 and 5 yrs) differ versus whether a particular year was  
28 significantly above the baseline mean. Furthermore, there may be major ecological differences  
29 between two successive years of concentrations that exceed the criteria versus two years  
30 separated in time, and the document does not discuss this. The Panel recommends that more  
31 thought be given to these implementation issues.

32  
33 The Panel recommends a reconsideration of the rationale for doing both a principal  
34 component and cluster analysis. EPA proposes to use a combination of principal component  
35 analysis (PCA) and cluster analysis to define coastal regions based on multivariate measurements  
36 with sites. As the goal of cluster analysis is precisely to identify groups of similar sites, it is  
37 unclear why PCA is also being proposed in this context.

38  
39 ***Past Alterations***

40 The coastal and estuarine waters of South Florida have experienced enormous changes  
41 over the last century. For example, the Florida Bay of the early 1900’s was a true estuary with  
42 low and highly variable salinities for most of the year. Following widespread damming for the  
43 Flagler railroad, salinities were lowered throughout much of the Bay; the effect on salinity was  
44 relatively minor, but the effect on residence time was significant. Then, after canal construction  
45 from the 1920s to mid-1960s, the vast majority of the water flowing out of Lake Okeechobee is  
46 now shunted out to sea before reaching the southern-most waters of the state (Steinman et al.

**Science Advisory Board Panel Draft Report (dated April 8, 2011) - Do not Cite or Quote**

This draft is a work in progress, does not reflect consensus advice or recommendations, has not been approved by the chartered SAB, and does not represent EPA policy.

1 2002b). Although there were a number of animal studies conducted, there were few nutrient or  
2 chlorophyll measurements made because the water was so clear that light penetrated to the  
3 bottom. In the 1970s, *Thalassia* covered Florida Bay, believed to be a result of the artificially  
4 high salinities resulting from the eastward and westward shunting of water that used to flow  
5 south into Florida Bay. A major drought in the mid-1980s resulted in Florida Bay salinity going  
6 as high as 70 ppt, which killed off *Thalassia* and other sea grasses. Although nutrients were not  
7 a cause of the sea grass dieoff, the result was that enormous amounts of nitrogen and phosphorus  
8 that had been sequestered as detritus in the sediment were no longer protected by the dense sea  
9 grasses. A subsequent large storm event then mixed large amounts of sediment nutrients into the  
10 water column. The result was eutrophication, yet the ultimate cause was a change in salinity that  
11 killed off sea grasses years before. Based on this brief history, the Panel has the following  
12 recommendations:

- 13  
14 • When setting reference conditions, EPA should consider historical water management  
15 and structural changes and regional climatic variability that have affected water delivery  
16 to South Florida estuaries and coastal waters, recognizing that current conditions do not  
17 reflect historical conditions.
  
- 18 • In particular, salinity should be considered a covariate because of its role in maintaining  
19 the water quality necessary for seagrasses. We note, however, that salinity is relevant  
20 (and in fact variable as a result of water management) only in a very restricted part of this  
21 domain.
  
- 22 • Seagrass coverage and the extent of epiphytic colonization should be considered as  
23 endpoints, in addition to water column chlorophyll (see also 3.2.2).

24  
25  
26 ***Geographic Areas to be Included***

27 South Florida contains a number of parks and marine protected areas. Management  
28 jurisdiction in the region has been clarified to a large degree by the formation of the National  
29 Marine Sanctuary. The document should clarify which coastal and estuarine areas will be under  
30 the jurisdiction of the EPA document under review vs. other regulations. We note that the  
31 Florida Keys National Marine Sanctuary (FKNMS) domain (and that of the three national parks:  
32 Biscayne, Dry Tortugas and Everglades – a.k.a. Florida Bay) are not the only federally protected  
33 waters, and there are also state-protected waters of various types. It is our understanding that  
34 what is set by EPA will constitute a “de minimus” standard for these areas, which could receive  
35 additional protection. Similarly, the EPA document should clarify the relationship of the South  
36 Florida coastal and estuarine nutrient criteria to the relevant indicators and performance  
37 measures already established by the Comprehensive Everglades Restoration Plan (CERP)  
38 through its Restoration, Coordination and Verification (RECOVER) program.  
39

1 **3.6. Downstream Protection Values**

2 **3.6.1. DPVs for Estuaries**

3 *Charge Question 6(a). Are the methods EPA is considering for deriving downstream*  
4 *protection values (DPVs) for estuaries (excluding marine waters in South Florida) as*  
5 *described in Section 6.1-6.4 appropriate to ensure attainment and maintenance of*  
6 *downstream water quality standards, given available data? Please describe additional*  
7 *approaches and their advantages and disadvantages that EPA should consider when*  
8 *developing numeric criteria to protect these downstream estuarine waters (excluding marine*  
9 *waters in South Florida), given available data?*

10

11 **Rationale for DPVs**

12 The 1972 Clean Water Act (CWA) states that:

13

14 In designating uses of a water body and the appropriate criteria for those uses, the  
15 State shall take into consideration the water quality standards of downstream waters  
16 and shall ensure that its water quality standards provide for the attainment and  
17 maintenance of the water quality standards of downstream waters.

18

19 This provision has been the basis for ensuring that water quality standards in one state provide  
20 for attainment and maintenance of water quality standards of downstream states and Tribes. The  
21 recently published nutrient criteria for Florida's lakes and flowing waters (75 FR75762-75807,  
22 December 6, 2010) explicitly included the concept of downstream protection values (DPV) as a  
23 concentration or loading value in a stream at the point of entry into a lake, set at a value to ensure  
24 that lake nutrient criteria are attained. The rule also notes that wasteload and/or load allocations  
25 from an approved total maximum daily load (TMDL) may be used as the DPV.

26

27 In the present document, the concept of DPV is included as a means of ensuring that  
28 upstream N and P water quality criteria will be set at levels that will protect downstream  
29 estuarine designated uses. This is particularly important because freshwater and estuarine  
30 systems may be sensitive to different nutrients. However, the Panel had concern with the  
31 overlap between DPV and the TMDL process. To illustrate this, consider Figure 2 below.  
32 Water quality criteria (WQC) are required for all waterbodies in this figure – the estuary and the  
33 streams. If streams A1, B1, and C1 meet their WQC, yet the estuary does not, additional  
34 pollutant load reductions to the estuary are required. These reductions could come from direct  
35 loading, the atmosphere, or the tributaries (A1, B1, C1). Standard practice in the TMDL process  
36 is to model the estuary and watershed to determine the additional pollutant load reduction  
37 needed, and then to allocate the load reduction based on input from state and local officials. One  
38 possible result of implementation of the required load reduction is that one or more streams may  
39 require WQC more stringent than those initially established in order to attain WQC in the  
40 estuary. Regardless, this regulatory-driven analysis achieves compliance with all WQC without  
41 the additional regulatory entity of DPV criteria.

42

43 The drawback of relying on TMDLs, rather than the DPVs, is that the TMDL process is  
44 not triggered until impairment has occurred in a waterbody. However, the TMDL process

**Science Advisory Board Panel Draft Report (dated April 8, 2011) - Do not Cite or Quote**

This draft is a work in progress, does not reflect consensus advice or recommendations, has not been approved by the chartered SAB, and does not represent EPA policy.

1 includes the opportunity to consider socio-economic factors when selecting an allocation scheme  
2 for load reduction. In principle, in an environment with no water quality impairment, the DPV  
3 concept could be captured within the designated use for each tributary and correspondingly  
4 reflected in the water quality criteria for each tributary. However, in situations where  
5 impairment exists, the DPV enforces rigidity that likely will not accommodate variations in  
6 natural and human factors affecting nutrient loading. Of particular note, nutrient trading between  
7 tributary watersheds, which might be viewed favorably from a social and an economic  
8 perspective, may be prohibited due to *a priori* nutrient load restrictions imposed by the rigidity  
9 of the DPVs.

10  
11 The EPA document includes an example of DPV developed using an equal load  
12 reduction for each tributary to a waterbody. The Panel notes that this equal allocation approach  
13 ignores the possibility of different background nutrient concentrations in upstream segments, and  
14 other characteristics of segments that would make nutrient load reductions impractical. In  
15 response to a Panel request for further information, the Agency described four options for  
16 distributing loads for calculation of DPV (Appendix C). Each of these options presents some  
17 issues, as discussed below.

- 18  
19 • **Option A** distributes the load based on discharge, making the assumption that high  
20 nutrient loads are associated with higher flow rates. This option could present several  
21 unresolvable issues. For example, a stream with a low discharge but high nutrient load  
22 may be required to reduce its load to unachievable values. This could occur if a stream  
23 drains an area with elevated natural background, and would result in criteria that would  
24 require load reductions below background. Conversely, a stream with a high discharge  
25 may not be required to reduce its load even if there were some relatively simple  
26 approaches for reducing its load.
- 27  
28 • **Option B** would distribute load based on watershed area drained. This would appear  
29 logical if the export of nutrients from all land uses were nearly equivalent, which it is not.  
30 As with Option A, this option might result in reduction targets that are unreasonable or  
31 unachievable in small subwatersheds while missing opportunities for reductions in larger  
32 watersheds. An integrated approach to load reduction should focus on watersheds, rather  
33 than on individual segments.
- 34  
35 • **Option C** would distribute load based on natural or background loading. This approach  
36 seems reasonable if it is possible to accurately arrive at the natural background nutrient  
37 load for a segment or watershed. Essentially, an estimate of the natural background for  
38 each segment, given a set of criteria (slope, soils, etc), would set the achievable “DPV”  
39 for the stream segment. Such an approach is intellectually appealing, but may not be  
40 practical in the time frame given for the project. It does, however, strongly suggest the  
41 need to have a GIS-based model such as SWAT developed for the project and later  
42 implementation.
- 43  
44 • **Option D** would distribute load based on average background, as described in Option C,  
45 or natural background, but include anthropogenic loading using existing data. The  
46 difference between natural and anthropogenic load would be the load that could be


**Science Advisory Board Panel Draft Report (dated April 8, 2011) - Do not Cite or Quote**

This draft is a work in progress, does not reflect consensus advice or recommendations, has not been approved by the chartered SAB, and does not represent EPA policy.

1 targeted for reduction. In the description of this option, it is suggested that average load  
2 for a stream/segment could be divided by the average discharge. This is the  
3 concentration, and makes one question the need to calculate loads or DPVs if we can  
4 simply express a maximum concentration for a stream or segment.  
5

6 The Panel notes a few additional points to think about when considering the wisdom of DPVs:  
7

- 8 • There are other sources, such as direct (e.g., groundwater and atmosphere) loadings to the  
9 estuary. How might those be addressed in the determination of tributary DPVs?
- 10 • Referring to Figure 2 (below), why should there not be DPVs for streams B2a and B2b in  
11 order to protect stream B1?  
12



13  
14  
15  
16  
17 ***Approach to Setting DPVs***

18 EPA's proposed assessment of DPVs is based on watershed modeling (to be undertaken  
19 using the LSPC model) which results in an apportioned pollutant load reduction for each  
20 tributary to the waterbody (e.g., estuary) of interest. In an example in the review document, EPA  
21 proposes to apportion the pollutant load reduction (required to achieve compliance with the  
22 waterbody water quality criterion) as an equal fractional load reduction for each tributary to the  
23 waterbody. As noted above, this approach would formalize, and unnecessarily restrict, the more  
24 flexible pollutant load allocation process that occurs for TMDL pollutant load allocation when a  
25 water quality standard violation occurs. That said, the Panel has the following suggestions for  
26 the modeling of load reduction apportionment for upstream segments:

**Science Advisory Board Panel Draft Report (dated April 8, 2011) - Do not Cite or Quote**

This draft is a work in progress, does not reflect consensus advice or recommendations, has not been approved by the chartered SAB, and does not represent EPA policy.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43

1. The watershed segment approach is valid, but care should be taken in selecting segments to take into account available data and other watershed characteristics such as predominant land-use.

Given a need to complete watershed modeling for the purpose of determining DPVs, the division of the watershed into segments for the purpose of predicting loadings at the “pour point” into the estuary or marine receiving waters should not be limited to simple hydrologic division of the watershed. This may conflict with the premise of using a 12-digit HUC, but the segmentation process needs to take into account predominant land uses for a segment, and those land uses that may be significantly different. For example, urban areas, with high impervious surface cover and altered stream channels, are likely to behave in a way that is distinctly different than less developed areas. Therefore, a simple model delineation of subwatersheds may not be suitable and some expert analysis and adjustment of the segments would be more appropriate.

2. The impacts of urban environments should be considered.

Urbanized areas have a distinct influence on normal stream processes given their large areas of impervious cover. In addition to changes in stream habitat, runoff from impervious surfaces as well as municipal and industrial discharges may contribute to stream nutrient loads. For this reason, the Panel recommends that large urbanized areas be given special consideration in any modeling approach that might be used to generate DPVs.

3. Given that a complete uncertainty analysis cannot be accomplished, it is essential that, in all text in the revised report where uncertainty is mentioned, readers are clearly told what is included (excluded) in any uncertainty analysis undertaken or contemplated.
4. EPA should provide justification for the choice of the LSPC model and explain why it is the most applicable model for this case.

The LSPC model is an updated version of the older HSPF program. While the model can be integrated with a Geographic Information System (GIS), it is not a GIS-based approach. Numerous models exist for watershed management, physical flow, and water quality modeling that may better utilize the strengths of current GIS platforms. For example, the SWAT (Soil and Water Assessment Tool) model is commonly used in the development of TMDLs and is run within a GIS. Given the complexity of watershed modeling at the proposed scale and the complex nature of the problem being addressed, it may be prudent to build watershed models that can take advantage of a wider array of GIS-based tools and data for the current project and in future applications during implementation. This would not be a significant increase in work load, as the development of the GIS and SWAT model would replace the development of the LSPC model.

**Science Advisory Board Panel Draft Report (dated April 8, 2011) - Do not Cite or Quote**

This draft is a work in progress, does not reflect consensus advice or recommendations, has not been approved by the chartered SAB, and does not represent EPA policy.

- 1 5. The time frame of modeling is important and should be linked to the response of the  
2 endpoints in the receiving waters.  
3

4 In the EPA presentation on development of DPVs, it was indicated that adjustments for  
5 seasonal effects and flow levels are being considered. This is a very important consideration and  
6 the EPA is encouraged to analyze available data in the context of seasonal changes in the  
7 watershed and for the differences between baseflow and storm event conditions. Seasonal  
8 changes in the watershed may result from both natural processes (e.g., biotic activity) and from  
9 anthropogenic factors (e.g., agricultural practices). The differences in loadings seen during  
10 baseflow and storm events may be dramatic, with the majority of loading of TN and TP coming  
11 during a few large storm events. This is particularly true for N and P species associated with  
12 suspended sediment. Using an annual average value may grossly underpredict the impact of  
13 large storm events. Therefore, EPA should evaluate the sensitivity of the selected biological  
14 endpoints to the potential influences of shorter-term (e.g., days to weeks) events that may result  
15 in high levels of TN and TP loading to determine if annual or seasonal averages are sufficient to  
16 protect estuarine biota.  
17

- 18 6. In-stream/watershed P transformations should be considered in greater depth for streams,  
19 lakes and canals.  
20

21 Species/fractions of N and P are often a part of TMDL modeling. If DPVs are to be  
22 developed in Florida, expressed as loads, and serve in a TMDL-like role, then DPVs might be  
23 expressed as nutrient fractions (for a biotic estuarine water quality criterion). In the discussion of  
24 nutrients, EPA correctly identifies the role of N species/fractions, but does not consider P  
25 species/fractions.  
26

27 The dynamics of P in watersheds, lakes and canals is important to any effort to produce  
28 DPVs or similar water quality criteria. Foremost is the need to recognize the mobility, reactivity  
29 and bioavailability of the different P species: soluble reactive phosphorus (SRP); dissolved  
30 phosphorus (DP), which is the sum of SRP and total hydrolysable P (THP); and total phosphorus  
31 (TP), which is the sum of DP and particulate phosphorus (PP). These phases exist in natural  
32 waters in varying degrees that are dependent on processes within the waterbody and on external  
33 inputs. Furthermore, transitions among forms occur during transport within watersheds and  
34 within sediments in streams, lakes and canals. Although the modeling effort will evaluate TP for  
35 DPVs, the transport and fate of P should not be oversimplified. A brief overview of some of the  
36 extensive literature on P cycling and transformations within watersheds is provided in Appendix  
37 B.  
38

- 39 7. How are nutrients, especially P, from natural or geologic sources separated from  
40 anthropogenic sources?  
41

42 Further complicating apportionment and determination of DPVs is the issue of  
43 background values for nutrients, especially P. Given that some areas of Florida have bedrock  
44 geology with high P concentrations, understanding background is critical. In watersheds where  
45 high P loadings are the result of natural factors, DPVs may not be applicable.  
46

**Science Advisory Board Panel Draft Report (dated April 8, 2011) - Do not Cite or Quote**

This draft is a work in progress, does not reflect consensus advice or recommendations, has not been approved by the chartered SAB, and does not represent EPA policy.

- 1 8. The continuum of fresh to saline waters in going from watersheds to the receiving estuarine  
2 or coastal marine waters must be considered in the process of determining DPVs.  
3

4 In many instances, fresh water systems are P-limited with respect to nutrient balance and  
5 the potential for the development of eutrophic conditions. The opposite is often the case for  
6 estuarine or marine waters, where N and P can be co-limiting or N is the limiting nutrient. This  
7 raises the potential where the application of upstream water quality standards that may be  
8 focused at reducing P inputs could be protective of the watershed, but create a situation in the  
9 downstream brackish or saline receiving waters that creates a nutrient imbalance (Elmgren and  
10 Larsson 2001; Paerl et al. 2004). This issue should be recognized during development of DPVs  
11 and implementation of recent inland water criteria.  
12

13 **3.6.2 DPVs for South Florida Estuarine and Coastal Waters**

14 *Charge Question 6(b). Are the methods that EPA is considering for deriving downstream*  
15 *protection values (DPVs) for marine waters in South Florida as described in Section 6.5*  
16 *appropriate to ensure attainment and maintenance of downstream water quality standards,*  
17 *given available data? Please describe additional approaches and their advantages and*  
18 *disadvantages that EPA should consider when developing numeric criteria to protect*  
19 *downstream marine waters in South Florida, given available data?*  
20

21 Unlike for estuaries in other parts of the state, the EPA document is not proposing an  
22 upstream apportionment of load reduction by stream segment because of the highly managed  
23 hydrology in South Florida. Instead, the document proposes setting a protective load at the  
24 terminal reach of each tributary, i.e., at the point where the tributary empties into estuarine or  
25 coastal waters. The EPA document discusses several schemes for allocating acceptable  
26 estimated nutrient loads among tributaries, including allocation based on flow-weighted  
27 concentration, flow-only, or total load for each tributary. The Panel's thoughts on various  
28 possible allocation schemes are presented in the previous section.  
29

30 As discussed in section 3.4.1, the Panel recommends that nutrient criteria for canals be  
31 focused on downstream, rather than instream, protection from nutrient impacts. The Panel has  
32 the following additional comments:  
33

- 34 1. Provide more information on how canals will be evaluated.  
35

36 A number of primary canals empty directly into coastal waters, so it will be important to  
37 incorporate all available data on N and P species for the terminal reach of these canals and to  
38 provide a more detailed approach on how DPV criteria will be developed.  
39

- 40 2. The time frame of modeling is important and should be linked to the response of the  
41 endpoints in the receiving waters.  
42

43 Given the wide variation in flow conditions for canals, the concentrations of nutrients in  
44 canal waters are likely highly variable. Hence average nutrient concentrations in canal waters  
45 when released to estuarine and coastal marine waters may not adequately represent the  
46 concentrations needed to protect receiving waters. If additional information is not available for

**Science Advisory Board Panel Draft Report (dated April 8, 2011) - Do not Cite or Quote**

This draft is a work in progress, does not reflect consensus advice or recommendations, has not been approved by the chartered SAB, and does not represent EPA policy.

1 nutrient concentrations in the canals, discharge of canal waters to the receiving waterbodies  
2 needs to take into consideration loading rates on a daily basis that will ensure the receiving  
3 waterbodies meet their water quality standards.

4  
5 3. In-stream/watershed P transformations should be considered in more depth for streams, lakes  
6 and canals.

7  
8 Although less is known about P transformations within the canals of South Florida than  
9 for streams, the physical and chemical processes that control P transport within a watershed  
10 should be the same for canals. Additional consideration, however, must be given to the special  
11 situations that result as a function of the wide-ranging flow situations for the canal system.  
12 Furthermore, it is important to understand the temporal parameters and their range of variability.  
13 These factors will determine, in part, the mechanisms that are most important under different sets  
14 of flow conditions.

15  
16 4. The continuum of fresh to saline waters must be considered in the process of determining  
17 DPVs.

18  
19 For canal waters discharging to estuarine and coastal marine waters, the issue of the  
20 continuum of fresh to saline water is the same as discussed earlier in our response to charge  
21 question 6(a), above.  
22

**Science Advisory Board Panel Draft Report (dated April 8, 2011) - Do not Cite or Quote**

This draft is a work in progress, does not reflect consensus advice or recommendations, has not been approved by the chartered SAB, and does not represent EPA policy.

**REFERENCES**

1  
2  
3  
4 Baden S., Boström, C., Tobiasson S, Arponen, H and Moksnes P-O. 2010. Relative importance  
5 of trophic interactions and nutrient enrichment in seagrass ecosystems: A broad-scale  
6 experimental assessment. *Limnology and Oceanography* **55**:1435-1448.  
7  
8 Banse, K. 1977. Determining the carbon-to-chlorophyll ratio of natural phytoplankton. *Marine*  
9 *Biology* **41**:199-212.  
10  
11 Bostrom, B., A.K. Pettersson, and I. Ahlgren. 1989. Seasonal dynamics of a cyanobacterial  
12 dominated microbial community in surface sediments of a shallow eutrophic lake. *Aquat.*  
13 *Sci.* **51**:153-178.  
14  
15 Boyer, J. and H. O. Briceño. 2010. Written comments provided to the SAB Nutrient Criteria  
16 Review Panel, December 2, 2010. Available at  
17 [http://yosemite.epa.gov/sab/sabproduct.nsf/C485AFF485895924852577F100569C8F/\\$File/Boyer+Letter+to+SAB+on+EPA+NNC+Approach+2010-12-02.pdf](http://yosemite.epa.gov/sab/sabproduct.nsf/C485AFF485895924852577F100569C8F/$File/Boyer+Letter+to+SAB+on+EPA+NNC+Approach+2010-12-02.pdf)  
18  
19  
20 Briceño, H., J. Boyer and P. Harlem. 2010. Proposed Methodology for the Assessment of  
21 Protective Numeric Nutrient Criteria for South Florida Estuaries and Coastal Waters.  
22 Comments submitted to the SAB Staff Office, December 6, 2010. Available at  
23 [http://yosemite.epa.gov/sab/sabproduct.nsf/3759224AA1DA4CAD852577F3005B533F/\\$File/Briceno+comments.pdf](http://yosemite.epa.gov/sab/sabproduct.nsf/3759224AA1DA4CAD852577F3005B533F/$File/Briceno+comments.pdf)  
24  
25  
26 Burkepile, D. E. and M.E. Hay. 2006. Herbivore vs. nutrient control of marine primary  
27 producers: context-dependent effects. *Ecology* **87**:3128-3139.  
28  
29 Burkholder, J.M., D.A. Tomasko and B.W. Touchette. 2007. Seagrasses and eutrophications. *J.*  
30 *Exper. Mar. Biol Ecol.* **350**:46-72.  
31  
32 Carlson, P.R. and K. Madley. 2007. Statewide Summary for Florida. In: Handley, L., D. Altsman  
33 and R. DeMay.(eds.). Seagrass status and trends in the northern Gulf of Mexico. pp. 99-114.  
34 USGS SIR 06-5287. <http://pubs.usgs.gov/sir/2006/5287>  
35  
36 Carrick, H.J. and A.D. Steinman. 2001. Variation in periphyton biomass and species  
37 composition in Lake Okeechobee, Florida (USA). *Archiv fur Hydrobiologie* **152**:411-  
38 438.  
39  
40 Cloern, J.E. 1982. Does the benthos control phytoplankton biomass in south San Francisco Bay?  
41 *Marine Ecology Progress Series* **9**:191-202.  
42  
43 Cohen, R.H., P.V. Dresler, E.J. P. Phillips, and R.L. Cory. 1984. The effect of the Asiatic clam,  
44 *Corbicula fluminea*, on phytoplankton of the Potomac River, Maryland. *Limnol. Oceanogr.*  
45 **29**:170-180.  
46

**Science Advisory Board Panel Draft Report (dated April 8, 2011) - Do not Cite or Quote**

This draft is a work in progress, does not reflect consensus advice or recommendations, has not been approved by the chartered SAB, and does not represent EPA policy.

- 1 Conley, D.J., H. W. Paerl, R.W. Howarth, D.F. Boesch, S.P. Seitzinger, K.E. Havens, C.  
2 Lancelot, and G.E. Likens. 2009. Controlling Eutrophication: Nitrogen and Phosphorus.  
3 *Science* **323**:1014-1015.  
4
- 5 DeBusk, T.A. 2010. Comments on Numeric Nutrient Criteria for Florida's Southern Inland  
6 Flowing Waters. Presentation to the SAB Nutrient Criteria Review Panel, December 13,  
7 2010.  
8
- 9 Dixon, K. 1999, Establishing light requirements for the seagrass *Thalassia testudinum*: an  
10 example for Tampa Bay, Florida. In: Bortone, SA (ed) *Seagrasses: Monitoring Ecology*  
11 *Physiology and Management*, pp. 81-98. CRC Press, Boca Raton, FL.  
12
- 13 Doering, P.H., R.H. Chamberlain, K.M. Donohue, and A.D. Steinman. 1999. Effect of salinity  
14 on the growth of *Vallisneria americana* Michx. from the Caloosahatchee Estuary,  
15 Florida. *Florida Scientist* **62**:89-105.  
16
- 17 Elmgren, R., and U. Larsson. 2001. Nitrogen and the Baltic Sea: Managing nitrogen in relation  
18 to phosphorus. *The Scientific Word* **1**: 371-377.  
19
- 20 Evans, D.J. and P.J. Johnes. 2004. Physico-chemical controls on phosphorus cycling in  
21 twolowland streams. Part 1 — the water column. *Sci. Total Environ.* **329**:145–163.  
22
- 23 Evans, D.J., P.J. Johnes, and D.S. Lawrence. 2004. Physico-chemical controls on phosphorus  
24 cycling in two lowland streams. Part 2—the sediment phase. *Sci. Total Environ.*  
25 **329**:165–182.  
26
- 27 Fore, L.S. 2004. *Development and Testing of Biomonitoring Tools for Macroinvertebrates in*  
28 *Florida Streams*. Prepared for Florida Department of Environmental Protection by  
29 Statistical Design, Seattle, WA. 74p. Available at  
30 [http://publicfiles.dep.state.fl.us/dear/labs/sas/sopdoc/sci\\_old.pdf](http://publicfiles.dep.state.fl.us/dear/labs/sas/sopdoc/sci_old.pdf) (accessed 01/15/11).  
31
- 32 Golterman, H.L. 1995. The role of the ironhydroxide–phosphate–sulphide system in the  
33 phosphate exchange between sediments and overlying water. *Hydrobiol.* **297**:43–54.  
34
- 35 Greening, H. 2010. Presentation on EPA's Proposed Approaches for Florida ENCs: Observations  
36 from Tampa Bay. Presentation to the SAB Nutrient Criteria Review Panel, December 13,  
37 2010.
- 38 Handley, L. D. Altsman and R. DeMay. 2007. Abstract. In: Handley, L., D. Altsman and R.  
39 DeMay.(eds.). *Seagrass status and trends in the northern Gulf of Mexico*. pp.1-3. USGS  
40 SIR 06-5287. <http://pubs.usgs.gov/sir/2006/5287>
- 41 Hauxwell, J., J. Cebrián, C. Furlong and I. Valiela, 2001. Macroalgal canopies contribute to  
42 eelgrass (*Zostera marina*) decline in temperate estuarine ecosystems. *Ecology* **82**:1007–  
43 1022.

**Science Advisory Board Panel Draft Report (dated April 8, 2011) - Do not Cite or Quote**

This draft is a work in progress, does not reflect consensus advice or recommendations, has not been approved by the chartered SAB, and does not represent EPA policy.

- 1 Hauxwell, J., J. Cebrián and I. Valiela. 2003. Eelgrass *Zostera marina* loss in temperate  
2 estuaries: relationship to land-derived nitrogen loads and effect of light limitation  
3 imposed by algae. *Mar. Ecol. Prog. Ser.* **247**:59–73.
- 4 He, R. and R.H. Weisberg. 2002. Tides on the West Florida shelf. *J. Phys. Oceanography* **33**:  
5 465-477.  
6
- 7 Heck, Jr., K.L. and J.F. Valentine. 2007. The primacy of top-down effects in shallow benthic  
8 ecosystems. *Estuaries and Coasts* **30**:371-381.  
9
- 10 Hilton, J., M. O'Hare, M.J. Bowes, and J.I Jones. 2006. How green is my river? A new paradigm  
11 of eutrophication in rivers. *Science of the Total Environment* **365**:66-83.  
12
- 13 House, W.A., 2003. Geochemical cycling of phosphorus in rivers. *Appl. Geochem.* **18**:739-748.  
14
- 15 Hu, C., Z. Chen, T.D. Clayton, P. Swarzenski, J.C. Brock, and F.E. Muller-Karger. 2005.  
16 Assessment of estuarine water-quality indicators using MODIS medium-resolution  
17 bands: Initial results from Tampa Bay, FL. *Remote Sensing of Environment* **94**:425-427.  
18
- 19 Huang, B.Q. and H.S. Hong. 1999. Alkaline phosphatase activity and utilization of dissolved  
20 organic phosphorus by algae in subtropical coastal waters. *Marine Pollution Bulletin*  
21 **39**:205-211.  
22
- 23 Hughes, A.R., K.J. Bando, L.F. Rodriguez and S.L. Williams. 2004. Relative effects of grazers  
24 and nutrients on seagrasses: a meta-analysis approach. *Mar Ecol Progr Ser* **282**:87-99.  
25
- 26 Jarvie, H.P., Neal, C., Jurgens, M.D., Sutton, E.J., Neal, M., Wickham, H.D., Hill, L.K., Harman,  
27 S.A., Davies, J.J.L., Warwick, A., Barrett, C., Griffiths, J., Binley, A., Swannack, N.,  
28 McIntyre, N., 2006. Within-river nutrient processing in Chalk streams: the Pang and  
29 Lambourn, UK. *J. Hydrol.* **330**:101–125.  
30
- 31 Jensen, H.S. and F.Ø. Andersen. 1992. Importance of temperature, nitrate, and pH for phosphate  
32 release from aerobic sediments of four shallow, eutrophic lakes. *Limnol. Oceanogr.*  
33 **37**:577–589.  
34
- 35 Johnes, P.J. and R.A. Hodgkinson. 1998. Phosphorus loss from agricultural catchments: pathways  
36 and implications for management. *Soil Use Manage* **14**:175–185.  
37
- 38 Koblinsky, C. J. 1981. The  $M_2$  tide on the West Florida Shelf. *Deep-Sea Res.* **28A**:1517–1532.  
39
- 40 Kraemer, G.P., R.H. Chamberlain, P.H. Doering, A.D. Steinman, and M.D. Hanisak 1999.  
41 Physiological Responses of Transplants of the Freshwater Angiosperm *Vallisneria*  
42 *americana* along a Salinity Gradient in the Caloosahatchee Estuary (Southwestern  
43 Florida). *Estuaries* **22**:138-148.

**Science Advisory Board Panel Draft Report (dated April 8, 2011) - Do not Cite or Quote**

This draft is a work in progress, does not reflect consensus advice or recommendations, has not been approved by the chartered SAB, and does not represent EPA policy.

- 1 Latimer, J.S. and S.A. Rego. 2010. Empirical relationships between eelgrass extent and predicted  
2 watershed-derived nitrogen loading for shallow New England estuaries. *Estuarine,*  
3 *Coastal and Shelf Science* **90**:231-240.  
4
- 5 Lee, T.N. and D.A. Mayer. 1977. Low-frequency current variability and spin-off eddies on the  
6 shelf off southeast Florida. *J. Mar. Res.* **35**:193-220.  
7
- 8 Lee, T.N., E. Williams, D. Wilson, E. Johns, and N. Smith. 2002. Transport processes linking  
9 south Florida coastal ecosystems. pp. 309-342, In: *The Everglades, Florida Bay and*  
10 *Coral Reefs of the Florida Keys: An Ecosystem Sourcebook*.  
11
- 12 Lewis, W.M. and J.H. McCutchan. 2010. Ecological responses to nutrients in streams and rivers  
13 of the Colorado mountains and foothills. *Freshwater Biology* **55**:1973-1983.  
14
- 15 McCormick, P.V. and M.B. O'Dell. 1996. Quantifying periphyton responses to phosphorus  
16 enrichment in the Florida Everglades: a synoptic-experimental approach. *Journal of the*  
17 *North American Benthological Society* **15**:450-468.  
18
- 19 McCormick, P.V., P.S. Rawlik, K. Lurding, E.P. Smith, and F.H. Sklar. 1996. Periphyton-water  
20 quality relationships along a nutrient gradient in the northern Everglades. *Journal of the*  
21 *North American Benthological Society* **15**:433-449.  
22
- 23 McCormick, P.V., S. Newman, S. Miao, D.E. Gawlick, D. Marley, K.R. Reddy, and T.D.  
24 Fontaine. 2002. Effects of anthropogenic phosphorus inputs on the Everglades. Pages 83-  
25 126 in: J. Porter and K. Porter (editors). *The Everglades, Florida Bay, and Coral Reefs of*  
26 *the Florida Keys. An Ecosystem Handbook*. CRC Press, Boca Raton, FL.  
27
- 28 Melack, J.M. 1995. Transport and transformations of P, fluvial and lacustrine ecosystems. In  
29 *Phosphorus in the global environment*, ed. H. Tniessen, Chapter 15. West  
30 Sussex, England: John Wiley and Sons Ltd.  
31
- 32 Muller-Karger, F. E., C. Hu, S. Andréfouët, and R. Varela. 2005. The Color of the Coastal Ocean  
33 and applications in the solution of research and management problems. *In: Remote*  
34 *Sensing of Coastal Aquatic Environments: Technologies, Techniques and Application*,  
35 R.L. Miller, C.E. Del Castillo and B.A. McKee [Eds.], Springer, 101-127.  
36
- 37 Owens, P.N. and D.E. Walling. 2002. The phosphorus content of fluvial sediment in ruraland  
38 industrialized river basins. *Water Res.* **36**:685–701.  
39
- 40 Paerl, H.W. 2009. Controlling Eutrophication along the freshwater–Marine Continuum: Dual  
41 Nutrient (N and P) Reductions are Essential. *Estuaries and Coasts* **32**:593-601.  
42
- 43 Paerl, H.W. and J. Huisman. 2008. Blooms like it hot. *Science* **320**:57-58.  
44
- 45 Paerl, H.W., L.M. Valdes, M.F. Piehler and M.E. Lebo. 2004. Solving problems resulting from  
46 solutions: The evolution of a dual nutrient management strategy for the eutrophying

**Science Advisory Board Panel Draft Report (dated April 8, 2011) - Do not Cite or Quote**

This draft is a work in progress, does not reflect consensus advice or recommendations, has not been approved by the chartered SAB, and does not represent EPA policy.

- 1 Neuse River Estuary, North Carolina, USA. *Environmental Science & Technology* **38**:  
2 3068-3073.  
3
- 4 Paerl, H.W., L.M. Valdes, A.R. Joyner, B.L. Peierls, C.P. Buzzelli, M. F. Piehler, S.R. Riggs, R.  
5 R. Christian, J.S. Ramus, E.J. Clesceri, L.A. Eby, L.W. Crowder, and R.A. Luettich .  
6 2006a. Ecological response to hurricane events in the Pamlico Sound System, NC and  
7 implications for assessment and management in a regime of increased frequency.  
8 *Estuaries and Coasts* **29**:1033-1045.  
9
- 10 Paerl, H.W., L.M. Valdes, J.E. Adolf, B.M. Peierls and L.W. Harding Jr. 2006b. Anthropogenic  
11 and climatic influences on the eutrophication of large estuarine ecosystems. *Limnology*  
12 *and Oceanography* **51**:448-462.  
13
- 14 Paerl, H.W., L.M. Valdes, A.R. Joyner, and V. Winkelmann. 2007. Phytoplankton Indicators of  
15 Ecological Change in the Nutrient and Climatically-Impacted Neuse River-Pamlico  
16 Sound System, North Carolina. *Ecological Applications* **17**:88-101.  
17
- 18 Palandro, D., C. Hu, S. Andréfouët, and F.E. Müller-Karger. 2004. Synoptic water clarity  
19 assessment in the Florida Keys using diffuse attenuation coefficient estimated from  
20 Landsat imagery. *Hydrobiologia* **530-531(1)**:489-493.  
21
- 22 Pauly, D. 1995. Anecdotes and the shifting baseline syndrome of fisheries TREE 10:430.  
23
- 24 Ryan, R.J., A.I. Packman, and S.S. Kilham. 2007. Relating phosphorus uptake to changes in  
25 transient storage and streambed sediment characteristics in headwater tributaries of Valley  
26 Creek, an urbanizing watershed. *J. Hydrol.* **336**:444-457.  
27
- 28 Rydin, E. 2000. Potentially mobile phosphorus in Lake Erken sediments. *Water Res.* **34**:2037-  
29 2042.  
30
- 31 Rydin, E. And A.K. Brunberg. 1998. Seasonal dynamics of phosphorus in Lake Erken surface  
32 sediments. *Arch. Hydrobiol. Spec. Issues Advanc. Limnol.* **51**:157-167.  
33
- 34 South Florida Water Management District (SFWMD). 2010. Canals in South Florida: A  
35 Technical Support Document (April 28, 2010). SFWMD, West Palm Beach, FL.  
36
- 37 Snyder, B., M. Barbour, and E. Leppo. 1998. *Development of a watershed-based approach for*  
38 *biomonitoring of fresh surface waters in coastal Florida canal system*. Prepared by  
39 TetraTech, Inc., Owings Mills, MD under contract with Metro-Dade Environmental  
40 Resources Management, 201 p.  
41
- 42 Steinman, A., K. Havens, L. Hornung. 2002a. The Managed Recession of Lake Okeechobee,  
43 Florida: Integrating Science and Natural Resource Management. *Conservation Ecology*  
44 **6**:243-256.  
45

**Science Advisory Board Panel Draft Report (dated April 8, 2011) - Do not Cite or Quote**

This draft is a work in progress, does not reflect consensus advice or recommendations, has not been approved by the chartered SAB, and does not represent EPA policy.

- 1 Steinman, A.D., K.E. Havens, H.J. Carrick, and R. VanZee. 2002b. The past, present, and future  
2 hydrology and ecology of Lake Okeechobee and its watersheds. Pages 19-37 in: J. Porter  
3 and K. Porter (editors). *The Everglades, Florida Bay, and Coral Reefs of the Florida*  
4 *Keys. An Ecosystem Handbook*. CRC Press, Boca Raton, FL.  
5
- 6 Steinman, A.D., M. Luttenton, and K.E. Havens. 2004. Sustainability of surface and subsurface  
7 water resources: case studies from Florida and Michigan. *Water Resources Update* **127**:  
8 100-107.  
9
- 10 Stumpf, R.P., M.E. Culver, P.A. Tester, M. Tomlinson, G.J. Kirkpatrick, B.A. Pederson, E.  
11 Truby, V. Ransibrahmanakul, and M. Soracco. 2003. Monitoring *Karenia brevis* blooms  
12 in the Gulf of Mexico using satellite ocean color imagery and other data. *Harmful Algae*  
13 **2**:147-160.  
14
- 15 Stumpf, R.P., M.C. Tomlinson, J.A. Calkins, B. Kirkpatrick, K. Fisher, K. Nierenberg, R.  
16 Currier, and T.T. Wynne. 2009. Skill assessment for an operational algal bloom forecast  
17 system. *J. Mar. Syst.* **76**:151-161.  
18
- 19 Thompson, J.K., J.R. Koseff, S.G. Monismith, and L.V. Lucas. 2008. Shallow water processes  
20 govern system-wide phytoplankton bloom dynamics: A field study. *Journal of Marine*  
21 *Systems* **74**:153-166.  
22
- 23 Tomlinson, M.C., R.P. Stumpf, V. Ransibrahmanakul, E.W. Truby, G.J. Kirkpatrick, B.A.  
24 Pederson, G.A. Vargo, and C.A. Heil. 2004. Evaluation of the use of SeaWiFS imagery  
25 for detecting *Karenia brevis* harmful algal blooms in the eastern Gulf of Mexico. *Remote*  
26 *Sensing of the Environment* **91**:293-303.  
27
- 28 U.S. EPA (Environmental Protection Agency). 2010a. *Methods and Approaches for Deriving*  
29 *Numeric Criteria for Nitrogen/Phosphorus Pollution in Florida's Estuaries, Coastal*  
30 *Waters, and Southern Inland Flowing Waters* (November 17, 2010 Draft).  
31
- 32 U.S. EPA (Environmental Protection Agency). 2010b. *Using Stressor-response Relationships to*  
33 *Derive Numeric Nutrient Criteria*. EPA Office of Water, November 2010. EPA-820-S-  
34 10-001.  
35
- 36 Valdes-Weaver, L.M., M.F. Piehler, J.L. Pinckney, K.E. Howe, K. Rosignol, and H.W. Paerl.  
37 2006. Long-term temporal and spatial trends in phytoplankton biomass and class-level  
38 taxonomic composition in the hydrologically variable Neuse-Pamlico estuarine  
39 continuum, NC, USA. *Limnology and Oceanography* **51**:1410-1420.  
40
- 41 Verdonchot, P.F.M. 1987. Aquatic oligochaetes in ditches. *Hydrobiologia* **155**:283-292.  
42
- 43 Walker, N.D. and N.N. Rabalais. 2006. Relationships among satellite chlorophyll a, river inputs,  
44 and hypoxia on the Louisiana continental shelf, Gulf of Mexico. *Estuaries and Coasts*  
45 **29**:1081-1093.  
46

**Science Advisory Board Panel Draft Report (dated April 8, 2011) - Do not Cite or Quote**

This draft is a work in progress, does not reflect consensus advice or recommendations, has not been approved by the chartered SAB, and does not represent EPA policy.

- 1 Webster, P.J., G.J. Holland, J.A. Curry, and H.R. Chang. 2005. Changes in tropical cyclone
- 2 number, duration, and intensity in a warming environment. *Science* **309**:1844-1846.
- 3

**Charge to the  
Science Advisory Board  
Nutrient Criteria Review Panel**

**Methods and Approaches for Deriving Numeric Criteria for Nitrogen/Phosphorus  
Pollution in Florida's Estuaries, Coastal Waters, and Southern Inland Flowing Waters**

**Background**

In 2011, EPA will propose numeric criteria for nitrogen/phosphorus pollution to protect estuaries, coastal areas and South Florida inland flowing waters that have been designated Class I, II and III<sup>1</sup>, as well as downstream protective values (DPVs) to protect estuarine and marine waters. This is the second phase of a rulemaking effort as EPA has already established numeric criteria for Florida's lakes, flowing waters<sup>2</sup> and springs within the State of Florida, including DPVs for lakes.

As part of this current effort to derive chlorophyll-a, total nitrogen (TN) and total phosphorus (TP) criteria to protect estuarine and coastal water bodies, EPA must also develop additional criteria to assure that upstream criteria will meet standards established for downstream estuarine and marine waters in Florida. These DPVs, will supplement the existing inland water TN and TP stream criteria, if the applicable DPV is more stringent.

Overall, these numeric criteria are being developed to translate and implement Florida's existing narrative nutrient criterion, to protect the designated use that Florida has previously set for these waters, at Rule 62-302.530(47)(b), F.A.C. which provides that

*"In no case shall nutrient concentrations of a body of water be altered so as to cause an imbalance in natural populations of aquatic flora or fauna."*

Under the Clean Water Act and EPA's implementing regulations, these numeric criteria must be based on sound scientific rationale and reflect the best available scientific knowledge.

EPA has previously published a series of peer reviewed technical guidance documents<sup>3</sup> to develop numeric criteria to address nitrogen/phosphorus pollution in different water body types. EPA recognizes that available and reliable data sources for use in numeric criteria development vary across estuarine and coastal waters in Florida and flowing waters in South Florida. In addition, scientifically defensible approaches for numeric criteria development have different requirements that must be taken into consideration in the context of the specific application and

---

<sup>1</sup> Class I Potable water supplies; Class II Shellfish propagation or harvesting; Class III Recreation, propagation, and maintenance of a healthy, well-balanced population of fish and wildlife.

<sup>2</sup> In the November 2010 rulemaking, EPA did not establish numeric criteria for inland flowing waters in South Florida. For the purpose of this effort, EPA has distinguished South Florida as those areas south of Lake Okeechobee and the Caloosahatchee River watershed to the west of Lake Okeechobee and the St. Lucie watershed to the east of Lake Okeechobee.

<sup>3</sup> U.S. EPA. 2000. Nutrient Criteria Technical Guidance Manual: Rivers and Streams. EPA-822-B-00-002; U.S. EPA. 2001. Nutrient Technical Guidance Manual: Estuarine and Coastal Marine Waters. EPA-822-B-01-003; U.S. EPA. 2010. Using Stressor-response Relationships to Derive Numeric Nutrient Criteria. EPA-820-S-10-001;

available information. This document describes the scientific approaches EPA is considering the derivation of numeric criteria to address nitrogen/phosphorus pollution in Florida estuarine and coastal waters, and inland flowing waters in South Florida, given the available data currently available.

**Review Document:** The SAB is asked to review the draft document, *Methods and Approaches for Deriving Numeric Criteria for Nitrogen/Phosphorus Pollution in Florida's Estuaries, Coastal Waters, and Southern Inland Flowing Waters*, and respond to the following charge questions.

### **Charge Questions:**

#### **1. General Approach**

- a) EPA has introduced a general conceptual model in Chapter 2, including the selection of assessment endpoint and indicator variables. What is your perspective of the general conceptual model?
- b) EPA has delineated the State of Florida into 4 general categories of waters—Florida estuaries, Florida coastal waters, South Florida inland flowing waters, and South Florida marine waters—for purposes of considering approaches to numeric nutrient criteria development. Are these categories appropriate and scientifically defensible? (Note that the details of segmentation of waters within these categories is addressed in subsequent charge questions.)

#### **Florida Estuaries (Chapter 3)**

EPA is considering three approaches: (1) reference conditions, (2) stressor/response models, and (3) water quality simulation modeling that could be used independently or in combination to develop numeric criteria for estuaries (exclusive of marine waters in South Florida that are covered in Chapter 5). Estuarine waters are defined as “a part of a river or stream or other body of water that has an unimpaired connection with the open sea and where the sea water is measurably diluted with fresh water derived from land drainage”<sup>4</sup>.

#### **2. Estuaries**

- a) Are the data sources identified appropriate for use in deriving numeric criteria in Florida's estuaries (as discussed in Sections 2.4 and 3.2)? Is the SAB aware of additional available, reliable data that EPA should consider in delineating estuaries or deriving criteria for estuarine waters? Please identify the additional data sources.
- b) Are the assessment endpoints identified in Sections 2.3 and 3.2 (healthy seagrass communities; balanced phytoplankton biomass and production; and balanced faunal

---

<sup>4</sup> U.S. EPA. 2000 Estuaries and Clean Waters Act of 2000, <http://water.epa.gov/type/oceb/nep/320.cfm> (accessed 11/1/2010)

communities) appropriate to translate Florida's narrative nutrient criterion (as cited above) into numeric criteria for Florida's estuaries, given currently available data? Does the SAB suggest modification or addition to these assessment endpoints? A literature review of endpoints considered can be found in Appendix B.

- c) EPA describes potential approaches in Section 3.3 (reference conditions, stressor response relationships, and water quality simulation models) for deriving numeric criteria in Florida's estuaries. Compare and contrast the ability of each approach to ensure the attainment and maintenance of natural populations of aquatic flora and fauna for different types of estuaries, given currently available data?

## Florida Coastal Waters (Chapter 4)

EPA is considering a reference-based approach to derive numeric criteria for most of Florida's coastal waters. Coastal waters are defined as marine waters up to three nautical miles from shore<sup>5</sup>. Specifically, EPA is considering the use of a remote sensing method to develop numeric Chl<sub>RS-a</sub> criteria for the Northwest Gulf Coast, West Gulf Coast, and Atlantic Coastal areas. Due to interference from colored dissolved organic matter and bottom reflectance on satellite measurements, EPA is not considering the derivation of Chl<sub>RS-a</sub> criteria using remote sensing data in coastal waters from Apalachicola Bay to Suwannee River (Big Bend) and South Florida.

### 3. Coastal Waters

- a) Are the data sources identified in Sections 2.4, 4.1.1 and 4.2 appropriate for use in deriving numeric criteria in Florida's coastal waters? Is the SAB aware of additional available, reliable data that EPA should consider in delineating coastal waters or deriving criteria for coastal waters? Please identify the additional data sources.
- b) Is the assessment endpoint identified in Section 4.2 (chlorophyll-a to measure balanced phytoplankton biomass and production) appropriate to translate Florida's narrative nutrient criteria (described above) into numeric criteria for Florida's coastal waters, given currently available data? Does the SAB suggest modification or addition to this assessment endpoint?
- c) Does the approach EPA describes in Section 4.2 appropriately apply remote sensing data to ensure attainment and maintenance of balanced natural populations of aquatic flora and fauna in Florida's coastal waters? If not, please provide an alternate methodology utilizing available reliable data and tools, and describe the corresponding advantages and disadvantages.

---

<sup>5</sup> Based on the Clean Water Act definition of "Waters of the United States"

## **South Florida Inland Flowing Waters (Chapter 5)**

EPA is considering a reference-based approach to derive numeric criteria for South Florida inland flowing waters using least-disturbed sites that support balanced natural populations of aquatic flora and fauna. Alternative methods of criteria derivation for inland flowing waters include stressor-response relationships between chlorophyll-a and TN and TP, and a distributional approach using all sites. South Florida inland flowing waters are defined for this effort as free-flowing, predominantly fresh surface water in a defined channel and include, streams, rivers, creeks, branches, canals, freshwater sloughs, and other similar water bodies located in the South Florida nutrient watershed region<sup>6</sup>.

### **4. South Florida Inland Flowing Waters**

- a) Are the data sources identified in Section 2.4 and 5.4 appropriate for use in deriving numeric criteria in South Florida's inland flowing waters (as discussed in Chapters 2 and 5)? Is the SAB aware of additional available, reliable data that EPA should consider in delineating or deriving criteria for South Florida's inland flowing waters? Please identify the additional data sources.
- b) Are the assessment endpoints identified in Section 5.4 (balanced faunal communities, i.e., aquatic macroinvertebrates, and balanced phytoplankton biomass and production) appropriate to translate Florida's narrative nutrient criteria (described above) into numeric criteria for South Florida's inland flowing waters, given currently available data? Does the SAB suggest modification or addition to these assessment endpoints?
- c) EPA describes two approaches in Section 5.4 (reference conditions and stressor-response relationships) for deriving numeric criteria in South Florida inland flowing waters. Compare and contrast the ability of each approach to ensure attainment and maintenance of balanced natural populations of aquatic flora and fauna in different types of flowing water or geographical areas, given currently available data?

---

<sup>6</sup> The South Florida nutrient watershed region is the area *south* of Lake Okeechobee and the Caloosahatchee River watershed to the west of Lake Okeechobee and the St. Lucie watershed to the east of Lake Okeechobee. EPA is not deriving criteria that would apply to waters located on the Seminole Indian Reservation or the Miccosukee Indian Reservation, waters located in stormwater treatment areas (STAs), wetlands, or marshes; or Class IV canals. EPA is also not establishing new TP criteria for the Everglades Protection Area (EvPA) in deference to the Everglades Forever Act (EFA).

### **South Florida Marine Waters (Chapter 5)**

EPA is considering a reference-based approach to derive numeric criteria in South Florida marine waters using least-disturbed sites that support balanced natural populations of aquatic flora and fauna. South Florida marine waters include estuarine and coastal waters extending three nautical miles offshore. Estuarine and coastal waters in South Florida are considered together because the watershed based approach for delineating water bodies is less suited in South Florida due to its highly managed inland flows and open-water dominated systems (i.e., Florida Bay and Keys)

#### **5. South Florida Marine Waters**

- a) Are the data sources identified in Section 2.4 and 5.5 appropriate for use in deriving numeric criteria in South Florida's marine waters (as discussed in Chapters 2 and 5)? Is the SAB aware of additional available, reliable data that EPA should consider in delineating or deriving criteria for South Florida's marine waters? Please identify the additional data sources.
  
- b) EPA describes two methods in Section 5.6 for using a reference condition approach for deriving numeric criteria in South Florida marine waters (least-disturbed sites or bionomial test). Compare and contrast the ability of each approach to ensure attainment and maintenance of balanced natural populations of aquatic flora and fauna in South Florida marine waters, given currently available data?

## **Downstream Protection Values for Florida Estuaries and South Florida Marine Waters (Chapter 6)**

The approach that EPA is considering for developing stream DPV criteria is to adjust upstream limits on TN and TP loading rates that are needed to support balanced natural populations of aquatic flora and fauna in downstream estuarine waters. The loading limits will be determined as part of the criteria development effort for estuarine and marine waters (as described in Chapters 3 and 5) and are scaled based on the average streamflow entering the estuary to determine criteria for TN and TP concentrations in streams as they discharge into estuaries or marine waters.

DPVs can be determined for upstream reaches within watersheds by accounting for expected loss or permanent retention of TN and TP within the stream network. Because of the complexities associated with the managed flows in South Florida inland flowing waters that are covered in Chapter 5, the fraction of TN or TP from the upstream tributary reach that eventually flows into the marine waters in South Florida cannot be estimated or predicted. Therefore, EPA is considering expressing DPVs at the terminal reach of the tributary into a South Florida estuary as protective concentrations or, alternatively, protective loads.

### **6. Downstream Protection Values for Florida Estuaries and South Florida Marine Waters**

- a. Are the methods EPA is considering for deriving downstream protection values (DPVs) for estuaries (excluding marine waters in South Florida) as described in Section 6.1-6.4 appropriate to ensure attainment and maintenance of downstream water quality standards, given available data? Please describe additional approaches and their advantages and disadvantages that EPA should consider when developing numeric criteria to protect these downstream estuarine waters (excluding marine waters in South Florida), given available data?
- b. Are the methods that EPA is considering for deriving downstream protection values (DPVs) for marine waters in South Florida as described in Section 6.5 appropriate to ensure attainment and maintenance of downstream water quality standards, given available data? Please describe additional approaches and their advantages and disadvantages that EPA should consider when developing numeric criteria to protect downstream marine waters in South Florida, given available data?

## APPENDIX B: Phosphorus Transport and Fate in Freshwater Systems

While streams are often viewed as simply a transfer mechanism for P, recent work has investigated processes that occur during transport. There are mechanisms that transform P within different physico-chemical fractions within the stream channel (Melack, 1995; Evans and Johnes, 2004; Evans et al., 2004), and the speciation of soluble P phases and fractionation of P are critical for any evaluation of transport or retention within a watershed. Various processes transform P including sorption, co-precipitation, and redox reactions (e.g., House 2003), and SRP interacts with stream sediments. Stream sediments act as both sinks and sources for SRP within the stream depending on the SRP concentration in the stream water and may change both temporally and spatially within a watershed (e.g., Jarvie et al., 2006; Ryan et al., 2007). This would suggest that EPA should evaluate existing data sets with regard to SRP and TP concentrations.

In comparing rural versus urbanized watersheds, Owens and Walling (2002) found that PP increased in stream sediments receiving point source discharge high in SRP, and that PP (inorganic and organic) may be the most significant mechanism for P transport. Up to 20% of the PP in stream sediment is likely to be easily bioavailable as inorganic P phases dominate. These mechanisms may also be active in lake or canal sediments. Given the short-term bioavailability of some fraction of the PP, it is important to evaluate TP in the context of SRP, DP and PP with some evaluation of the immediacy of the impact of each fraction.

Phosphorus retention within watersheds is typically dominated by calcite co-precipitation within bed sediment and physical trapping of sediment by reduction of flow velocity. Lake sediments may act as both sinks and sources for P cycling, with a large fraction of the inorganic P in surface sediments in equilibrium with the water column (Golterman 1995). The cycling of P is most prevalent in stratified lakes with anoxic hypolimnion, but significant cycling of P also occurs from oxic sediments (Bostrom et al., 1989; Jensen and Andersen, 1992; Rydin and Brunberg, 1998) found in nearshore environments, stream sediments and likely in canals.

P mobilization occurs under both oxic and anoxic conditions, and exchangeable and Fe-bound P are generally mobile (Rydin 2000). Organic-associated P is about 60% mobile, with greater mobility in anoxic sediments. P associated with Al and Ca is immobile and may be considered permanently bound. P release from aerobic sediments may deplete the Fe-bound P despite Fe remaining in the solid phase (Jensen and Andersen 1992). The release process involves a complex relationship between nitrate concentrations and microbial activity resulting in seasonal effect of increasing sediment P retention during winter with subsequent release during late summer and autumn. Biota also play a role in P cycling in lake sediments (e.g., bioturbation, rooted macrophytes that alter the sediment biogeochemistry). The likely lack of available data on the fractionation of P between the various physico-chemical phases will limit a detailed evaluation; however, it is important that modelling of P transport include some recognition of the biogeochemical processes involved in P cycling.



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, D.C. 20460

FEB 18 2011

**MEMORANDUM**

OFFICE OF  
WATER

**SUBJECT:** Science Advisory Board Nutrient Criteria Review Panel Request for Information

**FROM:** Elizabeth Behl, Director  
Health and Ecological Criteria Division  
Office of Science and Technology

**TO:** Stephanie Sanzone, Designated Federal Official  
Science Advisory Board

This memorandum is in response to the request for information from the U.S. Environmental Protection Agency Science Advisory Board (SAB) on the proposed methods and approaches to derive numeric criteria protective of downstream estuaries in Florida. In the SAB teleconference call on February 7, 2011, we indicated that there were several approaches that could be used to allocate nutrient loads among stream reaches within an estuarine watershed for the purpose of computing downstream protective values (DPVs). We provided one of those approaches to the Panel as an illustrative example at the SAB meeting in December (Option A below). EPA also is considering several other options, described below, including two (Options C and D) which take into consideration initial feedback EPA received from the SAB in December 2010.

**Options for Distributing Loads**

**Option A: Distribute Load in Proportion to Flow.** This option would distribute the watershed load among stream reaches according to the fraction of the total watershed discharge contributed by that reach, thereby addressing the fact that higher nutrient loading is often associated with higher freshwater flow. In developing this option, EPA also considered in-stream processing of nutrients, computing the aggregate loss and/or retention of nutrients within the stream network. Criteria derived using this approach could be higher for streams for which a significant quantity of transported nutrients are lost or retained before reaching estuaries. EPA described this approach in the Methods and Approaches document submitted to the EPA SAB for review.

**Option B: Distribute Load in Proportion to Area.** This option would distribute the watershed load among stream reaches according to the fraction of the total watershed area that is drained via that reach. DPV concentrations could be computed by dividing the loading limit for each reach by the average freshwater discharge from the respective reach. EPA does not intend to consider this approach further, recognizing that it does not take into account differences in freshwater yield (i.e., runoff per area), an important factor affecting nutrient transport from watersheds. Freshwater yield varies significantly among watersheds due to both natural and

anthropogenic causes. For example, a watershed with high relief (i.e., slope) or significant impervious surfaces may have higher freshwater yield than a low-relief, forested watershed. DPV criteria derived using this approach could be much lower for high-runoff watersheds, and in some cases could be unreasonably low.

**Option C: Distribute Load In Proportion to Natural or “Background” Loading.** This option distributes the load among terminal reaches in a watershed in proportion to an estimate of the loading that would occur from each watershed in the absence of anthropogenic influence, which we refer to as “background loading.” These estimates could be derived using the Loading Simulation Program in C++, or LSPC. This option would enable EPA to consider a diverse array of environmental information (such as slope, natural land cover type, soil types, local rainfall) in deriving numeric criteria. As an example, if the watershed TN or TP loading rate for an estuary is, in aggregate, 20% more than the estimate of background loading, the watershed load distributed to each terminal stream reach would be 20% more than the respective background load. DPV criteria could be computed by dividing the distributed loading by the average freshwater discharge. Since the estimates of these loads consider hydrological and other landscape factors, the DPV estimates will also reflect these factors. DPV criteria for upstream reaches could be determined recursively using the same process. DPV criteria derived using this approach limit loading to estuaries to the watershed load and DPVs criteria reflect a range of natural factors that impact nutrient concentrations in watersheds.

**Option D: Distribute Load in Proportion to Existing Loading.** This option distributes the watershed load among the terminal reaches in a watershed based on both the average background loading (as described above) and average existing loading (e.g., 1997-2009), recognizing that the difference between the estimate of existing loading and background loading is an estimate of anthropogenic loading. For example, if the aggregate watershed loading to the estuary is 30% less than the current loading rate, DPV criteria would be computed based on 100% of the background loading plus a fraction of anthropogenic loading, such that the total loading is equal to the watershed load. DPV criteria would be computed by dividing the divided load by the average freshwater flow. DPVs for unmodified watersheds with low nutrient yields would be relatively low, whereas DPVs for sub-watersheds with developed or agricultural lands would be expected to be higher, but lower than existing concentrations. DPV criteria derived using this approach limits loading to estuaries to the watershed load and DPVs reflect both natural and anthropogenic factors that impact nutrient concentrations in watersheds.