

TCE Meeting Presentation: Supporting Material for C. Norman
Paul Dugard
to:
Marc Rigas
05/03/2010 09:01 PM
Please respond to Paul Dugard
Show Details

Dear Dr. Rigas;

Please find attached two tables that Mr Norman wishes to have made available to the panel and which will be referred to in his presentation on May 10.

Thank you.

Paul Dugard

**Contaminated Water Supplies at Camp Lejeune,
Assessing Potential Health Effects
National Research Council of the National Academy of Sciences (2009)**

BOX 1 Five Categories Used by IOM to Classify Associations

Sufficient Evidence of a Causal Relationship

Evidence from available studies is sufficient to conclude that a causal relationship exists between exposure to a specific agent and a specific health outcome in humans, and the evidence is supported by experimental data. The evidence fulfills the guidelines for sufficient evidence of an association (below) and satisfies several of the guidelines used to assess causality: strength of association, dose-response relationship, consistency of association, biologic plausibility, and a temporal relationship.

Sufficient Evidence of an Association

Evidence from available studies is sufficient to conclude that there is a positive association. A consistent positive association has been observed between exposure to a specific agent and a specific health outcome in human studies in which chance and bias, including confounding, could be ruled out with reasonable confidence. For example, several high-quality studies report consistent positive associations, and the studies are sufficiently free of bias, including adequate control for confounding.

Limited/Suggestive Evidence of an Association

Evidence from available studies suggests an association between exposure to a specific agent and a specific health outcome in human studies, but the body of evidence is limited. . . .

Inadequate/Insufficient Evidence to Determine Whether an Association Exists

Evidence from available studies is of insufficient quantity, quality, or consistency to permit a conclusion regarding the existence of an association between exposure to a specific agent and a specific health outcome in humans.

Limited/Suggestive Evidence of No Association

Evidence from well-conducted studies is consistent in not showing a positive association between exposure to a specific agent and a specific health outcome after exposure of any magnitude. . . .

Source: IOM (Institute of Medicine). 2003. Gulf War and Health, Vol. 2, Insecticides and Solvents. Washington, DC: National Academies Press.

BOX 2 Categorization of Health Outcomes^a Reviewed in Relation to TCE, PCE, or Solvent Mixtures

Sufficient Evidence of a Causal Relationship

- No outcomes

Sufficient Evidence of an Association

- No outcomes

Limited/Suggestive Evidence of an Association

- Kidney cancer
- Adult leukemia (solvent mixtures)
- Multiple myeloma (solvent mixtures)
- Myelodysplastic syndromes (solvent mixtures)
- Scleroderma (solvent mixtures)
- Neurobehavioral effects (solvent mixtures)

Inadequate/Insufficient Evidence to Determine Whether an Association Exists

- Oral/pharyngeal cancer
- Nasal cancer
- Laryngeal cancer
- Esophageal cancer (TCE)
- Stomach cancer
- Colon cancer
- Rectal cancer
- Pancreatic cancer
- Hepatobiliary cancer
- Lung cancer (TCE)
- Bone cancer
- Soft tissue sarcoma
- Melanoma
- Non-melanoma skin cancer
- Breast cancer (TCE)
- Cervical cancer
- Ovarian/uterine cancer
- Prostate cancer
- Bladder cancer (TCE)
- Cancer of the brain or central nervous system
- Non-Hodgkin lymphoma
- Hodgkin disease
- Multiple myeloma
- Adult leukemia
- Myelodysplastic syndromes
- Childhood leukemia
- Childhood neuroblastoma
- Childhood brain cancer
- Aplastic anemia
- Congenital malformations
- Male infertility
- Female infertility (after exposure cessation)
- Miscarriage, preterm birth, or fetal growth restriction (from maternal preconception exposure or paternal exposure)
- Preterm birth or fetal growth restriction (from exposure during pregnancy)
- Cardiovascular effects
- Liver function or risk of cirrhosis
- Gastrointestinal effects
- Renal toxicity
- Amyotrophic lateral sclerosis
- Parkinson disease
- Multiple sclerosis
- Alzheimer disease
- Long-term reduction in color discrimination
- Long-term hearing loss
- Long-term reduction in olfactory function

Limited/Suggestive Evidence of No Association

- No outcomes

^aOutcomes for TCE and PCE unless otherwise specified*

* PCE-only outcomes omitted